

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Phenylsemicarbazide

Uzma Ashiq,^{a*} Rifat Ara Jamal,^a Muhammad Nadeem Arshad,^b Zahida Tasneem Maqsood^a and Islam Ullah Khan^b

^aDepartment of Chemistry, University of Karachi, Karachi 75270, Pakistan, and^bDepartment of Chemistry, Government College University, Lahore, Pakistan

Correspondence e-mail: uzzmma@yahoo.com


Received 11 August 2009; accepted 1 September 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.039; wR factor = 0.107; data-to-parameter ratio = 15.7.

The title compound, $\text{C}_7\text{H}_9\text{N}_3\text{O}$, crystallizes with two independent molecules per asymmetric unit. The structure is stabilized by four distinct intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. Four intramolecular interactions of the $\text{N}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ types are also observed.

Related literature

For related structures see: Ashiq, Jamal *et al.* (2008, 2009); Jamal *et al.* (2008, 2009); Kallel *et al.* (1992); Saraogi *et al.* (2002); For the biological activity of hydrazides, see: Ara *et al.* (2007); Ashiq, Ara *et al.* (2008).



Experimental

Crystal data

 $\text{C}_7\text{H}_9\text{N}_3\text{O}$ $M_r = 151.17$ Monoclinic, $P2_1/c$ $a = 16.5984$ (10) Å $b = 8.8862$ (4) Å $c = 10.3518$ (6) Å $\beta = 91.359$ (3)° $V = 1526.43$ (14) Å³ $Z = 8$ Mo $K\alpha$ radiation $\mu = 0.09$ mm⁻¹ $T = 296$ K $0.43 \times 0.15 \times 0.12$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer

Absorption correction: multi-scan (SADABS; Bruker, 2005)

 $T_{\min} = 0.979$, $T_{\max} = 0.990$

15394 measured reflections

3500 independent reflections

2258 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.030$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.039$ $wR(F^2) = 0.107$ $S = 1.03$

3500 reflections

223 parameters

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.16$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.15$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N11}-\text{H11N}\cdots\text{N13}$	0.849 (16)	2.130 (15)	2.6149 (19)	116.0 (13)
$\text{N12}-\text{H12N}\cdots\text{O21}^{\text{i}}$	0.872 (15)	2.071 (15)	2.9386 (15)	172.8 (14)
$\text{N13}-\text{H14N}\cdots\text{O11}^{\text{ii}}$	0.887 (18)	2.383 (17)	3.2149 (18)	156.1 (15)
$\text{N21}-\text{H21N}\cdots\text{N23}$	0.852 (16)	2.130 (17)	2.6093 (19)	115.3 (14)
$\text{N22}-\text{H22N}\cdots\text{O11}^{\text{iii}}$	0.901 (16)	2.079 (17)	2.9784 (17)	175.9 (14)
$\text{N23}-\text{H23N}\cdots\text{O21}^{\text{iv}}$	0.919 (17)	2.203 (17)	3.0850 (18)	160.5 (15)
$\text{C12}-\text{H12}\cdots\text{O11}$	0.93	2.33	2.9119 (18)	120
$\text{C22}-\text{H22}\cdots\text{O21}$	0.93	2.46	2.9833 (18)	116

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (iii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXL97.

The authors thank the Higher Education Commission Pakistan for providing the diffractometer at GCU, Lahore, and Bana International for their support in collecting the crystallographic data.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2182).

References

- Ara, R., Ashiq, U., Mahroof-Tahir, M., Maqsood, Z. T., Khan, K. M., Lodhi, M. A. & Choudhary, M. I. (2007). *Chem. Biodivers.* **4**, 58–71.
- Ashiq, U., Ara, R., Mahroof-Tahir, M., Maqsood, Z. T., Khan, K. M., Khan, S. N., Siddiqui, H. & Choudhary, M. I. (2008). *Chem. Biodivers.* **5**, 82–92.
- Ashiq, U., Jamal, R. A., Mahroof-Tahir, M., Keramidas, A. D., Maqsood, Z. T., Khan, K. M. & Tahir, M. N. (2008). *Anal. Sci.* **X**, **24**, 103–104.
- Ashiq, U., Jamal, R. A., Tahir, M. N., Yousuf, S. & Khan, I. U. (2009). *Acta Cryst.* **E65**, o1551.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Jamal, R. A., Ashiq, U., Arshad, M. N., Maqsood, Z. T. & Khan, I. U. (2008). *Acta Cryst.* **E64**, o2188.
- Jamal, R. A., Ashiq, U., Arshad, M. N., Maqsood, Z. T. & Khan, I. U. (2009). *Acta Cryst.* **E65**, o1764.
- Kallel, A., Amor, B. H., Svoboda, I. & Fuess, H. (1992). *Z. Kristallogr.* **198**, 137–140.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Saraogi, I., Mruthyunjayaswamy, B. H. M., Ijare, O. B., Jadegoud, Y. & Guru Row, T. N. (2002). *Acta Cryst.* **E58**, o1341–o1342.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o2360 [doi:10.1107/S1600536809035284]

4-Phenylsemicarbazide

U. Ashiq, R. A. Jamal, M. N. Arshad, Z. T. Maqsood and I. U. Khan

Comment

Hydrazides are known to have different biological activities (Ashiq, Ara *et al.*, 2008; Ara *et al.*, 2007). In order to study the biological activity of 4-phenylsemicarbazide, we undertook the synthesis of title compound and report its crystal structure in this paper. The title compound was found to be active against urease enzyme (Ara *et al.*, 2007). The structures of benzhydrazide (Kallel *et al.*, 1992), *para*-chloro (Saraogi *et al.*, 2002), *para*-bromo (Ashiq, Jamal *et al.*, 2008), *para*-iodo (Jamal *et al.*, 2008), *para*-methoxy (Ashiq, Jamal *et al.*, 2009) and *para*-hydroxy (Jamal *et al.*, 2009) analogues have already been reported.

The unit cell contains two crystallographically unique molecules, whose molecular structures are presented in Fig. 1. The bond distances and bond angles are similar to the corresponding distances and angles reported in the structures quoted above. The molecular packing diagram (Fig. 2) shows the presence of four intermolecular hydrogen bonds and four intramolecular hydrogen interactions. In the crystal structure, two adjacent molecules are held together by intermolecular hydrogen bonds between the N12—H12N \cdots O21, N13—H14N \cdots O11, N22—H22N \cdots O11 and N23—H23N \cdots O21 (details are given in Table 1).

Experimental

All reagent-grade chemicals were obtained from Aldrich and Sigma Chemical companies and were used without further purification. To a solution of phenyl urea (34 g, 0.25 moles) in 50 ml ethanol, hydrazine hydrate (25.0 ml, 0.45 moles) was added. The mixture was refluxed for 24 h and solid 4-phenylsemicarbazide was obtained upon removal of the solvent by rotary evaporation (yield 79%).

Refinement

H atoms were positioned geometrically, with C—H = 0.93 Å for aromatic C-atoms, and for N1 and N2 atoms were taken from fourier synthesis and their coordinates were refined with N—H = 0.849 (16)–0.918 (17)Å with U_{iso} set to 1.2 U_{eq} of their parent atoms.

Figures




Fig. 1. ORTEP diagram of the title compound with the ellipsoids drawn at the 50% probability level, showing the atomic labels.




Fig. 2. A packing diagram viewed down the *c* axis showing hydrogen bonds drawn as dashed lines. Hydrogen atoms not involved in H-bonding have been omitted.

4-Phenylsemicarbazide

Crystal data

$C_7H_9N_3O$

$M_r = 151.17$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 16.5984$ (10) Å

$b = 8.8862$ (4) Å

$c = 10.3518$ (6) Å

$\beta = 91.359$ (3)°

$V = 1526.43$ (14) Å³

$Z = 8$

$F_{000} = 640$

$D_x = 1.316$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3838 reflections

$\theta = 2.5$ – 24.4 °

$\mu = 0.09$ mm⁻¹

$T = 296$ K

Needle, colourless

$0.43 \times 0.15 \times 0.12$ mm

Data collection

Bruker Kappa APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 296$ K

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2005)

$T_{\min} = 0.979$, $T_{\max} = 0.990$

15394 measured reflections

3500 independent reflections

2258 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 27.5$ °

$\theta_{\min} = 2.5$ °

$h = -21 \rightarrow 20$

$k = -6 \rightarrow 11$

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.107$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0459P)^2 + 0.1614P]$

where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.03$	$(\Delta/\sigma)_{\max} < 0.001$
3500 reflections	$\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$
223 parameters	$\Delta\rho_{\min} = -0.15 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O11	0.01604 (6)	0.79747 (12)	0.48141 (10)	0.0553 (4)
N11	0.09359 (7)	0.58992 (13)	0.44101 (12)	0.0427 (4)
N12	-0.01754 (7)	0.64373 (13)	0.31698 (11)	0.0438 (4)
N13	-0.00041 (8)	0.51513 (16)	0.24513 (14)	0.0506 (5)
C11	0.15892 (8)	0.60879 (14)	0.52831 (12)	0.0354 (4)
C12	0.16095 (9)	0.71248 (16)	0.62779 (13)	0.0439 (5)
C13	0.22909 (10)	0.72368 (18)	0.70707 (15)	0.0508 (5)
C14	0.29449 (9)	0.63342 (19)	0.68932 (15)	0.0554 (6)
C15	0.29205 (10)	0.52914 (19)	0.59133 (16)	0.0572 (6)
C16	0.22502 (9)	0.51619 (17)	0.51125 (14)	0.0462 (5)
C17	0.03011 (8)	0.68266 (15)	0.41816 (13)	0.0378 (5)
O21	0.17516 (6)	0.28317 (12)	0.23819 (10)	0.0529 (4)
N21	0.28009 (7)	0.40092 (15)	0.13966 (13)	0.0500 (5)
N22	0.15623 (8)	0.38933 (15)	0.04272 (12)	0.0488 (4)
N23	0.18846 (8)	0.47117 (17)	-0.05959 (13)	0.0514 (5)
C21	0.34464 (8)	0.38042 (16)	0.22790 (14)	0.0433 (5)
C22	0.34917 (9)	0.26384 (18)	0.31535 (15)	0.0509 (5)
C23	0.41651 (9)	0.2490 (2)	0.39537 (16)	0.0593 (6)
C24	0.47943 (10)	0.3476 (2)	0.38889 (17)	0.0643 (6)
C25	0.47515 (10)	0.4626 (2)	0.30239 (18)	0.0707 (7)
C26	0.40834 (9)	0.48014 (19)	0.22223 (17)	0.0596 (6)
C27	0.20285 (8)	0.35311 (15)	0.14604 (14)	0.0405 (5)
H11N	0.0960 (9)	0.5210 (18)	0.3845 (15)	0.0510*
H12	0.11680	0.77440	0.64140	0.0530*
H12N	-0.0624 (9)	0.6927 (17)	0.3025 (14)	0.0530*
H13	0.23030	0.79400	0.77360	0.0610*
H13N	-0.0413 (10)	0.4498 (19)	0.2569 (15)	0.0610*
H14	0.34000	0.64240	0.74280	0.0660*

supplementary materials

H14N	0.0017 (10)	0.5393 (19)	0.1621 (17)	0.0610*
H15	0.33610	0.46660	0.57900	0.0690*
H16	0.22410	0.44500	0.44540	0.0550*
H21N	0.2880 (10)	0.4551 (18)	0.0733 (16)	0.0600*
H22	0.30700	0.19530	0.32060	0.0610*
H22N	0.1043 (10)	0.3598 (17)	0.0395 (15)	0.0590*
H23	0.41900	0.17040	0.45470	0.0710*
H23N	0.1844 (10)	0.4143 (19)	-0.1336 (16)	0.0620*
H24	0.52460	0.33620	0.44290	0.0770*
H24N	0.1591 (10)	0.5550 (19)	-0.0679 (15)	0.0620*
H25	0.51780	0.53010	0.29720	0.0850*
H26	0.40610	0.55970	0.16390	0.0720*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O11	0.0570 (7)	0.0548 (7)	0.0534 (7)	0.0192 (5)	-0.0156 (5)	-0.0142 (5)
N11	0.0424 (7)	0.0419 (7)	0.0434 (7)	0.0079 (6)	-0.0090 (6)	-0.0063 (5)
N12	0.0394 (7)	0.0465 (7)	0.0449 (7)	0.0100 (6)	-0.0088 (6)	-0.0061 (6)
N13	0.0484 (8)	0.0510 (8)	0.0519 (8)	0.0034 (6)	-0.0074 (7)	-0.0105 (7)
C11	0.0355 (7)	0.0362 (7)	0.0345 (7)	-0.0003 (6)	0.0002 (6)	0.0061 (6)
C12	0.0433 (8)	0.0468 (8)	0.0416 (8)	0.0066 (7)	-0.0014 (6)	-0.0018 (7)
C13	0.0546 (10)	0.0539 (9)	0.0435 (9)	-0.0006 (8)	-0.0050 (7)	-0.0052 (7)
C14	0.0438 (9)	0.0679 (11)	0.0537 (10)	0.0002 (8)	-0.0136 (7)	0.0009 (8)
C15	0.0435 (9)	0.0658 (11)	0.0619 (10)	0.0155 (8)	-0.0063 (8)	-0.0005 (8)
C16	0.0438 (8)	0.0480 (9)	0.0464 (9)	0.0087 (7)	-0.0042 (7)	-0.0025 (7)
C17	0.0372 (8)	0.0392 (8)	0.0370 (8)	0.0024 (6)	-0.0014 (6)	0.0032 (6)
O21	0.0386 (6)	0.0680 (7)	0.0520 (7)	-0.0068 (5)	-0.0024 (5)	0.0135 (5)
N21	0.0380 (7)	0.0630 (9)	0.0488 (8)	-0.0090 (6)	-0.0040 (6)	0.0102 (6)
N22	0.0372 (7)	0.0595 (8)	0.0495 (8)	-0.0050 (6)	-0.0050 (6)	0.0095 (6)
N23	0.0517 (8)	0.0591 (9)	0.0432 (8)	0.0001 (6)	-0.0026 (6)	0.0044 (7)
C21	0.0340 (8)	0.0526 (9)	0.0433 (8)	-0.0017 (6)	0.0008 (6)	-0.0063 (7)
C22	0.0384 (8)	0.0566 (10)	0.0578 (10)	-0.0019 (7)	0.0009 (7)	0.0013 (8)
C23	0.0455 (10)	0.0743 (12)	0.0579 (10)	0.0094 (8)	-0.0022 (8)	0.0068 (9)
C24	0.0405 (9)	0.0906 (13)	0.0612 (11)	0.0034 (9)	-0.0121 (8)	-0.0091 (10)
C25	0.0445 (10)	0.0823 (13)	0.0846 (13)	-0.0177 (9)	-0.0116 (9)	-0.0025 (11)
C26	0.0488 (10)	0.0628 (11)	0.0669 (11)	-0.0135 (8)	-0.0044 (8)	0.0043 (8)
C27	0.0355 (8)	0.0419 (8)	0.0440 (8)	-0.0013 (6)	-0.0016 (6)	-0.0033 (7)

Geometric parameters (\AA , $^\circ$)

O11—C17	1.2375 (17)	C12—C13	1.385 (2)
O21—C27	1.2362 (17)	C13—C14	1.366 (2)
N11—C11	1.4051 (18)	C14—C15	1.374 (2)
N11—C17	1.3541 (18)	C15—C16	1.376 (2)
N12—N13	1.3964 (18)	C12—H12	0.9300
N12—C17	1.3427 (18)	C13—H13	0.9300
N11—H11N	0.849 (16)	C14—H14	0.9300
N12—H12N	0.872 (15)	C15—H15	0.9300

N13—H14N	0.887 (18)	C16—H16	0.9300
N13—H13N	0.904 (17)	C21—C22	1.377 (2)
N21—C21	1.4032 (19)	C21—C26	1.382 (2)
N21—C27	1.3537 (18)	C22—C23	1.382 (2)
N22—C27	1.3443 (19)	C23—C24	1.366 (2)
N22—N23	1.4013 (19)	C24—C25	1.360 (3)
N21—H21N	0.852 (16)	C25—C26	1.378 (2)
N22—H22N	0.901 (16)	C22—H22	0.9300
N23—H23N	0.919 (17)	C23—H23	0.9300
N23—H24N	0.893 (17)	C24—H24	0.9300
C11—C12	1.3817 (19)	C25—H25	0.9300
C11—C16	1.386 (2)	C26—H26	0.9300
C11—N11—C17	128.70 (12)	C13—C12—H12	120.00
N13—N12—C17	120.18 (12)	C14—C13—H13	119.00
C11—N11—H11N	118.8 (10)	C12—C13—H13	119.00
C17—N11—H11N	111.6 (10)	C15—C14—H14	120.00
C17—N12—H12N	119.4 (10)	C13—C14—H14	120.00
N13—N12—H12N	120.0 (10)	C16—C15—H15	120.00
N12—N13—H14N	109.3 (11)	C14—C15—H15	120.00
H13N—N13—H14N	109.5 (15)	C15—C16—H16	120.00
N12—N13—H13N	106.9 (11)	C11—C16—H16	120.00
C21—N21—C27	129.56 (13)	C22—C21—C26	118.72 (14)
N23—N22—C27	120.18 (12)	N21—C21—C22	123.69 (13)
C21—N21—H21N	117.9 (11)	N21—C21—C26	117.53 (13)
C27—N21—H21N	112.4 (11)	C21—C22—C23	119.79 (14)
C27—N22—H22N	119.5 (10)	C22—C23—C24	121.19 (16)
N23—N22—H22N	120.4 (10)	C23—C24—C25	119.10 (16)
H23N—N23—H24N	110.4 (15)	C24—C25—C26	120.69 (16)
N22—N23—H24N	106.8 (10)	C21—C26—C25	120.50 (16)
N22—N23—H23N	108.8 (11)	O21—C27—N21	124.42 (13)
C12—C11—C16	119.04 (13)	O21—C27—N22	121.12 (13)
N11—C11—C16	116.64 (12)	N21—C27—N22	114.46 (13)
N11—C11—C12	124.32 (12)	C21—C22—H22	120.00
C11—C12—C13	119.65 (14)	C23—C22—H22	120.00
C12—C13—C14	121.24 (14)	C22—C23—H23	119.00
C13—C14—C15	119.07 (15)	C24—C23—H23	119.00
C14—C15—C16	120.69 (15)	C23—C24—H24	120.00
C11—C16—C15	120.30 (14)	C25—C24—H24	120.00
N11—C17—N12	114.90 (12)	C24—C25—H25	120.00
O11—C17—N11	124.42 (13)	C26—C25—H25	120.00
O11—C17—N12	120.67 (12)	C21—C26—H26	120.00
C11—C12—H12	120.00	C25—C26—H26	120.00
C17—N11—C11—C12	14.9 (2)	C16—C11—C12—C13	1.0 (2)
C17—N11—C11—C16	-164.97 (14)	N11—C11—C16—C15	178.94 (13)
C11—N11—C17—O11	-6.6 (2)	C11—C12—C13—C14	-0.3 (2)
C11—N11—C17—N12	171.74 (13)	C12—C13—C14—C15	-0.4 (2)
N13—N12—C17—O11	179.06 (13)	C13—C14—C15—C16	0.5 (2)
N13—N12—C17—N11	0.66 (18)	C14—C15—C16—C11	0.2 (2)

supplementary materials

C21—N21—C27—O21	2.2 (2)	N21—C21—C22—C23	177.50 (14)
C27—N21—C21—C22	24.3 (2)	C26—C21—C22—C23	0.2 (2)
C27—N21—C21—C26	-158.35 (15)	N21—C21—C26—C25	-177.24 (15)
C21—N21—C27—N22	-178.53 (14)	C22—C21—C26—C25	0.2 (2)
N23—N22—C27—N21	-0.5 (2)	C21—C22—C23—C24	-0.5 (2)
N23—N22—C27—O21	178.77 (13)	C22—C23—C24—C25	0.3 (3)
N11—C11—C12—C13	-178.84 (13)	C23—C24—C25—C26	0.1 (3)
C12—C11—C16—C15	-0.9 (2)	C24—C25—C26—C21	-0.4 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N11—H11N \cdots N13	0.849 (16)	2.130 (15)	2.6149 (19)	116.0 (13)
N12—H12N \cdots O21 ⁱ	0.872 (15)	2.071 (15)	2.9386 (15)	172.8 (14)
N13—H14N \cdots O11 ⁱⁱ	0.887 (18)	2.383 (17)	3.2149 (18)	156.1 (15)
N21—H21N \cdots N23	0.852 (16)	2.130 (17)	2.6093 (19)	115.3 (14)
N22—H22N \cdots O11 ⁱⁱⁱ	0.901 (16)	2.079 (17)	2.9784 (17)	175.9 (14)
N23—H23N \cdots O21 ^{iv}	0.919 (17)	2.203 (17)	3.0850 (18)	160.5 (15)
C12—H12 \cdots O11	0.93	2.33	2.9119 (18)	120
C22—H22 \cdots O21	0.93	2.46	2.9833 (18)	116

Symmetry codes: (i) $-x, y+1/2, -z+1/2$; (ii) $x, -y+3/2, z-1/2$; (iii) $-x, y-1/2, -z+1/2$; (iv) $x, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

