

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 4-Methyl-2H-1,3-oxazine-2,6(3H)-dione

Damon Parrish, Fredrick Leuschner, Gretchen M. Rehberg and Margaret E. Kastner\*

Department of Chemistry, Bucknell University, Lewisburg, PA 17837, USA

Correspondence e-mail: kastner@bucknell.edu


Received 13 August 2009; accepted 28 August 2009

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.049;  $wR$  factor = 0.191; data-to-parameter ratio = 15.6.

In the title compound,  $\text{C}_5\text{H}_5\text{NO}_3$ , the planar (maximum deviation = 0.075 Å for the ring O atom) molecules form  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds in a zigzag chain ( $\text{C}-\text{O}\cdots\text{N}$  bond angle  $\simeq 140^\circ$ ) between glide-related molecules.

## Related literature

For synthetic background, see: Warren *et al.* (1975); Rehberg & Glass (1995). For related structures, see: Copley *et al.* (2005); Parrish, Leuschner *et al.* (2009); Parrish, Tivitmahaisoon *et al.* (2009).



## Experimental

## Crystal data

$\text{C}_5\text{H}_5\text{NO}_3$   
 $M_r = 127.1$   
 Monoclinic,  $P2_1/n$

$a = 7.254$  (3) Å  
 $b = 6.683$  (2) Å  
 $c = 11.689$  (5) Å

$\beta = 98.11$  (4) $^\circ$   
 $V = 561.0$  (4) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

$\mu = 0.13$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.46 \times 0.30 \times 0.10$  mm

## Data collection

Bruker R3/V diffractometer  
 Absorption correction: none  
 1410 measured reflections  
 1294 independent reflections  
 910 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.012$   
 3 standard reflections  
 every 97 reflections  
 intensity decay: none

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.191$   
 $S = 0.93$   
 1294 reflections

83 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.23$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.24$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N3}-\text{H3}\cdots\text{O6}^i$	0.86	2.02	2.877 (3)	173

Symmetry code: (i)  $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *XSCANS* (Bruker, 1996); cell refinement: *XSCANS* (Bruker, 1996); data reduction: *XSCANS* (Bruker, 1996); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors thank the National Science Foundation for grant No. ILI8951058.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2197).

## References

- Bruker (1996). *XSCANS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Copley, R. C. B., Deprez, L. S., Lewis, T. C. & Price, S. L. (2005). *CrystEngComm*, **7**, 421–428.  
 Parrish, D., Leuschner, F., Rehberg, G. M. & Kastner, M. E. (2009). *Acta Cryst.* **E65**, o2356.  
 Parrish, D., Tivitmahaisoon, P., Rehberg, G. M. & Kastner, M. E. (2009). *Acta Cryst.* **E65**, o2355.  
 Rehberg, G. M. & Glass, B. M. (1995). *Org. Prep. Proced. Int.* **27**, 651–652.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Warren, J. D., MacMillan, J. H. & Washburne, S. S. (1975). *J. Org. Chem.* **40**, 743–746.

**supplementary materials**

*Acta Cryst.* (2009). E65, o2354 [ doi:10.1107/S1600536809034618 ]

## 4-Methyl-2*H*-1,3-oxazine-2,6(3*H*)-dione

D. Parrish, F. Leuschner, G. M. Rehberg and M. E. Kastner

### Comment

The synthesis of derivatives of 3-oxauracil has previously been reported (Warren *et al.*, 1975) and an improved synthesis of the unsubstituted 3-oxauracil was reported by Rehberg & Glass (1995). The structure of the unsubstituted 3-oxauracil and its monohydrate have been reported (Copley *et al.*, 2005). Three derivatives of 3-oxauracil (4-methyl, 4-bromo, and 4,5-dichloro) have been prepared in our laboratory in route to the synthesis of 1-aza-1,3-butadienes. In this paper, we report the crystal structure of the title compound, (I).

In the title compound (Fig. 1) only one intermolecular H-bond is formed between N3 and O6 of glide-related molecules (details are given in Table 1). Although the molecules of (I) are planar, the H-bonding chains are staggered as shown in Figure 2. The hydrogen bonding networks in (I) differs significantly from the hydrogen bonding in 4,5-dichloro (Parrish, Leuschner *et al.*, 2009) and 4-bromo (Parrish, Tivitmahaisoon *et al.*, 2009) derivatives.

### Experimental

Citraconic anhydride (3-methylfuran-2,5-dione, 2.0 ml, 22 mmol) and trimethylsilyl azide (3.0 ml, 23 mmol) were added to 10 ml dichloromethane at 273 K and stirred under nitrogen for 4 h. Upon warming to room temperature over night, a white precipitate formed. Ethanol (2.5 ml) was added, the mixture stirred 2 additional hours, and then the solvent was removed under reduced pressure to obtain the title compound; yield: 1.7 g (13 mmol, 59%). Crystals of the title compound were grown from a solution of acetone at room temperature by slow evaporation.

### Refinement

Hydrogen positions were calculated and refined using a riding model using the following C—H distances: methyl 0.96 Å, methylene 0.93 Å, and N—H 0.88 Å with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C5/N3})$  and  $1.5U_{\text{eq}}(\text{C7})$ .

### Figures




Fig. 1. The molecular structure of the title compound, with atom labels and 50% probability displacement ellipsoids for non-H atoms.




Fig. 2. The packing of the title compound viewed down the *a* axis; intermolecular hydrogen bonds have been represented by dashed lines.

## 4-Methyl-2H-1,3-oxazine-2,6(3H)-dione

### Crystal data

$C_5H_5NO_3$

$M_r = 127.1$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 7.254$  (3) Å

$b = 6.683$  (2) Å

$c = 11.689$  (5) Å

$\beta = 98.11$  (4)°

$V = 561.0$  (4) Å<sup>3</sup>

$Z = 4$

$F_{000} = 264$

$D_x = 1.505$  Mg m<sup>-3</sup>

$D_m = 1.46$  Mg m<sup>-3</sup>

$D_m$  measured by floatation in bromoform/hexane solution

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 20 reflections

$\theta = 10$ – $12.5^\circ$

$\mu = 0.13$  mm<sup>-1</sup>

$T = 293$  K

Plates, colorless

$0.46 \times 0.30 \times 0.10$  mm

### Data collection

Bruker R3/V  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$  K

$\theta$  –  $2\theta$  scans

Absorption correction: none

1410 measured reflections

1294 independent reflections

910 reflections with  $I > 2\sigma(I)$

$R_{int} = 0.012$

$\theta_{max} = 27.6^\circ$

$\theta_{min} = 3.1^\circ$

$h = 0 \rightarrow 9$

$k = 0 \rightarrow 8$

$l = -15 \rightarrow 15$

3 standard reflections

every 97 reflections

intensity decay: none

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.049$

$wR(F^2) = 0.191$

$S = 0.93$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1301P)^2 + 0.1905P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{max} = 0.005$

1294 reflections  $\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$   
 83 parameters  $\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$   
 Primary atom site location: structure-invariant direct methods Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.6042 (2)	0.1383 (2)	0.68042 (13)	0.0466 (5)
C2	0.4998 (3)	0.2078 (3)	0.58306 (19)	0.0422 (5)
O2	0.3337 (2)	0.2161 (3)	0.57888 (19)	0.0694 (6)
N3	0.5939 (2)	0.2571 (3)	0.49490 (14)	0.0406 (5)
H3	0.5305	0.2967	0.4312	0.049*
C4	0.7833 (3)	0.2477 (3)	0.50089 (18)	0.0392 (5)
C5	0.8847 (3)	0.1896 (3)	0.59971 (19)	0.0430 (5)
H5	1.0137	0.1846	0.6048	0.052*
C6	0.7988 (3)	0.1357 (3)	0.69668 (18)	0.0435 (5)
O6	0.8682 (3)	0.0868 (3)	0.79242 (15)	0.0695 (7)
C7	0.8619 (4)	0.3002 (5)	0.3938 (2)	0.0630 (8)
H7A	0.9955	0.2979	0.4093	0.095*
H7B	0.8210	0.4317	0.3687	0.095*
H7C	0.8201	0.2050	0.3343	0.095*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0448 (9)	0.0555 (10)	0.0417 (8)	-0.0028 (7)	0.0137 (6)	0.0020 (7)
C2	0.0329 (10)	0.0455 (12)	0.0490 (11)	0.0003 (8)	0.0086 (8)	-0.0068 (9)
O2	0.0324 (9)	0.0847 (14)	0.0936 (15)	-0.0020 (9)	0.0174 (9)	-0.0026 (11)
N3	0.0342 (9)	0.0511 (10)	0.0350 (9)	0.0041 (7)	0.0003 (7)	0.0010 (7)
C4	0.0374 (10)	0.0384 (10)	0.0445 (11)	0.0029 (8)	0.0152 (8)	-0.0025 (9)
C5	0.0294 (9)	0.0458 (12)	0.0534 (12)	0.0012 (9)	0.0049 (8)	-0.0021 (10)
C6	0.0447 (11)	0.0395 (11)	0.0435 (11)	-0.0024 (9)	-0.0038 (9)	-0.0025 (9)
O6	0.0831 (14)	0.0687 (13)	0.0494 (10)	-0.0096 (10)	-0.0167 (9)	0.0110 (9)
C7	0.0651 (16)	0.0733 (18)	0.0573 (14)	0.0053 (13)	0.0313 (12)	0.0133 (13)

## supplementary materials

---

### Geometric parameters (Å, °)

O1—C2	1.357 (3)	C4—C7	1.489 (3)
O1—C6	1.398 (3)	C5—C6	1.415 (3)
C2—O2	1.200 (3)	C5—H5	0.9300
C2—N3	1.354 (3)	C6—O6	1.206 (3)
N3—C4	1.368 (3)	C7—H7A	0.9600
N3—H3	0.8600	C7—H7B	0.9600
C4—C5	1.337 (3)	C7—H7C	0.9600
C2—O1—C6	123.50 (17)	C4—C5—H5	119.5
O2—C2—N3	124.6 (2)	C6—C5—H5	119.5
O2—C2—O1	119.2 (2)	O6—C6—O1	114.3 (2)
N3—C2—O1	116.10 (18)	O6—C6—C5	129.7 (2)
C2—N3—C4	124.08 (18)	O1—C6—C5	115.99 (18)
C2—N3—H3	118.0	C4—C7—H7A	109.5
C4—N3—H3	118.0	C4—C7—H7B	109.5
C5—C4—N3	118.95 (18)	H7A—C7—H7B	109.5
C5—C4—C7	124.5 (2)	C4—C7—H7C	109.5
N3—C4—C7	116.6 (2)	H7A—C7—H7C	109.5
C4—C5—C6	121.03 (19)	H7B—C7—H7C	109.5
C6—O1—C2—O2	-175.5 (2)	N3—C4—C5—C6	0.9 (3)
C6—O1—C2—N3	6.7 (3)	C7—C4—C5—C6	-177.8 (2)
O2—C2—N3—C4	179.7 (2)	C2—O1—C6—O6	173.2 (2)
O1—C2—N3—C4	-2.6 (3)	C2—O1—C6—C5	-6.9 (3)
C2—N3—C4—C5	-1.1 (3)	C4—C5—C6—O6	-177.3 (2)
C2—N3—C4—C7	177.7 (2)	C4—C5—C6—O1	2.8 (3)

### Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N3—H3 $\cdots$ O6 <sup>i</sup>	0.86	2.02	2.877 (3)	173

Symmetry codes: (i)  $x-1/2, -y+1/2, z-1/2$ .

Fig. 1




Fig. 2

