

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

10-Allyl-2,3-dihydro-1H-pyrrolo[2,1-c]-[1,4]benzodiazepine-5,11(10H,11aH)-dione

Hanane Benzeid,^a El Mokhtar Essassi,^a Nathalie Saffon,^b Bernard Garrigues^c and Seik Weng Ng^{d*}

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, ^bService commun Rayons X, Université Paul Sabatier, Bâtiment 2R1, 118 route de Narbonne, 31062 Toulouse, France, ^cHétérochimie Fondamentale et Appliquée, Université Paul Sabatier, UMR 5069, 118 Route de Narbonne, 31062 Toulouse, France, and ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 24 August 2009; accepted 27 August 2009

Key indicators: single-crystal X-ray study; $T = 193$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.049; wR factor = 0.146; data-to-parameter ratio = 12.7.

The compound, $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_2$, features a pyrroline ring fused with a seven-membered diazepine ring; the latter system adopts a boat conformation (with the methine C atom as the prow and the two C atoms of the aromatic ring as the stern). A CH_2-CH_2 segment of the pyrroline ring is disordered over two positions in a 1:1 ratio.

Related literature

Pyrrolo[2,1-c][1,4]benzodiazepines are potent, naturally occurring antitumor antibiotics produced by *Streptomyces* species; see: Cargill *et al.* (1974); Thurston *et al.* (1993). For the design and synthesis of DNA inter-strand cross-linking as well as conjugate agents to enhance the sequence selectivity and to increase selectivity for tumor cells, see: Bose *et al.* (1992); Gregson *et al.* (2004).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_2$
 $M_r = 256.30$
 Orthorhombic, $P2_12_12_1$
 $a = 7.0988$ (1) Å
 $b = 11.7166$ (2) Å
 $c = 15.6592$ (3) Å
 $V = 1302.44$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 193$ K
 $0.30 \times 0.30 \times 0.20$ mm

Data collection

Bruker APEXII diffractometer
 Absorption correction: none
 20329 measured reflections
 2263 independent reflections
 1900 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.146$
 $S = 1.04$
 2263 reflections
 178 parameters
 15 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.53$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.30$ e Å⁻³

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

We thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2637).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bose, D. S., Thompson, A. S., Ching, J. A., Hartley, J. A., Berardini, M. D., Jenkins, T. C., Neidle, S., Hurley, L. H. & Thurston, D. E. (1992). *J. Am. Chem. Soc.* **114**, 4939–4941.
 Bruker (2005). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cargill, C., Bachmann, E. & Zbinden, G. (1974). *J. Natl. Cancer Inst.* **53**, 481–486.
 Gregson, S. T., Howard, P. W., Gullick, D. R., Hamaguchi, A., Corcoran, K. E., Brooks, N. A., Hartley, J. A., Jenkins, T. C., Patel, S., Guille, M. J. & Thurston, D. E. (2004). *J. Med. Chem.* **47**, 1161–1174.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Thurston, D. E., Neidle, S. & Waring, M. J. (1993). *Molecular Aspects of Anticancer Drug-DNA Interactions*, Vol. 1, pp. 54–88. New York: Macmillan Press.
 Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, o2322 [doi:10.1107/S1600536809034266]

10-Allyl-2,3-dihydro-1*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5,11(10*H*,11*aH*)-dione

H. Benzeid, E. M. Essassi, N. Saffon, B. Garrigues and S. W. Ng

Experimental

2,3-Dihydro-1*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5,11(10*H*,11*aH*)-dione (1 g, 4.6 mmol), allyl bromide (0.64 g, 4.6 mmol) and potassium carbonate (0.64 g, 4.6 mmol) along with a catalytic amount of tetra-*n*-butyammonium bromide were stirred in *N,N*-dimethylformamide (20 ml) for 12 h. After the completion of the reaction (as monitored by TLC), the solid material was removed by filtration and the solvent evaporated under vacuum. Dichloromethane (20 ml) was added and the solution filtered. The solvent was removed and the product purified by recrystallization from dichloromethane to afford colorless crystals in 80% yield. The formulation was established by proton and carbon-13 NMR spectroscopy in CDCl₃.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95 to 0.99 Å) and were included in the refinement in the riding model approximation, with *U*(H) set to 1.2*U*(C).

Two of the carbon atoms in the tetrahydropyrrolyl ring are disordered over two positions; the occupancy could not be refined, and was assumed to be 50:50. The pairs of carbon-carbon (C10—C11, C10'—C11') distances were restrained to within 0.01 Å of each other, and the temperature factors of the primed atoms were restrained to those of the unprimed ones. Their anisotropic temperature factors were restrained to nearly isotropic values.

In the absence of significant anomalous dispersion effects, Friedel pairs were merged.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of C₁₅H₁₆N₂O₂ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

\ 10-Allyl-2,3-dihydro-1*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5,\ 11(10*H*,11*aH*)-dione

Crystal data

C₁₅H₁₆N₂O₂

M_r = 256.30

*F*₀₀₀ = 544

D_x = 1.307 Mg m⁻³

supplementary materials

Orthorhombic, $P2_12_12_1$
Hall symbol: P 2ac 2ab
 $a = 7.0988$ (1) Å
 $b = 11.7166$ (2) Å
 $c = 15.6592$ (3) Å
 $V = 1302.44$ (4) Å³
 $Z = 4$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 7433 reflections
 $\theta = 2.2$ – 25.3°
 $\mu = 0.09$ mm⁻¹
 $T = 193$ K
Block, colorless
 $0.30 \times 0.30 \times 0.20$ mm

Data collection

Bruker APEXII
diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 193$ K
 φ and ω scans
Absorption correction: None
20329 measured reflections
2263 independent reflections

1900 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$
 $\theta_{\text{max}} = 30.5^\circ$
 $\theta_{\text{min}} = 5.2^\circ$
 $h = -10 \rightarrow 10$
 $k = -16 \rightarrow 14$
 $l = -22 \rightarrow 19$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.146$
 $S = 1.04$
2263 reflections
178 parameters
15 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.086P)^2 + 0.2922P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.53$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.30$ e Å⁻³
Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.3349 (3)	0.89582 (17)	0.62000 (12)	0.0545 (5)	
O2	-0.1387 (3)	0.62142 (13)	0.76109 (11)	0.0423 (4)	
N1	0.2785 (2)	0.86934 (15)	0.76059 (12)	0.0312 (4)	
N2	-0.0624 (3)	0.77145 (15)	0.67775 (12)	0.0327 (4)	
C1	0.2213 (4)	0.88657 (18)	0.67841 (15)	0.0355 (5)	
C2	-0.0695 (3)	0.71719 (17)	0.75353 (14)	0.0301 (4)	
C3	-0.0008 (3)	0.78280 (17)	0.82960 (13)	0.0280 (4)	
C4	0.1542 (3)	0.85792 (16)	0.83096 (13)	0.0279 (4)	
C5	0.1940 (3)	0.91758 (17)	0.90646 (13)	0.0333 (4)	

H5	0.2989	0.9678	0.9083	0.040*	
C6	0.0828 (4)	0.9044 (2)	0.97834 (14)	0.0396 (5)	
H6	0.1090	0.9476	1.0283	0.048*	
C7	-0.0675 (4)	0.8278 (2)	0.97772 (14)	0.0407 (5)	
H7	-0.1425	0.8172	1.0274	0.049*	
C8	-0.1059 (3)	0.76768 (19)	0.90409 (14)	0.0359 (5)	
H8	-0.2069	0.7144	0.9040	0.043*	
C9	-0.1524 (4)	0.7262 (2)	0.60032 (14)	0.0399 (5)	
H9A	-0.2678	0.6823	0.6137	0.048*	0.50
H9B	-0.0651	0.6780	0.5667	0.048*	0.50
H9C	-0.2759	0.6916	0.6143	0.048*	0.50
H9D	-0.0719	0.6671	0.5736	0.048*	0.50
C10	-0.197 (3)	0.8367 (11)	0.5551 (12)	0.059 (3)	0.50
H10A	-0.2187	0.8236	0.4935	0.071*	0.50
H10B	-0.3102	0.8732	0.5800	0.071*	0.50
C11	-0.025 (2)	0.9097 (17)	0.5692 (5)	0.047 (2)	0.50
H11A	-0.0500	0.9912	0.5570	0.056*	0.50
H11B	0.0832	0.8835	0.5342	0.056*	0.50
C10'	-0.178 (3)	0.8258 (11)	0.5408 (12)	0.059 (3)	0.50
H10C	-0.3124	0.8465	0.5366	0.071*	0.50
H10D	-0.1312	0.8061	0.4830	0.071*	0.50
C11'	-0.066 (2)	0.9241 (16)	0.5769 (5)	0.047 (2)	0.50
H11C	-0.1475	0.9923	0.5823	0.056*	0.50
H11D	0.0400	0.9430	0.5384	0.056*	0.50
C12	0.0096 (3)	0.88869 (18)	0.66551 (14)	0.0339 (5)	
H12	-0.0551	0.9459	0.7027	0.041*	0.50
H12'	-0.0479	0.9394	0.7098	0.041*	0.50
C13	0.4837 (3)	0.8775 (2)	0.77691 (19)	0.0433 (6)	
H13A	0.5522	0.8322	0.7335	0.052*	
H13B	0.5121	0.8446	0.8337	0.052*	
C14	0.5520 (4)	1.0005 (2)	0.7740 (2)	0.0490 (6)	
H14	0.5020	1.0490	0.7309	0.059*	
C15	0.6751 (4)	1.0438 (3)	0.8269 (2)	0.0565 (7)	
H15A	0.7278	0.9976	0.8707	0.068*	
H15B	0.7119	1.1214	0.8216	0.068*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0621 (12)	0.0550 (11)	0.0466 (10)	-0.0140 (10)	0.0231 (9)	-0.0031 (9)
O2	0.0473 (9)	0.0272 (7)	0.0525 (10)	-0.0097 (7)	-0.0042 (8)	0.0039 (7)
N1	0.0253 (7)	0.0269 (8)	0.0413 (10)	-0.0014 (6)	0.0036 (7)	0.0012 (7)
N2	0.0361 (8)	0.0276 (8)	0.0345 (8)	-0.0072 (7)	-0.0013 (8)	-0.0020 (7)
C1	0.0425 (11)	0.0278 (9)	0.0362 (10)	-0.0075 (9)	0.0072 (9)	-0.0012 (8)
C2	0.0273 (8)	0.0252 (8)	0.0380 (10)	-0.0003 (7)	0.0003 (9)	0.0007 (8)
C3	0.0266 (8)	0.0245 (8)	0.0331 (9)	0.0012 (7)	-0.0001 (8)	0.0042 (7)
C4	0.0267 (8)	0.0225 (8)	0.0345 (9)	0.0010 (7)	-0.0026 (8)	0.0043 (7)
C5	0.0358 (10)	0.0277 (9)	0.0362 (10)	-0.0004 (8)	-0.0083 (9)	0.0034 (8)

supplementary materials

C6	0.0487 (13)	0.0388 (11)	0.0312 (10)	0.0040 (10)	-0.0062 (10)	0.0022 (9)
C7	0.0467 (12)	0.0436 (12)	0.0320 (10)	0.0029 (11)	0.0042 (10)	0.0098 (9)
C8	0.0345 (10)	0.0360 (10)	0.0372 (10)	-0.0027 (9)	0.0015 (9)	0.0095 (9)
C9	0.0446 (12)	0.0394 (11)	0.0357 (11)	-0.0074 (10)	-0.0010 (10)	-0.0097 (9)
C10	0.088 (4)	0.059 (3)	0.031 (5)	-0.015 (3)	-0.015 (4)	0.000 (3)
C11	0.061 (6)	0.042 (4)	0.0371 (16)	-0.008 (4)	-0.012 (3)	0.010 (2)
C10'	0.088 (4)	0.059 (3)	0.031 (5)	-0.015 (3)	-0.015 (4)	0.000 (3)
C11'	0.061 (6)	0.042 (4)	0.0371 (16)	-0.008 (4)	-0.012 (3)	0.010 (2)
C12	0.0440 (11)	0.0259 (9)	0.0319 (10)	-0.0067 (9)	-0.0066 (9)	0.0041 (8)
C13	0.0253 (9)	0.0403 (12)	0.0643 (16)	0.0012 (9)	0.0044 (10)	-0.0007 (11)
C14	0.0315 (11)	0.0506 (14)	0.0648 (16)	-0.0065 (11)	0.0037 (12)	0.0040 (13)
C15	0.0413 (13)	0.0502 (14)	0.078 (2)	-0.0061 (12)	0.0002 (15)	-0.0023 (14)

Geometric parameters (Å, °)

O1—C1	1.224 (3)	C9—H9C	0.9900
O2—C2	1.231 (3)	C9—H9D	0.9900
N1—C1	1.365 (3)	C10—C11	1.509 (9)
N1—C4	1.418 (3)	C10—H10A	0.9900
N1—C13	1.482 (3)	C10—H10B	0.9900
N2—C2	1.347 (3)	C11—C12	1.547 (6)
N2—C9	1.470 (3)	C11—H11A	0.9900
N2—C12	1.478 (3)	C11—H11B	0.9900
C1—C12	1.517 (4)	C10'—C11'	1.507 (9)
C2—C3	1.499 (3)	C10'—H10C	0.9900
C3—C8	1.396 (3)	C10'—H10D	0.9900
C3—C4	1.409 (3)	C11'—C12	1.545 (6)
C4—C5	1.402 (3)	C11'—H11C	0.9900
C5—C6	1.383 (3)	C11'—H11D	0.9900
C5—H5	0.9500	C12—H12	1.0000
C6—C7	1.395 (4)	C12—H12'	1.0000
C6—H6	0.9500	C13—C14	1.522 (3)
C7—C8	1.378 (3)	C13—H13A	0.9900
C7—H7	0.9500	C13—H13B	0.9900
C8—H8	0.9500	C14—C15	1.306 (4)
C9—C10'	1.504 (7)	C14—H14	0.9500
C9—C10	1.510 (7)	C15—H15A	0.9500
C9—H9A	0.9900	C15—H15B	0.9500
C9—H9B	0.9900		
C1—N1—C4	124.17 (18)	C11—C10—H10A	110.9
C1—N1—C13	116.5 (2)	C9—C10—H10A	110.9
C4—N1—C13	118.9 (2)	C11—C10—H10B	110.9
C2—N2—C9	122.70 (18)	C9—C10—H10B	110.9
C2—N2—C12	124.44 (18)	H10A—C10—H10B	108.9
C9—N2—C12	112.27 (18)	C10—C11—C12	100.3 (12)
O1—C1—N1	121.5 (2)	C10—C11—H11A	111.7
O1—C1—C12	123.5 (2)	C12—C11—H11A	111.7
N1—C1—C12	115.02 (19)	C10—C11—H11B	111.7
O2—C2—N2	122.0 (2)	C12—C11—H11B	111.7

O2—C2—C3	121.4 (2)	H11A—C11—H11B	109.5
N2—C2—C3	116.48 (18)	C9—C10'—C11'	107.4 (12)
C8—C3—C4	118.96 (19)	C9—C10'—H10C	110.2
C8—C3—C2	115.13 (18)	C11'—C10'—H10C	110.2
C4—C3—C2	125.89 (18)	C9—C10'—H10D	110.2
C5—C4—C3	118.78 (19)	C11'—C10'—H10D	110.2
C5—C4—N1	118.87 (18)	H10C—C10'—H10D	108.5
C3—C4—N1	122.21 (18)	C10'—C11'—C12	108.3 (12)
C6—C5—C4	121.0 (2)	C10'—C11'—H11C	110.0
C6—C5—H5	119.5	C12—C11'—H11C	110.0
C4—C5—H5	119.5	C10'—C11'—H11D	110.0
C5—C6—C7	120.2 (2)	C12—C11'—H11D	110.0
C5—C6—H6	119.9	H11C—C11'—H11D	108.4
C7—C6—H6	119.9	N2—C12—C1	108.08 (19)
C8—C7—C6	119.1 (2)	N2—C12—C11'	104.2 (8)
C8—C7—H7	120.5	C1—C12—C11'	117.9 (7)
C6—C7—H7	120.5	N2—C12—C11	102.7 (8)
C7—C8—C3	121.9 (2)	C1—C12—C11	106.7 (6)
C7—C8—H8	119.1	N2—C12—H12	112.9
C3—C8—H8	119.1	C1—C12—H12	112.9
N2—C9—C10'	106.4 (8)	C11'—C12—H12	100.6
N2—C9—C10	99.7 (8)	C11—C12—H12	112.9
N2—C9—H9A	111.8	N2—C12—H12'	108.8
C10'—C9—H9A	115.9	C1—C12—H12'	108.8
C10—C9—H9A	111.8	C11'—C12—H12'	108.8
N2—C9—H9B	111.8	C11—C12—H12'	121.1
C10'—C9—H9B	100.8	N1—C13—C14	111.7 (2)
C10—C9—H9B	111.8	N1—C13—H13A	109.3
H9A—C9—H9B	109.6	C14—C13—H13A	109.3
N2—C9—H9C	110.5	N1—C13—H13B	109.3
C10'—C9—H9C	110.5	C14—C13—H13B	109.3
C10—C9—H9C	105.6	H13A—C13—H13B	108.0
H9B—C9—H9C	116.1	C15—C14—C13	124.2 (3)
N2—C9—H9D	110.5	C15—C14—H14	117.9
C10'—C9—H9D	110.5	C13—C14—H14	117.9
C10—C9—H9D	121.5	C14—C15—H15A	120.0
H9A—C9—H9D	101.8	C14—C15—H15B	120.0
H9C—C9—H9D	108.6	H15A—C15—H15B	120.0
C11—C10—C9	104.3 (14)		
C4—N1—C1—O1	-180.0 (2)	C2—N2—C9—C10	150.1 (10)
C13—N1—C1—O1	7.7 (3)	C12—N2—C9—C10	-21.4 (10)
C4—N1—C1—C12	-2.2 (3)	N2—C9—C10—C11	41.2 (14)
C13—N1—C1—C12	-174.50 (19)	C10'—C9—C10—C11	-87 (8)
C9—N2—C2—O2	6.1 (3)	C9—C10—C11—C12	-45.3 (16)
C12—N2—C2—O2	176.6 (2)	N2—C9—C10'—C11'	11.4 (17)
C9—N2—C2—C3	-169.74 (19)	C10—C9—C10'—C11'	65 (8)
C12—N2—C2—C3	0.8 (3)	C9—C10'—C11'—C12	-6.8 (19)
O2—C2—C3—C8	-35.3 (3)	C2—N2—C12—C1	70.3 (3)
N2—C2—C3—C8	140.6 (2)	C9—N2—C12—C1	-118.3 (2)

supplementary materials

O2—C2—C3—C4	146.2 (2)	C2—N2—C12—C11'	-163.4 (7)
N2—C2—C3—C4	-38.0 (3)	C9—N2—C12—C11'	7.9 (7)
C8—C3—C4—C5	-1.9 (3)	C2—N2—C12—C11	-177.1 (6)
C2—C3—C4—C5	176.60 (19)	C9—N2—C12—C11	-5.7 (6)
C8—C3—C4—N1	173.72 (18)	O1—C1—C12—N2	107.3 (3)
C2—C3—C4—N1	-7.8 (3)	N1—C1—C12—N2	-70.5 (3)
C1—N1—C4—C5	-134.8 (2)	O1—C1—C12—C11'	-10.4 (9)
C13—N1—C4—C5	37.4 (3)	N1—C1—C12—C11'	171.8 (8)
C1—N1—C4—C3	49.6 (3)	O1—C1—C12—C11	-2.6 (8)
C13—N1—C4—C3	-138.2 (2)	N1—C1—C12—C11	179.7 (8)
C3—C4—C5—C6	-0.7 (3)	C10'—C11'—C12—N2	-0.5 (14)
N1—C4—C5—C6	-176.43 (19)	C10'—C11'—C12—C1	119.3 (11)
C4—C5—C6—C7	2.4 (3)	C10'—C11'—C12—C11	85 (7)
C5—C6—C7—C8	-1.4 (4)	C10—C11—C12—N2	30.4 (12)
C6—C7—C8—C3	-1.2 (4)	C10—C11—C12—C1	144.0 (10)
C4—C3—C8—C7	2.9 (3)	C10—C11—C12—C11'	-68 (6)
C2—C3—C8—C7	-175.8 (2)	C1—N1—C13—C14	73.4 (3)
C2—N2—C9—C10'	159.3 (10)	C4—N1—C13—C14	-99.4 (3)
C12—N2—C9—C10'	-12.3 (10)	N1—C13—C14—C15	139.5 (3)

Fig. 1

