

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[trimethyltin(IV)]- μ -2-(2-chlorophenyl)acetato]

Liyuan Wen, Handong Yin* and Wenkuan Li

College of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China

Correspondence e-mail: handongyin@163.com


Received 13 September 2009; accepted 25 September 2009

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.014$ Å; R factor = 0.047; wR factor = 0.107; data-to-parameter ratio = 17.4.

In the title polymeric coordination compound, $[\text{Sn}(\text{CH}_3)_3(\text{C}_8\text{H}_6\text{ClO}_2)]_n$, the Sn atoms exhibit a distorted trigonal-bipyramidal geometry with the carboxylate O atoms of the 2-chlorophenylacetato ligands in axial positions and with the equatorial sites occupied by the three methyl groups. Adjacent Sn atoms are bridged by coordination to the two O atoms of each 2-chlorophenylacetato ligand, forming a chain structure.

Related literature

For the biological activity of organotin compounds, see: Wang *et al.* (2007). For related structures, see: Wang *et al.* (2007); Ma *et al.* (2006).



Experimental

Crystal data

 $[\text{Sn}(\text{CH}_3)_3(\text{C}_8\text{H}_6\text{ClO}_2)]_n$
 $M_r = 333.39$

 Monoclinic, $P2_1/n$
 $a = 7.0754$ (9) Å

 $b = 28.306$ (3) Å

 $c = 13.6721$ (15) Å

 $\beta = 93.117$ (2)°

 $V = 2734.1$ (5) Å³
 $Z = 8$

 Mo $K\alpha$ radiation

 $\mu = 2.05$ mm⁻¹
 $T = 298$ K

 $0.49 \times 0.32 \times 0.15$ mm

Data collection

Siemens SMART CCD area-

detector diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.434$, $T_{\max} = 0.749$

14063 measured reflections

4820 independent reflections

 3411 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.056$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.107$
 $S = 1.05$

4820 reflections

277 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 1.09$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.92$ e Å⁻³

Table 1

Selected bond lengths (Å).

Sn1—C17	2.107 (7)	Sn2—C22	2.102 (7)
Sn1—C18	2.117 (7)	Sn2—C20	2.113 (7)
Sn1—C19	2.120 (7)	Sn2—C21	2.117 (7)
Sn1—O3	2.194 (4)	Sn2—O2	2.207 (5)
Sn1—O1	2.396 (5)	Sn2—O4 ⁱ	2.432 (5)

 Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We acknowledge the National Natural Science Foundation of China (20771053) and the Natural Science Foundation of Shandong Province (Y2008B48) for financial support. This work was also supported by the Shangdong 'Tai-Shan Scholar Research Fund'.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2651).

References

- Ma, C., Li, J., Zhang, R. & Wang, D. (2006). *J. Organomet. Chem.* **691**, 1713–1721.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). SMART and SAINT. Siemens Analytical X-ray Systems, Inc., Madison, Wisconsin, USA.
- Wang, H., Yin, H. & Wang, D. (2007). *Acta Cryst.* **E63**, m2955.

supplementary materials

Acta Cryst. (2009). E65, m1261 [doi:10.1107/S1600536809038872]

catena-Poly[[trimethyltin(IV)]- μ -2-(2-chlorophenyl)acetato]

L. Wen, H. Yin and W. Li

Comment

The chemistry of organotin(IV) derivatives is a subject of study with growing interest due to their significant antimicrobial properties as well as antitumor activities (Wang *et al.*, 2007). As a part of our ongoing investigations in this field we have synthesized the title compound and present its crystal structure here. The title compound, which is shown in Fig.1 forms an extended one-dimensional chain structure arising from Sn—O bridges formed by the 2-(2-chlorophenyl)acetato ligands. The Sn—O bond distances in the compound (Sn(1)—O(1) = 2.396 (5) Å; Sn(1)—O(3) = 2.194 (4) Å) are comparable to those found in related organotin carboxylates (Ma *et al.*, 2006). The Sn atom assumes a slightly distorted trigonal-bipyramidal coordination geometry, provided by and three methyl groups in the equatorial positions and two O atoms of symmetry related carboxylate groups in the axial positions.

Experimental

The reaction was carried out under a nitrogen atmosphere. 2-(2-chlorophenyl)acetic acid (1 mmol) and sodium ethoxide (1.2 mmol) were added to a stirred solution of benzene (30 ml) in a Schlenk flask and stirred for 0.5 h. Trimethyltin chloride (1 mmol) was then added to the reactor and the reaction mixture was stirred for 12 h at room temperature. The resulting clear solution was evaporated under vacuum. The product was crystallized from a solution of dichloromethane/methanol (1:1) to yield colourless blocks of the title compound (yield 81%. m.p.390 K). Anal. Calcd (%) for C₁₁H₁₅Cl₁O₂Sn₁ (Mr = 333.37): C,39.63; H, 4.54; Cl, 10.63. Found (%): C, 39.51; H, 4.64; Cl, 10.75.

Refinement

The H atoms were positioned geometrically, with methyl C—H distances of 0.96 Å and aromatic C—H distances of 0.93 Å, and refined as riding on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{O})$ or $1.5 U_{\text{eq}}(\text{C})$ for the methyl groups.

Figures




Fig. 1. The asymmetric unit of the title compound, showing 50% probability displacement ellipsoids. H atoms have been omitted for clarity.




Fig. 2. A view of the one-dimensional extended chain structure in the title compound.

catena-Poly[[trimethyltin(IV)]- μ -2-(2-chlorophenyl)acetato]

Crystal data

[Sn(CH₃)₃(C₈H₆ClO₂)]

$M_r = 333.39$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 7.0754$ (9) Å

$b = 28.306$ (3) Å

$c = 13.6721$ (15) Å

$\beta = 93.117$ (2)°

$V = 2734.1$ (5) Å³

$Z = 8$

$F_{000} = 1312$

$D_x = 1.620$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4326 reflections

$\theta = 2.6$ – 24.8 °

$\mu = 2.05$ mm⁻¹

$T = 298$ K

Block, colourless

$0.49 \times 0.32 \times 0.15$ mm

Data collection

Siemens SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298$ K

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.434$, $T_{\max} = 0.749$

14063 measured reflections

4820 independent reflections

3411 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.056$

$\theta_{\text{max}} = 25.0$ °

$\theta_{\text{min}} = 1.7$ °

$h = -8 \rightarrow 8$

$k = -33 \rightarrow 33$

$l = -10 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.047$

$wR(F^2) = 0.107$

$S = 1.05$

4820 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0224P)^2 + 9.0463P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.001$

$\Delta\rho_{\text{max}} = 1.09$ e Å⁻³

277 parameters

$$\Delta\rho_{\min} = -0.92 \text{ e } \text{\AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.02624 (7)	0.138217 (16)	0.88059 (3)	0.04764 (16)
Sn2	0.20583 (7)	0.212967 (17)	1.21572 (3)	0.04848 (16)
Cl1	0.4073 (6)	0.03321 (16)	1.1310 (3)	0.1655 (15)
Cl2	-0.2267 (4)	0.14040 (10)	0.36127 (15)	0.0943 (8)
O1	0.1070 (7)	0.14009 (16)	1.0530 (3)	0.0583 (13)
O2	0.1197 (7)	0.13802 (16)	1.2145 (3)	0.0572 (13)
O3	-0.0431 (7)	0.13733 (16)	0.7223 (3)	0.0564 (13)
O4	-0.1907 (7)	0.20535 (18)	0.7318 (3)	0.0620 (14)
C1	0.0842 (11)	0.1192 (3)	1.1319 (5)	0.0553 (19)
C2	0.0044 (13)	0.0697 (3)	1.1322 (6)	0.073 (2)
H2A	-0.1323	0.0720	1.1333	0.088*
H2B	0.0328	0.0546	1.0710	0.088*
C3	0.0721 (17)	0.0382 (3)	1.2138 (7)	0.079 (3)
C4	0.2486 (19)	0.0191 (4)	1.2200 (8)	0.103 (3)
C5	0.316 (2)	-0.0112 (4)	1.2941 (10)	0.127 (4)
H5	0.4382	-0.0237	1.2975	0.153*
C6	0.184 (3)	-0.0208 (4)	1.3620 (10)	0.121 (5)
H6	0.2224	-0.0410	1.4131	0.145*
C7	0.001 (2)	-0.0036 (4)	1.3623 (9)	0.115 (4)
H7	-0.0816	-0.0121	1.4100	0.138*
C8	-0.0504 (18)	0.0264 (3)	1.2880 (8)	0.100 (3)
H8	-0.1708	0.0397	1.2861	0.120*
C9	-0.1339 (10)	0.1720 (3)	0.6832 (5)	0.0504 (18)
C10	-0.1744 (10)	0.1690 (3)	0.5734 (5)	0.0553 (19)
H10A	-0.0690	0.1534	0.5443	0.066*
H10B	-0.1837	0.2007	0.5469	0.066*
C11	-0.3557 (10)	0.1423 (2)	0.5447 (5)	0.0514 (18)
C12	-0.3923 (11)	0.1278 (3)	0.4478 (6)	0.062 (2)
C13	-0.5520 (14)	0.1049 (3)	0.4182 (7)	0.079 (3)
H13	-0.5713	0.0958	0.3531	0.095*
C14	-0.6863 (13)	0.0950 (3)	0.4838 (7)	0.080 (3)
H14	-0.7967	0.0792	0.4633	0.097*
C15	-0.6576 (12)	0.1084 (3)	0.5797 (7)	0.072 (2)
H15	-0.7488	0.1020	0.6244	0.086*
C16	-0.4919 (11)	0.1317 (3)	0.6100 (6)	0.064 (2)
H16	-0.4723	0.1403	0.6754	0.077*
C17	0.1855 (13)	0.0758 (3)	0.8683 (6)	0.079 (3)
H17A	0.2891	0.0759	0.9167	0.119*
H17B	0.2338	0.0741	0.8041	0.119*
H17C	0.1062	0.0489	0.8785	0.119*
C18	-0.2588 (11)	0.1335 (3)	0.9193 (6)	0.077 (2)

supplementary materials

H18A	-0.3086	0.1030	0.9012	0.115*
H18B	-0.3322	0.1576	0.8855	0.115*
H18C	-0.2649	0.1378	0.9887	0.115*
C19	0.1885 (11)	0.2012 (3)	0.8801 (5)	0.069 (2)
H19A	0.3035	0.1969	0.9196	0.103*
H19B	0.1171	0.2266	0.9062	0.103*
H19C	0.2183	0.2086	0.8141	0.103*
C20	-0.0375 (10)	0.2385 (3)	1.1363 (6)	0.067 (2)
H20A	-0.0001	0.2560	1.0803	0.100*
H20B	-0.1072	0.2588	1.1776	0.100*
H20C	-0.1158	0.2124	1.1149	0.100*
C21	0.2034 (13)	0.2164 (3)	1.3703 (5)	0.075 (3)
H21A	0.1478	0.2458	1.3891	0.112*
H21B	0.3308	0.2145	1.3980	0.112*
H21C	0.1306	0.1906	1.3939	0.112*
C22	0.4641 (10)	0.2008 (3)	1.1510 (6)	0.071 (2)
H22A	0.4750	0.1678	1.1356	0.107*
H22B	0.5669	0.2099	1.1957	0.107*
H22C	0.4684	0.2191	1.0920	0.107*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.0547 (3)	0.0418 (3)	0.0458 (3)	0.0031 (2)	-0.0036 (2)	-0.0021 (2)
Sn2	0.0490 (3)	0.0512 (3)	0.0451 (3)	-0.0101 (2)	0.0020 (2)	-0.0036 (2)
Cl1	0.138 (3)	0.195 (4)	0.163 (3)	0.045 (3)	0.005 (3)	0.002 (3)
Cl2	0.1043 (18)	0.130 (2)	0.0492 (12)	-0.0172 (16)	0.0086 (12)	-0.0131 (13)
O1	0.083 (4)	0.046 (3)	0.045 (3)	-0.007 (3)	-0.002 (2)	0.000 (2)
O2	0.083 (4)	0.047 (3)	0.041 (3)	-0.015 (3)	-0.001 (2)	-0.003 (2)
O3	0.074 (3)	0.049 (3)	0.045 (3)	0.017 (3)	-0.008 (2)	-0.007 (2)
O4	0.073 (4)	0.058 (3)	0.054 (3)	0.022 (3)	-0.009 (3)	-0.005 (3)
C1	0.071 (5)	0.046 (4)	0.048 (4)	-0.003 (4)	-0.007 (4)	0.001 (4)
C2	0.102 (7)	0.057 (5)	0.059 (5)	-0.019 (5)	-0.013 (5)	0.001 (4)
C3	0.122 (9)	0.044 (5)	0.068 (6)	-0.019 (5)	-0.016 (6)	-0.006 (4)
C4	0.145 (11)	0.071 (7)	0.088 (7)	-0.001 (7)	-0.021 (8)	-0.002 (6)
C5	0.181 (14)	0.084 (9)	0.113 (10)	0.012 (9)	-0.035 (10)	-0.006 (8)
C6	0.195 (16)	0.064 (8)	0.101 (10)	-0.014 (9)	-0.027 (10)	0.006 (7)
C7	0.178 (14)	0.078 (8)	0.089 (8)	-0.040 (9)	-0.007 (9)	-0.002 (7)
C8	0.153 (10)	0.060 (6)	0.084 (7)	-0.026 (6)	-0.011 (7)	-0.003 (6)
C9	0.051 (4)	0.057 (5)	0.043 (4)	0.007 (4)	-0.002 (3)	0.001 (4)
C10	0.055 (5)	0.063 (5)	0.047 (4)	0.003 (4)	-0.006 (3)	-0.001 (4)
C11	0.050 (4)	0.053 (5)	0.051 (4)	0.007 (3)	-0.007 (3)	0.000 (3)
C12	0.064 (5)	0.061 (5)	0.060 (5)	0.002 (4)	-0.009 (4)	-0.010 (4)
C13	0.087 (7)	0.074 (6)	0.074 (6)	-0.010 (5)	-0.012 (5)	-0.010 (5)
C14	0.077 (6)	0.068 (6)	0.094 (7)	-0.012 (5)	-0.012 (5)	-0.002 (5)
C15	0.063 (5)	0.070 (6)	0.083 (6)	-0.010 (4)	0.008 (5)	0.005 (5)
C16	0.068 (5)	0.066 (5)	0.058 (5)	0.000 (4)	0.000 (4)	0.000 (4)
C17	0.106 (7)	0.071 (6)	0.060 (5)	0.036 (5)	-0.013 (5)	-0.009 (4)

C18	0.059 (5)	0.093 (7)	0.078 (6)	-0.008 (5)	0.006 (4)	0.006 (5)
C19	0.074 (5)	0.075 (6)	0.057 (5)	-0.017 (4)	0.000 (4)	0.012 (4)
C20	0.050 (5)	0.063 (5)	0.086 (6)	0.003 (4)	0.001 (4)	-0.010 (4)
C21	0.117 (7)	0.068 (6)	0.039 (4)	-0.029 (5)	0.005 (4)	-0.004 (4)
C22	0.057 (5)	0.082 (6)	0.076 (5)	-0.006 (4)	0.006 (4)	-0.003 (5)

Geometric parameters (Å, °)

Sn1—C17	2.107 (7)	C10—C11	1.522 (9)
Sn1—C18	2.117 (7)	C10—H10A	0.9700
Sn1—C19	2.120 (7)	C10—H10B	0.9700
Sn1—O3	2.194 (4)	C11—C16	1.381 (10)
Sn1—O1	2.396 (5)	C11—C12	1.397 (10)
Sn2—C22	2.102 (7)	C12—C13	1.345 (11)
Sn2—C20	2.113 (7)	C13—C14	1.371 (12)
Sn2—C21	2.117 (7)	C13—H13	0.9300
Sn2—O2	2.207 (5)	C14—C15	1.369 (11)
Sn2—O4 ⁱ	2.432 (5)	C14—H14	0.9300
C11—C4	1.746 (12)	C15—C16	1.389 (11)
C12—C12	1.747 (8)	C15—H15	0.9300
O1—C1	1.249 (8)	C16—H16	0.9300
O2—C1	1.263 (8)	C17—H17A	0.9600
O3—C9	1.273 (8)	C17—H17B	0.9600
O4—C9	1.235 (8)	C17—H17C	0.9600
O4—Sn2 ⁱⁱ	2.432 (5)	C18—H18A	0.9600
C1—C2	1.510 (10)	C18—H18B	0.9600
C2—C3	1.487 (11)	C18—H18C	0.9600
C2—H2A	0.9700	C19—H19A	0.9600
C2—H2B	0.9700	C19—H19B	0.9600
C3—C4	1.359 (14)	C19—H19C	0.9600
C3—C8	1.411 (13)	C20—H20A	0.9600
C4—C5	1.395 (15)	C20—H20B	0.9600
C5—C6	1.380 (17)	C20—H20C	0.9600
C5—H5	0.9300	C21—H21A	0.9600
C6—C7	1.387 (17)	C21—H21B	0.9600
C6—H6	0.9300	C21—H21C	0.9600
C7—C8	1.358 (14)	C22—H22A	0.9600
C7—H7	0.9300	C22—H22B	0.9600
C8—H8	0.9300	C22—H22C	0.9600
C9—C10	1.515 (9)		
C17—Sn1—C18	119.2 (4)	C9—C10—H10B	108.9
C17—Sn1—C19	114.4 (4)	C11—C10—H10B	108.9
C18—Sn1—C19	125.2 (3)	H10A—C10—H10B	107.7
C17—Sn1—O3	90.2 (2)	C16—C11—C12	116.7 (7)
C18—Sn1—O3	94.6 (3)	C16—C11—C10	123.3 (6)
C19—Sn1—O3	95.6 (2)	C12—C11—C10	120.0 (7)
C17—Sn1—O1	89.8 (2)	C13—C12—C11	122.4 (8)
C18—Sn1—O1	86.3 (3)	C13—C12—C12	118.5 (7)

supplementary materials

C19—Sn1—O1	83.4 (2)	C11—C12—C12	119.1 (6)
O3—Sn1—O1	178.93 (18)	C12—C13—C14	120.1 (8)
C22—Sn2—C20	122.9 (3)	C12—C13—H13	119.9
C22—Sn2—C21	118.9 (3)	C14—C13—H13	119.9
C20—Sn2—C21	116.5 (3)	C15—C14—C13	119.9 (8)
C22—Sn2—O2	94.9 (3)	C15—C14—H14	120.1
C20—Sn2—O2	96.2 (3)	C13—C14—H14	120.1
C21—Sn2—O2	92.0 (2)	C14—C15—C16	119.7 (8)
C22—Sn2—O4 ⁱ	86.0 (3)	C14—C15—H15	120.1
C20—Sn2—O4 ⁱ	87.3 (2)	C16—C15—H15	120.1
C21—Sn2—O4 ⁱ	83.4 (2)	C11—C16—C15	121.2 (7)
O2—Sn2—O4 ⁱ	175.08 (16)	C11—C16—H16	119.4
C1—O1—Sn1	143.5 (5)	C15—C16—H16	119.4
C1—O2—Sn2	117.0 (4)	Sn1—C17—H17A	109.5
C9—O3—Sn1	119.0 (4)	Sn1—C17—H17B	109.5
C9—O4—Sn2 ⁱⁱ	141.7 (5)	H17A—C17—H17B	109.5
O1—C1—O2	122.9 (7)	Sn1—C17—H17C	109.5
O1—C1—C2	120.6 (6)	H17A—C17—H17C	109.5
O2—C1—C2	116.5 (6)	H17B—C17—H17C	109.5
C3—C2—C1	116.9 (7)	Sn1—C18—H18A	109.5
C3—C2—H2A	108.1	Sn1—C18—H18B	109.5
C1—C2—H2A	108.1	H18A—C18—H18B	109.5
C3—C2—H2B	108.1	Sn1—C18—H18C	109.5
C1—C2—H2B	108.1	H18A—C18—H18C	109.5
H2A—C2—H2B	107.3	H18B—C18—H18C	109.5
C4—C3—C8	117.4 (10)	Sn1—C19—H19A	109.5
C4—C3—C2	122.9 (10)	Sn1—C19—H19B	109.5
C8—C3—C2	119.8 (10)	H19A—C19—H19B	109.5
C3—C4—C5	124.7 (13)	Sn1—C19—H19C	109.5
C3—C4—C11	119.3 (9)	H19A—C19—H19C	109.5
C5—C4—C11	116.0 (12)	H19B—C19—H19C	109.5
C6—C5—C4	113.0 (14)	Sn2—C20—H20A	109.5
C6—C5—H5	123.5	Sn2—C20—H20B	109.5
C4—C5—H5	123.5	H20A—C20—H20B	109.5
C5—C6—C7	127.0 (13)	Sn2—C20—H20C	109.5
C5—C6—H6	116.5	H20A—C20—H20C	109.5
C7—C6—H6	116.5	H20B—C20—H20C	109.5
C8—C7—C6	115.4 (13)	Sn2—C21—H21A	109.5
C8—C7—H7	122.3	Sn2—C21—H21B	109.5
C6—C7—H7	122.3	H21A—C21—H21B	109.5
C7—C8—C3	122.5 (12)	Sn2—C21—H21C	109.5
C7—C8—H8	118.8	H21A—C21—H21C	109.5
C3—C8—H8	118.8	H21B—C21—H21C	109.5
O4—C9—O3	122.4 (6)	Sn2—C22—H22A	109.5
O4—C9—C10	121.7 (6)	Sn2—C22—H22B	109.5
O3—C9—C10	115.9 (6)	H22A—C22—H22B	109.5
C9—C10—C11	113.2 (6)	Sn2—C22—H22C	109.5
C9—C10—H10A	108.9	H22A—C22—H22C	109.5

C11—C10—H10A	108.9	H22B—C22—H22C	109.5
C17—Sn1—O1—C1	-67.1 (9)	C11—C4—C5—C6	179.9 (9)
C18—Sn1—O1—C1	52.3 (9)	C4—C5—C6—C7	-0.1 (19)
C19—Sn1—O1—C1	178.4 (9)	C5—C6—C7—C8	-1.2 (19)
O3—Sn1—O1—C1	-159 (9)	C6—C7—C8—C3	2.0 (15)
C22—Sn2—O2—C1	-66.8 (6)	C4—C3—C8—C7	-1.6 (14)
C20—Sn2—O2—C1	57.1 (6)	C2—C3—C8—C7	177.3 (8)
C21—Sn2—O2—C1	174.0 (6)	Sn2 ⁱⁱ —O4—C9—O3	160.1 (5)
O4 ⁱ —Sn2—O2—C1	-167 (2)	Sn2 ⁱⁱ —O4—C9—C10	-22.1 (12)
C17—Sn1—O3—C9	-174.5 (6)	Sn1—O3—C9—O4	-2.7 (9)
C18—Sn1—O3—C9	66.2 (6)	Sn1—O3—C9—C10	179.3 (4)
C19—Sn1—O3—C9	-60.0 (6)	O4—C9—C10—C11	-91.2 (9)
O1—Sn1—O3—C9	-83 (10)	O3—C9—C10—C11	86.7 (8)
Sn1—O1—C1—O2	-163.3 (5)	C9—C10—C11—C16	14.6 (10)
Sn1—O1—C1—C2	14.4 (13)	C9—C10—C11—C12	-166.5 (7)
Sn2—O2—C1—O1	4.5 (10)	C16—C11—C12—C13	0.3 (11)
Sn2—O2—C1—C2	-173.3 (5)	C10—C11—C12—C13	-178.7 (8)
O1—C1—C2—C3	148.0 (8)	C16—C11—C12—C12	179.6 (6)
O2—C1—C2—C3	-34.2 (11)	C10—C11—C12—C12	0.6 (10)
C1—C2—C3—C4	-72.9 (11)	C11—C12—C13—C14	0.1 (13)
C1—C2—C3—C8	108.2 (9)	C12—C12—C13—C14	-179.1 (7)
C8—C3—C4—C5	0.2 (15)	C12—C13—C14—C15	-0.1 (14)
C2—C3—C4—C5	-178.7 (9)	C13—C14—C15—C16	-0.4 (13)
C8—C3—C4—C11	-179.1 (7)	C12—C11—C16—C15	-0.8 (11)
C2—C3—C4—C11	2.0 (13)	C10—C11—C16—C15	178.1 (7)
C3—C4—C5—C6	0.6 (17)	C14—C15—C16—C11	0.9 (12)

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (ii) $x-1/2, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

