

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(μ -cyclohexane-1,3-dicarboxylato)- $\kappa^3 O^1:O^4, O^4'; \kappa^3 O^1, O^1':O^4$ -bis[aqua(1,10-phenanthroline- $\kappa^2 N, N'$)zinc(II)]

Mohd. Razali Rizal and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 31 August 2009; accepted 1 September 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.014$ Å; R factor = 0.069; wR factor = 0.177; data-to-parameter ratio = 12.6.

The cyclohexane-1,3-dicarboxylate dianion in the dinuclear centrosymmetric title compound, $[Zn_2(C_8H_{10}O_4)_2(C_{12}H_8N_2)_2(H_2O)_2]$, has a chair conformation with both carboxylate groups in equatorial positions. One carboxylate group chelates a Zn^{II} atom, whereas the other binds through one O atom only to confer a six-coordinate status to the N -heterocycle-chelated water-coordinated Zn^{II} atom. Adjacent dinuclear molecules are linked by $O-H \cdots O$ hydrogen bonds into a linear chain.

Related literature

For the isostructural manganese(II) analog, see: Thirumurugan *et al.* (2006). For a review of the molecular architectures of metal carboxylate adducts of 2,2'-bipyridine-like ligands, see: Ye *et al.* (2003).



Experimental

Crystal data

$[Zn_2(C_8H_{10}O_4)_2(C_{12}H_8N_2)_2(H_2O)_2]$
 $M_r = 867.50$
 Monoclinic, $P2_1/c$
 $a = 9.6172$ (2) Å
 $b = 17.4722$ (5) Å
 $c = 11.4822$ (3) Å
 $\beta = 104.393$ (2)°
 $V = 1868.84$ (8) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.35$ mm⁻¹
 $T = 100$ K
 $0.20 \times 0.16 \times 0.15$ mm

Data collection

Bruker SMART APEX diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{min} = 0.774$, $T_{max} = 0.823$
 9952 measured reflections
 3289 independent reflections
 2706 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.069$
 $wR(F^2) = 0.177$
 $S = 1.09$
 3289 reflections
 261 parameters
 74 restraints
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{max} = 1.17$ e Å⁻³
 $\Delta\rho_{min} = -0.51$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O1w-H11 \cdots O2$	0.84 (6)	1.80 (7)	2.614 (6)	162 (8)
$O1w-H12 \cdots O4^i$	0.84 (6)	1.89 (3)	2.701 (6)	161 (8)

Symmetry code: (i) $x - 1, y, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XSELL (Sheldrick, 2008); software used to prepare material for publication: publCIF (Westrip, 2009).

We thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2534).

References

- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). Acta Cryst. A64, 112–122.
 Thirumurugan, T., Avinash, M. B. & Rao, C. N. R. (2006). Dalton Trans. pp. 221–228.
 Westrip, S. P. (2009). publCIF. In preparation.
 Ye, B.-H., Tong, M.-L. & Chen, X.-M. (2003). Coord. Chem. Rev. 246, 185–202.

supplementary materials

Acta Cryst. (2009). E65, m1178 [doi:10.1107/S1600536809035144]

Bis(μ -cyclohexane-1,3-dicarboxylato)- $\kappa^3 O^1:O^4,O^4';\kappa^3 O^1,O^1':O^4$ -bis[aqua(1,10-phenanthroline- $\kappa^2 N,N'$)zinc(II)]

M. R. Rizal and S. W. Ng

Experimental

Zinc acetate (0.10 g, 0.46 mmol), cyclohexane-1,3-dicarboxylic acid (mixture of *cis*- and *trans*-isomers) (0.08 g, 0.46 mmol) and 1,10-phenanthroline (0.09 g, 0.46 mmol) along with water (18 ml) were heated in a 23-ml Teflon-lined stainless-steel Parr bomb. The bomb was heated at 403 K for 3 days. The bomb was cooled to room temperature at 5 K per hour. Tiny crystals were isolated from the solution.

Refinement

Hydrogen atoms were included in the refinement in the riding model approximation with C–H 0.95 – 1.00 Å, and with $U(H)$ $1.2U_{eq}(C)$. The water H-atoms were located in a difference Fourier map, and were refined with a distance restraint of O–H 0.84±0.01 Å; their displacement factors were refined.

The carbon atoms of the 1,10-phenanthroline molecule displayed somewhat elongated thermal ellipsoids. As such, their anisotropic displacement factors were restrained to be nearly isotropic.

The final difference Fourier map had a peak in the vicinity of the C1 and C6 atoms.

Figures




Fig. 1. 50% Probability displacement ellipsoid plot of $Zn_2(H_2O)_2(C_{12}H_8N_2)_2(C_8H_{10}O_4)_2$. Hydrogen atoms are drawn as spheres of arbitrary radius. Symmetry code (i): 2-x, 1-y, 1-z.

Bis(μ -cyclohexane-1,3-dicarboxylato)- $\kappa^3 O^1: O^4,O^4';\kappa^3 O^1,O^1':O^4$ - bis[aqua(1,10-phenanthroline- $\kappa^2 N,N'$)zinc(II)]

Crystal data

$[Zn_2(C_8H_{10}O_4)_2(C_{12}H_8N_2)_2(H_2O)_2]$

$M_r = 867.50$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 9.6172$ (2) Å

$b = 17.4722$ (5) Å

$F_{000} = 896$

$D_x = 1.542$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2661 reflections

$\theta = 2.2$ – 26.3°

$\mu = 1.35$ mm⁻¹

supplementary materials

$c = 11.4822 (3) \text{ \AA}$
 $\beta = 104.393 (2)^\circ$
 $V = 1868.84 (8) \text{ \AA}^3$
 $Z = 2$

$T = 100 \text{ K}$
Block, colorless
 $0.20 \times 0.16 \times 0.15 \text{ mm}$

Data collection

Bruker SMART APEX diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 100 \text{ K}$
 ω scans
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.774, T_{\max} = 0.823$
9952 measured reflections

3289 independent reflections
2706 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$
 $\theta_{\max} = 25.0^\circ$
 $\theta_{\min} = 2.2^\circ$
 $h = -11 \rightarrow 11$
 $k = -20 \rightarrow 20$
 $l = -10 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.069$
 $wR(F^2) = 0.177$
 $S = 1.09$
3289 reflections
261 parameters
74 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0653P)^2 + 9.3251P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.17 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.51 \text{ e \AA}^{-3}$
Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.70076 (7)	0.51170 (4)	0.71412 (7)	0.0348 (3)
O1	0.8609 (5)	0.5855 (3)	0.7036 (5)	0.0516 (13)
O2	0.7359 (6)	0.6719 (3)	0.5775 (5)	0.0541 (14)
O3	1.1660 (5)	0.5852 (3)	0.3118 (4)	0.0461 (12)
O4	1.3286 (4)	0.5440 (3)	0.4667 (4)	0.0414 (11)
O1w	0.5440 (5)	0.5921 (3)	0.6510 (4)	0.0367 (10)
H11	0.593 (7)	0.626 (3)	0.627 (7)	0.06 (3)*
H12	0.471 (5)	0.572 (4)	0.606 (6)	0.05 (2)*
N1	0.5460 (6)	0.4333 (3)	0.7566 (5)	0.0394 (13)
N2	0.7539 (6)	0.5156 (4)	0.9034 (5)	0.0495 (16)
C1	1.1790 (11)	0.7240 (5)	0.4694 (11)	0.085 (3)

H1A	1.1088	0.7318	0.3912	0.102*
H1B	1.2750	0.7386	0.4598	0.102*
C2	1.1393 (10)	0.7760 (5)	0.5650 (11)	0.090 (4)
H2A	1.2152	0.7730	0.6410	0.108*
H2B	1.1322	0.8298	0.5371	0.108*
C3	0.9953 (11)	0.7504 (5)	0.5869 (12)	0.096 (4)
H3A	0.9754	0.7809	0.6536	0.116*
H3B	0.9179	0.7609	0.5139	0.116*
C4	0.9932 (9)	0.6721 (5)	0.6155 (10)	0.070 (3)
H4	1.0652	0.6653	0.6943	0.084*
C5	1.0389 (7)	0.6182 (4)	0.5277 (7)	0.0426 (16)
H5A	0.9630	0.6162	0.4516	0.051*
H5B	1.0517	0.5659	0.5620	0.051*
C6	1.1801 (9)	0.6455 (5)	0.5022 (9)	0.064 (2)
H6	1.2534	0.6415	0.5808	0.077*
C7	1.2268 (6)	0.5891 (4)	0.4205 (6)	0.0369 (15)
C8	0.8505 (7)	0.6411 (4)	0.6334 (8)	0.0499 (19)
C9	0.4467 (8)	0.3928 (4)	0.6845 (8)	0.0502 (18)
H9	0.4302	0.4013	0.6004	0.060*
C10	0.3633 (9)	0.3378 (5)	0.7238 (10)	0.066 (2)
H10	0.2932	0.3094	0.6675	0.079*
C11	0.3834 (10)	0.3256 (5)	0.8412 (11)	0.074 (3)
H11A	0.3270	0.2884	0.8689	0.089*
C12	0.4866 (10)	0.3671 (5)	0.9239 (8)	0.068 (3)
C13	0.5188 (13)	0.3622 (6)	1.0544 (11)	0.089 (3)
H13	0.4655	0.3274	1.0899	0.107*
C14	0.6173 (15)	0.4034 (7)	1.1242 (10)	0.094 (4)
H14	0.6321	0.3976	1.2086	0.113*
C15	0.7051 (12)	0.4574 (6)	1.0810 (8)	0.079 (3)
C16	0.8171 (14)	0.5024 (7)	1.1488 (10)	0.096 (4)
H16	0.8410	0.4975	1.2338	0.115*
C17	0.8914 (12)	0.5518 (7)	1.0997 (10)	0.092 (4)
H17	0.9654	0.5822	1.1484	0.110*
C18	0.8568 (8)	0.5576 (5)	0.9736 (7)	0.065 (2)
H18	0.9087	0.5927	0.9373	0.078*
C19	0.6786 (9)	0.4657 (5)	0.9548 (6)	0.053 (2)
C20	0.5701 (8)	0.4215 (4)	0.8787 (7)	0.0470 (18)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0269 (4)	0.0414 (4)	0.0374 (4)	-0.0027 (3)	0.0104 (3)	-0.0022 (3)
O1	0.031 (2)	0.049 (3)	0.076 (4)	-0.009 (2)	0.014 (2)	-0.004 (3)
O2	0.056 (3)	0.037 (3)	0.078 (4)	-0.011 (2)	0.033 (3)	-0.001 (2)
O3	0.032 (2)	0.060 (3)	0.044 (3)	0.002 (2)	0.004 (2)	0.001 (2)
O4	0.028 (2)	0.059 (3)	0.037 (3)	-0.005 (2)	0.0079 (19)	0.004 (2)
O1w	0.026 (2)	0.049 (3)	0.035 (3)	-0.002 (2)	0.008 (2)	0.002 (2)
N1	0.037 (3)	0.040 (3)	0.046 (3)	0.005 (2)	0.021 (3)	0.007 (3)

supplementary materials

N2	0.040 (3)	0.067 (4)	0.034 (3)	0.026 (3)	-0.005 (3)	-0.017 (3)
C1	0.082 (6)	0.062 (6)	0.138 (10)	-0.014 (5)	0.079 (7)	-0.008 (6)
C2	0.080 (6)	0.057 (5)	0.164 (11)	-0.035 (5)	0.087 (7)	-0.049 (6)
C3	0.086 (7)	0.068 (6)	0.166 (11)	-0.039 (5)	0.090 (8)	-0.057 (7)
C4	0.057 (5)	0.051 (5)	0.120 (8)	-0.012 (4)	0.057 (5)	-0.016 (5)
C5	0.038 (4)	0.040 (4)	0.055 (5)	-0.008 (3)	0.019 (3)	-0.001 (3)
C6	0.055 (5)	0.056 (5)	0.098 (7)	-0.016 (4)	0.049 (5)	-0.025 (5)
C7	0.028 (3)	0.046 (4)	0.040 (4)	-0.011 (3)	0.015 (3)	-0.005 (3)
C8	0.036 (4)	0.047 (4)	0.075 (5)	-0.006 (3)	0.028 (4)	-0.022 (4)
C9	0.044 (4)	0.045 (4)	0.069 (5)	-0.008 (3)	0.028 (4)	-0.001 (3)
C10	0.054 (4)	0.049 (4)	0.109 (7)	-0.001 (3)	0.047 (5)	0.004 (4)
C11	0.065 (5)	0.050 (4)	0.126 (7)	0.019 (4)	0.059 (5)	0.028 (5)
C12	0.085 (5)	0.065 (5)	0.079 (5)	0.053 (4)	0.064 (5)	0.043 (4)
C13	0.107 (7)	0.086 (6)	0.100 (7)	0.064 (5)	0.073 (6)	0.056 (5)
C14	0.122 (7)	0.112 (7)	0.062 (6)	0.084 (6)	0.050 (6)	0.036 (5)
C15	0.091 (6)	0.090 (6)	0.055 (5)	0.067 (5)	0.020 (5)	0.009 (5)
C16	0.112 (7)	0.108 (7)	0.056 (5)	0.080 (6)	-0.003 (5)	-0.018 (5)
C17	0.077 (6)	0.102 (7)	0.071 (6)	0.055 (5)	-0.029 (5)	-0.044 (5)
C18	0.049 (4)	0.078 (5)	0.053 (4)	0.035 (4)	-0.015 (4)	-0.028 (4)
C19	0.062 (4)	0.067 (5)	0.035 (4)	0.046 (4)	0.019 (3)	0.012 (3)
C20	0.056 (4)	0.050 (4)	0.045 (4)	0.029 (3)	0.031 (3)	0.018 (3)

Geometric parameters (Å, °)

Zn1—O1	2.035 (5)	C4—C5	1.521 (10)
Zn1—O3 ⁱ	2.188 (5)	C4—C8	1.536 (9)
Zn1—O4 ⁱ	2.247 (5)	C4—H4	1.0000
Zn1—O1w	2.056 (5)	C5—C6	1.536 (9)
Zn1—N1	2.166 (5)	C5—H5A	0.9900
Zn1—N2	2.107 (6)	C5—H5B	0.9900
O1—C8	1.250 (10)	C6—C7	1.504 (10)
O2—C8	1.251 (9)	C6—H6	1.0000
O3—C7	1.241 (8)	C9—C10	1.397 (10)
O3—Zn1 ⁱ	2.188 (5)	C9—H9	0.9500
O4—C7	1.266 (8)	C10—C11	1.331 (14)
O4—Zn1 ⁱ	2.247 (5)	C10—H10	0.9500
O1w—H11	0.84 (6)	C11—C12	1.394 (14)
O1w—H12	0.84 (6)	C11—H11A	0.9500
N1—C9	1.306 (9)	C12—C20	1.423 (11)
N1—C20	1.378 (9)	C12—C13	1.456 (15)
N2—C18	1.331 (10)	C13—C14	1.297 (16)
N2—C19	1.358 (11)	C13—H13	0.9500
C1—C6	1.422 (12)	C14—C15	1.434 (17)
C1—C2	1.544 (12)	C14—H14	0.9500
C1—H1A	0.9900	C15—C19	1.416 (11)
C1—H1B	0.9900	C15—C16	1.403 (16)
C2—C3	1.535 (10)	C16—C17	1.331 (17)
C2—H2A	0.9900	C16—H16	0.9500

C2—H2B	0.9900	C17—C18	1.407 (14)
C3—C4	1.408 (13)	C17—H17	0.9500
C3—H3A	0.9900	C18—H18	0.9500
C3—H3B	0.9900	C19—C20	1.412 (12)
O1—Zn1—O3 ⁱ	90.3 (2)	C6—C5—H5A	109.6
O1—Zn1—O4 ⁱ	98.2 (2)	C4—C5—H5B	109.6
O1—Zn1—O1w	92.6 (2)	C6—C5—H5B	109.6
O1—Zn1—N2	92.7 (2)	H5A—C5—H5B	108.1
O1—Zn1—N1	170.6 (2)	C1—C6—C7	116.8 (8)
O3 ⁱ —Zn1—O4 ⁱ	58.8 (2)	C1—C6—C5	113.7 (7)
O3 ⁱ —Zn1—O1w	152.1 (2)	C7—C6—C5	109.3 (6)
O3 ⁱ —Zn1—N1	90.0 (2)	C1—C6—H6	105.3
O3 ⁱ —Zn1—N2	99.5 (2)	C7—C6—H6	105.3
O4 ⁱ —Zn1—O1w	93.3 (2)	C5—C6—H6	105.3
O4 ⁱ —Zn1—N1	89.9 (2)	O3—C7—O4	120.6 (6)
O4 ⁱ —Zn1—N2	155.5 (2)	O3—C7—C6	121.6 (7)
O1w—Zn1—N1	91.6 (2)	O4—C7—C6	117.8 (7)
O1w—Zn1—N2	108.1 (2)	O2—C8—O1	125.7 (6)
N1—Zn1—N2	78.0 (2)	O2—C8—C4	118.9 (8)
C8—O1—Zn1	126.2 (5)	O1—C8—C4	115.4 (7)
C7—O3—Zn1 ⁱ	91.9 (4)	N1—C9—C10	123.7 (8)
C7—O4—Zn1 ⁱ	88.6 (4)	N1—C9—H9	118.1
Zn1—O1w—H11	100 (6)	C10—C9—H9	118.1
Zn1—O1w—H12	111 (5)	C11—C10—C9	119.2 (9)
H11—O1w—H12	123 (8)	C11—C10—H10	120.4
C9—N1—C20	118.4 (6)	C9—C10—H10	120.4
C9—N1—Zn1	129.4 (5)	C10—C11—C12	120.3 (8)
C20—N1—Zn1	111.9 (5)	C10—C11—H11A	119.8
C18—N2—C19	119.2 (7)	C12—C11—H11A	119.8
C18—N2—Zn1	126.6 (6)	C11—C12—C20	118.1 (8)
C19—N2—Zn1	114.1 (5)	C11—C12—C13	127.1 (9)
C6—C1—C2	111.5 (8)	C20—C12—C13	114.9 (10)
C6—C1—H1A	109.3	C14—C13—C12	122.6 (10)
C2—C1—H1A	109.3	C14—C13—H13	118.7
C6—C1—H1B	109.3	C12—C13—H13	118.7
C2—C1—H1B	109.3	C13—C14—C15	123.7 (10)
H1A—C1—H1B	108.0	C13—C14—H14	118.2
C3—C2—C1	109.7 (7)	C15—C14—H14	118.2
C3—C2—H2A	109.7	C14—C15—C19	116.7 (11)
C1—C2—H2A	109.7	C14—C15—C16	127.9 (11)
C3—C2—H2B	109.7	C19—C15—C16	115.4 (11)
C1—C2—H2B	109.7	C17—C16—C15	123.2 (10)
H2A—C2—H2B	108.2	C17—C16—H16	118.4
C4—C3—C2	112.8 (8)	C15—C16—H16	118.4
C4—C3—H3A	109.0	C16—C17—C18	117.9 (11)
C2—C3—H3A	109.0	C16—C17—H17	121.1

supplementary materials

C4—C3—H3B	109.0	C18—C17—H17	121.1
C2—C3—H3B	109.0	N2—C18—C17	122.2 (11)
H3A—C3—H3B	107.8	N2—C18—H18	118.9
C3—C4—C5	115.1 (8)	C17—C18—H18	118.9
C3—C4—C8	116.0 (7)	N2—C19—C20	118.3 (6)
C5—C4—C8	106.6 (6)	N2—C19—C15	122.0 (9)
C3—C4—H4	106.1	C20—C19—C15	119.6 (9)
C5—C4—H4	106.1	N1—C20—C19	117.3 (6)
C8—C4—H4	106.1	N1—C20—C12	120.2 (8)
C4—C5—C6	110.4 (6)	C19—C20—C12	122.5 (8)
C4—C5—H5A	109.6		
O1w—Zn1—O1—C8	-26.9 (6)	C3—C4—C8—O2	29.2 (13)
N2—Zn1—O1—C8	-135.2 (6)	C5—C4—C8—O2	-100.4 (8)
O3 ⁱ —Zn1—O1—C8	125.3 (6)	C3—C4—C8—O1	-151.5 (9)
O4 ⁱ —Zn1—O1—C8	66.8 (6)	C5—C4—C8—O1	78.9 (9)
O1w—Zn1—N1—C9	72.9 (6)	C20—N1—C9—C10	-0.2 (10)
N2—Zn1—N1—C9	-178.9 (6)	Zn1—N1—C9—C10	172.7 (5)
O3 ⁱ —Zn1—N1—C9	-79.2 (6)	N1—C9—C10—C11	0.8 (12)
O4 ⁱ —Zn1—N1—C9	-20.4 (6)	C9—C10—C11—C12	-0.2 (12)
O1w—Zn1—N1—C20	-113.8 (4)	C10—C11—C12—C20	-0.9 (11)
N2—Zn1—N1—C20	-5.6 (4)	C10—C11—C12—C13	178.2 (7)
O3 ⁱ —Zn1—N1—C20	94.1 (4)	C11—C12—C13—C14	-180.0 (9)
O4 ⁱ —Zn1—N1—C20	152.9 (4)	C20—C12—C13—C14	-0.9 (12)
O1—Zn1—N2—C18	2.1 (6)	C12—C13—C14—C15	-0.5 (14)
O1w—Zn1—N2—C18	-91.6 (5)	C13—C14—C15—C19	1.1 (13)
N1—Zn1—N2—C18	-179.3 (6)	C13—C14—C15—C16	-177.6 (9)
O3 ⁱ —Zn1—N2—C18	92.8 (6)	C14—C15—C16—C17	-179.2 (9)
O4 ⁱ —Zn1—N2—C18	118.8 (6)	C19—C15—C16—C17	2.2 (12)
O1—Zn1—N2—C19	-173.6 (4)	C15—C16—C17—C18	-1.2 (14)
O1w—Zn1—N2—C19	92.7 (5)	C19—N2—C18—C17	0.5 (10)
N1—Zn1—N2—C19	5.0 (4)	Zn1—N2—C18—C17	-175.0 (5)
O3 ⁱ —Zn1—N2—C19	-82.9 (4)	C16—C17—C18—N2	-0.2 (12)
O4 ⁱ —Zn1—N2—C19	-56.9 (7)	C18—N2—C19—C20	-179.8 (6)
C6—C1—C2—C3	-55.1 (14)	Zn1—N2—C19—C20	-3.8 (7)
C1—C2—C3—C4	53.5 (14)	C18—N2—C19—C15	0.6 (9)
C2—C3—C4—C5	-52.1 (13)	Zn1—N2—C19—C15	176.6 (5)
C2—C3—C4—C8	-177.5 (9)	C14—C15—C19—N2	179.3 (6)
C3—C4—C5—C6	48.8 (11)	C16—C15—C19—N2	-1.8 (10)
C8—C4—C5—C6	178.9 (8)	C14—C15—C19—C20	-0.2 (10)
C2—C1—C6—C7	-176.1 (8)	C16—C15—C19—C20	178.6 (6)
C2—C1—C6—C5	55.2 (12)	C9—N1—C20—C19	179.7 (6)
C4—C5—C6—C1	-50.4 (11)	Zn1—N1—C20—C19	5.5 (7)
C4—C5—C6—C7	177.0 (8)	C9—N1—C20—C12	-0.9 (9)
Zn1 ⁱ —O3—C7—O4	1.5 (6)	Zn1—N1—C20—C12	-175.0 (5)
Zn1 ⁱ —O3—C7—C6	-176.8 (5)	N2—C19—C20—N1	-1.3 (9)
Zn1 ⁱ —O4—C7—O3	-1.4 (6)	C15—C19—C20—N1	178.3 (6)

Zn1 ⁱ —O4—C7—C6	176.9 (5)	N2—C19—C20—C12	179.3 (6)
C1—C6—C7—O3	-56.5 (10)	C15—C19—C20—C12	-1.2 (10)
C5—C6—C7—O3	74.4 (9)	C11—C12—C20—N1	1.4 (9)
C1—C6—C7—O4	125.2 (9)	C13—C12—C20—N1	-177.7 (6)
C5—C6—C7—O4	-103.9 (8)	C11—C12—C20—C19	-179.1 (6)
Zn1—O1—C8—O2	17.5 (11)	C13—C12—C20—C19	1.7 (9)
Zn1—O1—C8—C4	-161.7 (5)		

Symmetry codes: (i) $-x+2, -y+1, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1w—H11 \cdots O2	0.84 (6)	1.80 (7)	2.614 (6)	162 (8)
O1w—H12 \cdots O4 ⁱⁱ	0.84 (6)	1.89 (3)	2.701 (6)	161 (8)

Symmetry codes: (ii) $x-1, y, z$.

