

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(5,5-diphenylhydantoinato- κN^3)-(ethylenediamine)zinc(II)

Xilan Hu,^{a*} Xingyou Xu,^b Daqi Wang^c and Yu Zhang^a^aHuaihai Institute of Technology, Jiangsu 222005, People's Republic of China,^bHuaiyin Institute of Technology, Jiangsu 223003, People's Republic of China, and^cCollege of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China

Correspondence e-mail: huxilan@hhit.edu.cn


Received 10 September 2009; accepted 20 October 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.034; wR factor = 0.089; data-to-parameter ratio = 13.2.

In the title compound, $[Zn(C_{15}H_{11}N_2O_2)_2(C_2H_8N_2)]$, the Zn^{II} atom is coordinated in a distorted tetrahedral geometry. Intramolecular $N-H\cdots O$, $C-H\cdots O$ and $C-H\cdots N$ hydrogen bonds occur. In the crystal, molecules are linked by intermolecular $N-H\cdots O$ hydrogen bonds, forming a three-dimensional network.

Related literature

5,5-Diphenylimidazoline-2,4-dione (phenytoin) is widely used in the treatment of epilepsy and should be an excellent ligand for transition metal complexes, see: Milne *et al.* (1999); Akitsu & Einaga (2005); Akitsu *et al.* (1997). For complexes with 5,5-diphenylhydantoinate, see: Hu, Xu, Wang & Xu (2006); Hu, Xu, Xu & Wang (2006).



Experimental

Crystal data

 $[Zn(C_{15}H_{11}N_2O_2)_2(C_2H_8N_2)]$ $M_r = 627.99$ Triclinic, $P\bar{1}$ $a = 9.702$ (1) Å $b = 13.052$ (2) Å $c = 13.293$ (2) Å
 $\alpha = 109.114$ (2)°
 $\beta = 109.462$ (2)°
 $\gamma = 93.020$ (10)°
 $V = 1473.1$ (3) Å³
 $Z = 2$
Mo $K\alpha$ radiation $\mu = 0.88$ mm⁻¹ $T = 298$ K $0.50 \times 0.46 \times 0.32$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996) $T_{\min} = 0.667$, $T_{\max} = 0.766$

7725 measured reflections

5124 independent reflections

4452 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.017$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.034$ $wR(F^2) = 0.089$ $S = 1.03$

5124 reflections

388 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.61$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.45$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N6-H6B\cdots O2$	0.90	2.18	2.858 (3)	132
$C9-H9\cdots O2$	0.93	2.36	2.992 (3)	125
$C11-H11\cdots N1$	0.93	2.52	2.860 (4)	102
$C24-H24\cdots O4$	0.93	2.39	3.042 (4)	127
$C26-H26\cdots N3$	0.93	2.51	2.854 (4)	102
$N1-H1\cdots O1^i$	0.86	2.16	3.008 (3)	167
$N3-H3\cdots O3^{ii}$	0.86	2.04	2.843 (3)	156
$N5-H5A\cdots O4^{iii}$	0.90	1.97	2.855 (3)	170
$N5-H5B\cdots O1^{iii}$	0.90	2.18	3.003 (3)	152

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, -y+2, -z+1$; (iii) $-x, -y+1, -z+1$.

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We are grateful for the financial support from the Key Project for Fundamental Research of the Education Committee of Jiangsu Province (07KJA15011) and the Natural Science Foundation of Huaihai Institute of Technology (KX07042).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2240).

References

- Akitsu, T. & Einaga, Y. (2005). *Acta Cryst.* **E61**, m183–m186.
 Akitsu, T., Komorita, S., Kushi, Y., Li, C., Kanehisa, N. & Kai, Y. (1997). *Bull. Chem. Soc. Jpn.* **70**, 821–827.
 Hu, X.-L., Xu, X.-Y., Wang, D.-Q. & Xu, T.-T. (2006). *Acta Cryst.* **E62**, m1922–m1923.
 Hu, X., Xu, X., Xu, T. & Wang, D. (2006). *Acta Cryst.* **E62**, m2221–m2223.
 Milne, P., Ho, M. & Weaver, D. F. (1999). *J. Mol. Struct. (THEOCHEM)*, **492**, 19–28.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

supplementary materials

Acta Cryst. (2009). E65, m1426 [doi:10.1107/S160053680904313X]

Bis(5,5-diphenylhydantoinato- κN^3)(ethylenediamine)zinc(II)

X. Hu, X. Xu, D. Wang and Y. Zhang

Comment

5,5-diphenylimidazoline-2,4-dione (phenytoin) compound is a widely used drug in the treatment of epilepsy and should be an excellent ligand for transition metal complex (Milne *et al.*, 1999; Akitsu, Komorita, Kushi *et al.*, 1997; Akitsu, Einaga, 2005). We have designed and synthesized a series of complexes with 5,5-diphenylhydantoinate (Hu, Xu, Wang *et al.*, 2006). We report here the crystal structure of the title compound (I). The compound (Fig. 1) consists of $[Zn(pht)_2(en)]$ (pht=5,5-diphenylhydantoinato; en=ethylenediamine) complex neutral molecule. The Zn atom is coordinated by two nitrogen atoms from two pht ligands and two nitrogen atoms from two en ligands and is in a distorted tetrahedron ZnN_4 coordination environment. The Zn—N bond distances lie in the range of 1.9506 (18) Å to 2.057 (2) Å. There are intra- and intermolecular N—H \cdots O=C hydrogen bonds, forming a three-dimensional network in the crystal structure, Table 1.

Experimental

To a solution of pht (1.00 mmol) in methanol (10 ml) was added zinc(II) acetate tetrahydrate (0.5 mmol) and the solution of Ethylenediamine(0.5 mmol) in methanol (10 ml). Then the mixture was sealed in a 25 ml stainless steel vessel with Teflon liner and heated to 393 K for 50 h the fill rate being 80%. After cooling to room temperature, the colorless single crystals were obtained by slow evaporation from the filtrate.

Refinement

The space group was uniquely assigned from the systematic absences. All H atoms were placed at calculated positions, with N—H = 0.86–0.90 Å, and with $U_{iso}(H)$ values were set at $1.2U_{eq}$, and C—H = 0.93 Å (phenyl), 0.97 Å (methylene), respectively, and with $U_{iso}(H)$ values were set at $1.2 U_{eq}(C)$ (phenyl, methylene).

Figures




Fig. 1. The molecular structure of the title complex. Displacement ellipsoids are drawn at the 30% probability level. The H-atoms have been omitted for clarity.

Bis(5,5-diphenylhydantoinato- κN^3)(ethylenediamine)zinc(II)

Crystal data

$[Zn(C_{15}H_{11}N_2O_2)_2(C_2H_8N_2)]$

$M_r = 627.99$

$Z = 2$

$F_{000} = 652$

supplementary materials

Triclinic, $P\bar{1}$	$D_x = 1.416 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.7020 (10) \text{ \AA}$	Cell parameters from 4727 reflections
$b = 13.0520 (15) \text{ \AA}$	$\theta = 2.6\text{--}27.7^\circ$
$c = 13.2930 (16) \text{ \AA}$	$\mu = 0.88 \text{ mm}^{-1}$
$\alpha = 109.114 (2)^\circ$	$T = 298 \text{ K}$
$\beta = 109.462 (2)^\circ$	Block, colorless
$\gamma = 93.3020 (10)^\circ$	$0.50 \times 0.46 \times 0.32 \text{ mm}$
$V = 1473.1 (3) \text{ \AA}^3$	

Data collection

Bruker SMART CCD area-detector diffractometer	5124 independent reflections
Radiation source: fine-focus sealed tube	4452 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.017$
$T = 298 \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.7^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.667$, $T_{\text{max}} = 0.766$	$k = -15 \rightarrow 15$
7725 measured reflections	$l = -14 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.034$	H-atom parameters constrained
$wR(F^2) = 0.089$	$w = 1/[\sigma^2(F_o^2) + (0.0432P)^2 + 0.9132P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
5124 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
388 parameters	$\Delta\rho_{\text{max}} = 0.61 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.45 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -

factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on F^2 are statistically about twice as large as those based on *F*, and *R*- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	-0.00468 (3)	0.61290 (2)	0.37871 (2)	0.02823 (10)
N1	0.4053 (2)	0.53299 (18)	0.37370 (16)	0.0358 (5)
H1	0.4894	0.5154	0.4032	0.043*
N2	0.1898 (2)	0.59269 (15)	0.36211 (16)	0.0290 (4)
N3	-0.0141 (3)	0.92085 (15)	0.58602 (18)	0.0380 (5)
H3	0.0040	0.9910	0.6028	0.046*
N4	-0.0279 (2)	0.74051 (15)	0.49503 (17)	0.0328 (4)
N5	-0.1098 (2)	0.46617 (15)	0.36800 (17)	0.0335 (4)
H5A	-0.0432	0.4228	0.3835	0.040*
H5B	-0.1548	0.4787	0.4190	0.040*
N6	-0.1472 (2)	0.5694 (2)	0.21093 (19)	0.0480 (6)
H6A	-0.2288	0.5998	0.2067	0.058*
H6B	-0.1020	0.5906	0.1699	0.058*
O1	0.33097 (18)	0.56196 (14)	0.52547 (13)	0.0337 (4)
O2	0.1300 (2)	0.61465 (16)	0.18764 (15)	0.0455 (5)
O3	0.0303 (2)	0.86389 (14)	0.41865 (17)	0.0510 (5)
O4	-0.1104 (2)	0.66928 (13)	0.60608 (16)	0.0401 (4)
C1	0.3111 (2)	0.56151 (18)	0.42891 (19)	0.0278 (5)
C2	0.2089 (3)	0.5866 (2)	0.2636 (2)	0.0318 (5)
C3	0.3484 (3)	0.5354 (2)	0.25864 (19)	0.0319 (5)
C4	0.2992 (3)	0.4187 (2)	0.1657 (2)	0.0333 (5)
C5	0.3729 (3)	0.3350 (2)	0.1842 (2)	0.0433 (6)
H5	0.4527	0.3499	0.2528	0.052*
C6	0.3292 (4)	0.2292 (2)	0.1016 (3)	0.0537 (7)
H6	0.3807	0.1741	0.1151	0.064*
C7	0.2103 (4)	0.2046 (2)	-0.0001 (3)	0.0523 (7)
H7	0.1812	0.1334	-0.0550	0.063*
C8	0.1354 (3)	0.2864 (2)	-0.0196 (2)	0.0498 (7)
H8	0.0545	0.2705	-0.0878	0.060*
C9	0.1796 (3)	0.3926 (2)	0.0618 (2)	0.0420 (6)
H9	0.1288	0.4476	0.0470	0.050*
C10	0.4530 (3)	0.6143 (2)	0.2406 (2)	0.0325 (5)
C11	0.5446 (3)	0.7065 (2)	0.3331 (2)	0.0440 (6)
H11	0.5478	0.7163	0.4064	0.053*
C12	0.6311 (3)	0.7837 (2)	0.3181 (3)	0.0490 (7)
H12	0.6919	0.8447	0.3811	0.059*
C13	0.6278 (3)	0.7708 (2)	0.2107 (3)	0.0500 (7)
H13	0.6852	0.8233	0.2006	0.060*
C14	0.5390 (3)	0.6797 (3)	0.1181 (3)	0.0553 (8)
H14	0.5364	0.6706	0.0451	0.066*
C15	0.4527 (3)	0.6011 (2)	0.1331 (2)	0.0469 (7)
H15	0.3943	0.5391	0.0701	0.056*

supplementary materials

C16	-0.0009 (3)	0.84551 (19)	0.4940 (2)	0.0346 (6)
C17	-0.0734 (3)	0.74627 (18)	0.5818 (2)	0.0303 (5)
C18	-0.0629 (3)	0.86969 (18)	0.6538 (2)	0.0313 (5)
C19	-0.2101 (3)	0.89940 (19)	0.6608 (2)	0.0341 (5)
C20	-0.2238 (4)	1.0100 (2)	0.6886 (3)	0.0608 (9)
H20	-0.1437	1.0630	0.7039	0.073*
C21	-0.3546 (5)	1.0409 (3)	0.6938 (4)	0.0841 (13)
H21	-0.3623	1.1150	0.7134	0.101*
C22	-0.4754 (4)	0.9628 (3)	0.6699 (4)	0.0769 (11)
H22	-0.5651	0.9839	0.6709	0.092*
C23	-0.4618 (3)	0.8539 (3)	0.6447 (3)	0.0568 (8)
H23	-0.5422	0.8011	0.6295	0.068*
C24	-0.3285 (3)	0.8228 (2)	0.6420 (2)	0.0422 (6)
H24	-0.3190	0.7494	0.6274	0.051*
C25	0.0588 (3)	0.89167 (19)	0.7714 (2)	0.0357 (6)
C26	0.2070 (3)	0.9131 (2)	0.7872 (3)	0.0530 (7)
H26	0.2329	0.9217	0.7286	0.064*
C27	0.3172 (4)	0.9218 (3)	0.8887 (3)	0.0681 (10)
H27	0.4166	0.9367	0.8982	0.082*
C28	0.2816 (4)	0.9088 (3)	0.9754 (3)	0.0697 (10)
H28	0.3562	0.9137	1.0433	0.084*
C29	0.1353 (4)	0.8884 (3)	0.9616 (3)	0.0718 (10)
H29	0.1104	0.8803	1.0208	0.086*
C30	0.0239 (3)	0.8797 (3)	0.8599 (3)	0.0554 (8)
H30	-0.0753	0.8657	0.8512	0.066*
C31	-0.2203 (4)	0.4122 (3)	0.2507 (3)	0.0715 (10)
H31A	-0.2261	0.3330	0.2275	0.086*
H31B	-0.3172	0.4277	0.2499	0.086*
C32	-0.1855 (5)	0.4482 (3)	0.1696 (3)	0.0890 (14)
H32A	-0.2703	0.4216	0.0975	0.107*
H32B	-0.1024	0.4167	0.1557	0.107*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.03145 (16)	0.02810 (15)	0.03137 (16)	0.01168 (11)	0.01542 (12)	0.01390 (11)
N1	0.0317 (11)	0.0594 (13)	0.0302 (11)	0.0233 (10)	0.0153 (9)	0.0273 (10)
N2	0.0279 (10)	0.0367 (10)	0.0280 (10)	0.0121 (8)	0.0125 (8)	0.0160 (8)
N3	0.0587 (14)	0.0218 (9)	0.0486 (13)	0.0109 (9)	0.0358 (11)	0.0151 (9)
N4	0.0423 (12)	0.0242 (9)	0.0404 (11)	0.0116 (9)	0.0239 (10)	0.0127 (9)
N5	0.0362 (11)	0.0334 (10)	0.0375 (11)	0.0081 (9)	0.0163 (9)	0.0185 (9)
N6	0.0366 (13)	0.0734 (16)	0.0464 (13)	0.0160 (12)	0.0117 (11)	0.0402 (13)
O1	0.0337 (9)	0.0471 (10)	0.0257 (8)	0.0093 (7)	0.0127 (7)	0.0183 (7)
O2	0.0418 (11)	0.0743 (13)	0.0428 (10)	0.0326 (10)	0.0212 (9)	0.0405 (10)
O3	0.0859 (15)	0.0372 (10)	0.0595 (12)	0.0229 (10)	0.0542 (12)	0.0252 (9)
O4	0.0506 (11)	0.0293 (9)	0.0547 (11)	0.0117 (8)	0.0299 (9)	0.0222 (8)
C1	0.0269 (12)	0.0297 (11)	0.0285 (12)	0.0061 (9)	0.0102 (10)	0.0128 (10)
C2	0.0284 (12)	0.0412 (13)	0.0326 (13)	0.0129 (10)	0.0130 (10)	0.0193 (11)

C3	0.0291 (13)	0.0496 (14)	0.0273 (12)	0.0168 (11)	0.0129 (10)	0.0231 (11)
C4	0.0311 (13)	0.0462 (14)	0.0347 (13)	0.0105 (11)	0.0181 (11)	0.0232 (11)
C5	0.0392 (15)	0.0525 (16)	0.0471 (15)	0.0186 (13)	0.0187 (13)	0.0253 (13)
C6	0.061 (2)	0.0490 (17)	0.066 (2)	0.0235 (15)	0.0309 (17)	0.0291 (15)
C7	0.064 (2)	0.0444 (16)	0.0512 (17)	0.0022 (14)	0.0278 (16)	0.0163 (14)
C8	0.0516 (18)	0.0568 (17)	0.0396 (15)	0.0017 (14)	0.0126 (13)	0.0219 (14)
C9	0.0432 (16)	0.0499 (16)	0.0390 (14)	0.0127 (12)	0.0145 (12)	0.0240 (13)
C10	0.0271 (12)	0.0461 (14)	0.0335 (13)	0.0159 (11)	0.0146 (10)	0.0213 (11)
C11	0.0457 (16)	0.0524 (16)	0.0381 (14)	0.0149 (13)	0.0185 (13)	0.0181 (13)
C12	0.0421 (16)	0.0460 (16)	0.0527 (17)	0.0083 (13)	0.0136 (13)	0.0146 (13)
C13	0.0414 (16)	0.0540 (17)	0.067 (2)	0.0096 (13)	0.0243 (15)	0.0329 (16)
C14	0.0559 (19)	0.073 (2)	0.0485 (17)	0.0045 (16)	0.0241 (15)	0.0330 (16)
C15	0.0444 (16)	0.0629 (18)	0.0347 (14)	−0.0003 (13)	0.0147 (12)	0.0215 (13)
C16	0.0427 (15)	0.0273 (12)	0.0445 (14)	0.0128 (10)	0.0267 (12)	0.0148 (11)
C17	0.0305 (13)	0.0267 (11)	0.0373 (13)	0.0088 (10)	0.0148 (10)	0.0134 (10)
C18	0.0386 (14)	0.0249 (11)	0.0377 (13)	0.0078 (10)	0.0212 (11)	0.0131 (10)
C19	0.0419 (14)	0.0357 (13)	0.0344 (13)	0.0146 (11)	0.0207 (11)	0.0171 (11)
C20	0.068 (2)	0.0454 (16)	0.101 (3)	0.0266 (15)	0.057 (2)	0.0371 (17)
C21	0.094 (3)	0.067 (2)	0.153 (4)	0.055 (2)	0.088 (3)	0.067 (3)
C22	0.065 (2)	0.093 (3)	0.123 (3)	0.050 (2)	0.063 (2)	0.069 (3)
C23	0.0432 (17)	0.071 (2)	0.068 (2)	0.0131 (15)	0.0288 (16)	0.0306 (17)
C24	0.0461 (16)	0.0405 (14)	0.0478 (16)	0.0121 (12)	0.0248 (13)	0.0179 (12)
C25	0.0374 (14)	0.0278 (12)	0.0418 (14)	0.0053 (10)	0.0180 (12)	0.0096 (11)
C26	0.0434 (17)	0.0584 (18)	0.0517 (17)	0.0047 (14)	0.0224 (14)	0.0095 (14)
C27	0.0370 (17)	0.078 (2)	0.069 (2)	0.0039 (16)	0.0139 (16)	0.0100 (19)
C28	0.055 (2)	0.070 (2)	0.060 (2)	0.0040 (17)	−0.0032 (17)	0.0200 (18)
C29	0.062 (2)	0.100 (3)	0.0502 (19)	−0.002 (2)	0.0093 (17)	0.0381 (19)
C30	0.0393 (16)	0.080 (2)	0.0496 (17)	0.0003 (15)	0.0144 (14)	0.0311 (16)
C31	0.078 (2)	0.059 (2)	0.053 (2)	−0.0192 (18)	0.0014 (18)	0.0198 (16)
C32	0.110 (3)	0.081 (3)	0.0382 (18)	−0.027 (2)	−0.0048 (19)	0.0157 (18)

Geometric parameters (Å, °)

Zn1—N4	1.9506 (18)	C11—C12	1.381 (4)
Zn1—N2	1.9941 (19)	C11—H11	0.9300
Zn1—N5	2.0561 (19)	C12—C13	1.370 (4)
Zn1—N6	2.057 (2)	C12—H12	0.9300
N1—C1	1.349 (3)	C13—C14	1.374 (4)
N1—C3	1.455 (3)	C13—H13	0.9300
N1—H1	0.8600	C14—C15	1.392 (4)
N2—C2	1.360 (3)	C14—H14	0.9300
N2—C1	1.390 (3)	C15—H15	0.9300
N3—C16	1.346 (3)	C17—C18	1.558 (3)
N3—C18	1.457 (3)	C18—C19	1.525 (3)
N3—H3	0.8600	C18—C25	1.535 (4)
N4—C17	1.348 (3)	C19—C24	1.377 (4)
N4—C16	1.386 (3)	C19—C20	1.394 (4)
N5—C31	1.469 (4)	C20—C21	1.370 (4)
N5—H5A	0.9000	C20—H20	0.9300

supplementary materials

N5—H5B	0.9000	C21—C22	1.384 (5)
N6—C32	1.472 (4)	C21—H21	0.9300
N6—H6A	0.9000	C22—C23	1.374 (5)
N6—H6B	0.9000	C22—H22	0.9300
O1—C1	1.231 (3)	C23—C24	1.386 (4)
O2—C2	1.224 (3)	C23—H23	0.9300
O3—C16	1.226 (3)	C24—H24	0.9300
O4—C17	1.221 (3)	C25—C30	1.379 (4)
C2—C3	1.555 (3)	C25—C26	1.380 (4)
C3—C10	1.534 (3)	C26—C27	1.379 (5)
C3—C4	1.537 (3)	C26—H26	0.9300
C4—C5	1.384 (3)	C27—C28	1.365 (5)
C4—C9	1.396 (4)	C27—H27	0.9300
C5—C6	1.386 (4)	C28—C29	1.368 (5)
C5—H5	0.9300	C28—H28	0.9300
C6—C7	1.377 (4)	C29—C30	1.386 (4)
C6—H6	0.9300	C29—H29	0.9300
C7—C8	1.371 (4)	C30—H30	0.9300
C7—H7	0.9300	C31—C32	1.433 (5)
C8—C9	1.384 (4)	C31—H31A	0.9700
C8—H8	0.9300	C31—H31B	0.9700
C9—H9	0.9300	C32—H32A	0.9700
C10—C15	1.382 (3)	C32—H32B	0.9700
C10—C11	1.389 (4)		
N4—Zn1—N2	123.12 (8)	C12—C13—C14	119.5 (3)
N4—Zn1—N5	112.66 (8)	C12—C13—H13	120.2
N2—Zn1—N5	108.94 (8)	C14—C13—H13	120.2
N4—Zn1—N6	118.42 (9)	C13—C14—C15	120.3 (3)
N2—Zn1—N6	102.08 (9)	C13—C14—H14	119.8
N5—Zn1—N6	84.83 (9)	C15—C14—H14	119.8
C1—N1—C3	112.29 (18)	C10—C15—C14	120.7 (3)
C1—N1—H1	123.9	C10—C15—H15	119.7
C3—N1—H1	123.9	C14—C15—H15	119.7
C2—N2—C1	108.39 (18)	O3—C16—N3	126.6 (2)
C2—N2—Zn1	120.78 (15)	O3—C16—N4	123.3 (2)
C1—N2—Zn1	129.63 (15)	N3—C16—N4	110.1 (2)
C16—N3—C18	112.08 (19)	O4—C17—N4	126.9 (2)
C16—N3—H3	124.0	O4—C17—C18	123.8 (2)
C18—N3—H3	124.0	N4—C17—C18	109.18 (18)
C17—N4—C16	109.22 (18)	N3—C18—C19	111.23 (19)
C17—N4—Zn1	130.24 (15)	N3—C18—C25	112.6 (2)
C16—N4—Zn1	120.51 (16)	C19—C18—C25	113.46 (19)
C31—N5—Zn1	107.60 (17)	N3—C18—C17	99.11 (17)
C31—N5—H5A	110.2	C19—C18—C17	114.35 (19)
Zn1—N5—H5A	110.2	C25—C18—C17	105.10 (18)
C31—N5—H5B	110.2	C24—C19—C20	118.8 (2)
Zn1—N5—H5B	110.2	C24—C19—C18	123.3 (2)
H5A—N5—H5B	108.5	C20—C19—C18	117.9 (2)
C32—N6—Zn1	103.62 (18)	C21—C20—C19	120.3 (3)

C32—N6—H6A	111.0	C21—C20—H20	119.8
Zn1—N6—H6A	111.0	C19—C20—H20	119.8
C32—N6—H6B	111.0	C20—C21—C22	120.6 (3)
Zn1—N6—H6B	111.0	C20—C21—H21	119.7
H6A—N6—H6B	109.0	C22—C21—H21	119.7
O1—C1—N1	124.9 (2)	C23—C22—C21	119.5 (3)
O1—C1—N2	124.9 (2)	C23—C22—H22	120.3
N1—C1—N2	110.25 (19)	C21—C22—H22	120.3
O2—C2—N2	127.1 (2)	C22—C23—C24	120.0 (3)
O2—C2—C3	123.4 (2)	C22—C23—H23	120.0
N2—C2—C3	109.58 (18)	C24—C23—H23	120.0
N1—C3—C10	112.7 (2)	C19—C24—C23	120.7 (3)
N1—C3—C4	111.98 (19)	C19—C24—H24	119.7
C10—C3—C4	114.11 (18)	C23—C24—H24	119.7
N1—C3—C2	98.85 (17)	C30—C25—C26	118.3 (3)
C10—C3—C2	108.59 (19)	C30—C25—C18	121.1 (2)
C4—C3—C2	109.47 (19)	C26—C25—C18	120.3 (2)
C5—C4—C9	117.7 (2)	C27—C26—C25	120.7 (3)
C5—C4—C3	120.1 (2)	C27—C26—H26	119.6
C9—C4—C3	122.2 (2)	C25—C26—H26	119.6
C4—C5—C6	120.8 (3)	C28—C27—C26	120.5 (3)
C4—C5—H5	119.6	C28—C27—H27	119.7
C6—C5—H5	119.6	C26—C27—H27	119.7
C7—C6—C5	120.8 (3)	C27—C28—C29	119.5 (3)
C7—C6—H6	119.6	C27—C28—H28	120.2
C5—C6—H6	119.6	C29—C28—H28	120.2
C8—C7—C6	119.3 (3)	C28—C29—C30	120.3 (3)
C8—C7—H7	120.4	C28—C29—H29	119.9
C6—C7—H7	120.4	C30—C29—H29	119.9
C7—C8—C9	120.3 (3)	C25—C30—C29	120.6 (3)
C7—C8—H8	119.8	C25—C30—H30	119.7
C9—C8—H8	119.8	C29—C30—H30	119.7
C8—C9—C4	121.1 (3)	C32—C31—N5	112.7 (3)
C8—C9—H9	119.4	C32—C31—H31A	109.1
C4—C9—H9	119.4	N5—C31—H31A	109.1
C15—C10—C11	118.0 (2)	C32—C31—H31B	109.1
C15—C10—C3	122.0 (2)	N5—C31—H31B	109.1
C11—C10—C3	119.8 (2)	H31A—C31—H31B	107.8
C12—C11—C10	121.1 (3)	C31—C32—N6	111.7 (3)
C12—C11—H11	119.4	C31—C32—H32A	109.3
C10—C11—H11	119.4	N6—C32—H32A	109.3
C13—C12—C11	120.3 (3)	C31—C32—H32B	109.3
C13—C12—H12	119.8	N6—C32—H32B	109.3
C11—C12—H12	119.8	H32A—C32—H32B	107.9
N4—Zn1—N2—C2	-118.65 (18)	C3—C10—C11—C12	-174.7 (2)
N5—Zn1—N2—C2	106.10 (18)	C10—C11—C12—C13	0.2 (4)
N6—Zn1—N2—C2	17.5 (2)	C11—C12—C13—C14	-0.8 (4)
N4—Zn1—N2—C1	75.4 (2)	C12—C13—C14—C15	0.1 (5)
N5—Zn1—N2—C1	-59.8 (2)	C11—C10—C15—C14	-1.7 (4)

supplementary materials

N6—Zn1—N2—C1	-148.45 (19)	C3—C10—C15—C14	174.0 (3)
N2—Zn1—N4—C17	-118.7 (2)	C13—C14—C15—C10	1.1 (5)
N5—Zn1—N4—C17	15.1 (2)	C18—N3—C16—O3	175.9 (3)
N6—Zn1—N4—C17	111.7 (2)	C18—N3—C16—N4	-4.2 (3)
N2—Zn1—N4—C16	63.7 (2)	C17—N4—C16—O3	-174.0 (3)
N5—Zn1—N4—C16	-162.46 (18)	Zn1—N4—C16—O3	4.0 (4)
N6—Zn1—N4—C16	-65.9 (2)	C17—N4—C16—N3	6.1 (3)
N4—Zn1—N5—C31	120.3 (2)	Zn1—N4—C16—N3	-175.88 (17)
N2—Zn1—N5—C31	-99.4 (2)	C16—N4—C17—O4	178.0 (2)
N6—Zn1—N5—C31	1.6 (2)	Zn1—N4—C17—O4	0.2 (4)
N4—Zn1—N6—C32	-137.5 (2)	C16—N4—C17—C18	-5.5 (3)
N2—Zn1—N6—C32	83.8 (2)	Zn1—N4—C17—C18	176.76 (16)
N5—Zn1—N6—C32	-24.5 (2)	C16—N3—C18—C19	-119.9 (2)
C3—N1—C1—O1	177.2 (2)	C16—N3—C18—C25	111.5 (2)
C3—N1—C1—N2	-4.1 (3)	C16—N3—C18—C17	0.8 (3)
C2—N2—C1—O1	177.2 (2)	O4—C17—C18—N3	179.6 (2)
Zn1—N2—C1—O1	-15.6 (3)	N4—C17—C18—N3	2.9 (2)
C2—N2—C1—N1	-1.5 (3)	O4—C17—C18—C19	-62.1 (3)
Zn1—N2—C1—N1	165.76 (16)	N4—C17—C18—C19	121.2 (2)
C1—N2—C2—O2	-174.8 (2)	O4—C17—C18—C25	63.0 (3)
Zn1—N2—C2—O2	16.6 (4)	N4—C17—C18—C25	-113.7 (2)
C1—N2—C2—C3	6.1 (3)	N3—C18—C19—C24	132.8 (2)
Zn1—N2—C2—C3	-162.49 (15)	C25—C18—C19—C24	-99.0 (3)
C1—N1—C3—C10	121.6 (2)	C17—C18—C19—C24	21.6 (3)
C1—N1—C3—C4	-108.2 (2)	N3—C18—C19—C20	-47.8 (3)
C1—N1—C3—C2	7.1 (3)	C25—C18—C19—C20	80.4 (3)
O2—C2—C3—N1	173.0 (2)	C17—C18—C19—C20	-159.1 (2)
N2—C2—C3—N1	-7.9 (2)	C24—C19—C20—C21	-1.9 (5)
O2—C2—C3—C10	55.4 (3)	C18—C19—C20—C21	178.7 (3)
N2—C2—C3—C10	-125.5 (2)	C19—C20—C21—C22	-0.9 (6)
O2—C2—C3—C4	-69.8 (3)	C20—C21—C22—C23	2.2 (7)
N2—C2—C3—C4	109.3 (2)	C21—C22—C23—C24	-0.8 (6)
N1—C3—C4—C5	-32.8 (3)	C20—C19—C24—C23	3.4 (4)
C10—C3—C4—C5	96.7 (3)	C18—C19—C24—C23	-177.2 (2)
C2—C3—C4—C5	-141.3 (2)	C22—C23—C24—C19	-2.1 (5)
N1—C3—C4—C9	146.4 (2)	N3—C18—C25—C30	160.6 (2)
C10—C3—C4—C9	-84.1 (3)	C19—C18—C25—C30	33.1 (3)
C2—C3—C4—C9	37.8 (3)	C17—C18—C25—C30	-92.6 (3)
C9—C4—C5—C6	0.3 (4)	N3—C18—C25—C26	-26.0 (3)
C3—C4—C5—C6	179.5 (2)	C19—C18—C25—C26	-153.5 (2)
C4—C5—C6—C7	-0.8 (4)	C17—C18—C25—C26	80.9 (3)
C5—C6—C7—C8	0.4 (4)	C30—C25—C26—C27	0.2 (4)
C6—C7—C8—C9	0.5 (4)	C18—C25—C26—C27	-173.4 (3)
C7—C8—C9—C4	-1.0 (4)	C25—C26—C27—C28	0.4 (5)
C5—C4—C9—C8	0.6 (4)	C26—C27—C28—C29	-0.9 (6)
C3—C4—C9—C8	-178.6 (2)	C27—C28—C29—C30	0.8 (6)
N1—C3—C10—C15	155.5 (2)	C26—C25—C30—C29	-0.4 (5)
C4—C3—C10—C15	26.3 (3)	C18—C25—C30—C29	173.2 (3)
C2—C3—C10—C15	-96.1 (3)	C28—C29—C30—C25	-0.1 (6)

N1—C3—C10—C11	-28.9 (3)	Zn1—N5—C31—C32	23.6 (4)
C4—C3—C10—C11	-158.1 (2)	N5—C31—C32—N6	-47.9 (5)
C2—C3—C10—C11	79.5 (3)	Zn1—N6—C32—C31	45.3 (4)
C15—C10—C11—C12	1.0 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N6—H6B \cdots O2	0.90	2.18	2.858 (3)	132
C9—H9 \cdots O2	0.93	2.36	2.992 (3)	125
C11—H11 \cdots N1	0.93	2.52	2.860 (4)	102
C24—H24 \cdots O4	0.93	2.39	3.042 (4)	127
C26—H26 \cdots N3	0.93	2.51	2.854 (4)	102
N1—H1 \cdots O1 ⁱ	0.86	2.16	3.008 (3)	167
N3—H3 \cdots O3 ⁱⁱ	0.86	2.04	2.843 (3)	156
N5—H5A \cdots O4 ⁱⁱⁱ	0.90	1.97	2.855 (3)	170
N5—H5B \cdots O1 ⁱⁱⁱ	0.90	2.18	3.003 (3)	152

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, -y+2, -z+1$; (iii) $-x, -y+1, -z+1$.

Fig. 1

