

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

{4-Bromo-2-[3-(diethylammonio)propyliminomethyl]phenolato}diiodidozinc(II) methanol solvate

Xue-Wen Zhu,* Zhi-Gang Yin, Xu-Zhao Yang, Gang-Sen Li and Chun-Xia Zhang

Key Laboratory of Surface and Interface Science of Henan, School of Material & Chemical Engineering, Zhengzhou University of Light Industry, Zhengzhou 450002, People's Republic of China

Correspondence e-mail: xuewen-zhu@163.com


Received 10 September 2009; accepted 22 September 2009

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; disorder in main residue; R factor = 0.048; wR factor = 0.105; data-to-parameter ratio = 20.7.

In the title complex, $[\text{ZnI}_2(\text{C}_{14}\text{H}_{21}\text{BrN}_2\text{O})]\cdot\text{CH}_3\text{OH}$, the asymmetric unit consists of a mononuclear zinc(II) complex molecule and a methanol solvent molecule. The compound was derived from the zwitterionic form of the Schiff base 4-bromo-2-[3-(diethylamino)propyliminomethyl]phenol. The Zn^{II} atom is four-coordinated by the imine N and phenolate O atoms of the Schiff base ligand and by two iodide ions in a distorted tetrahedral coordination. In the crystal structure, the methanol molecules are linked to the Schiff base molecules through $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. One I atom is disordered over two positions in a 0.702 (19):0.298 (19) ratio.

Related literature

For background to the chemistry of Schiff base complexes, see: Ali *et al.* (2008); Biswas *et al.* (2008); Chen *et al.* (2008); Darensbourg & Frantz (2007); Habibi *et al.* (2007); Kawamoto *et al.* (2008); Lipscomb & Sträter (1996); Tomat *et al.* (2007); Wu *et al.* (2008); Yuan *et al.* (2007). For related structures, see: Zhu (2008); Zhu & Yang (2008*a,b,c*); Qiu (2006*a,b*); Wei *et al.* (2007); Zhu *et al.* (2007). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

 $[\text{ZnI}_2(\text{C}_{14}\text{H}_{21}\text{BrN}_2\text{O})]\cdot\text{CH}_4\text{O}$
 $M_r = 664.45$

 Monoclinic, $P2_1/n$
 $a = 10.869$ (2) Å

 $b = 17.562$ (3) Å

 $c = 11.377$ (2) Å

 $\beta = 94.358$ (3)°

 $V = 2165.4$ (7) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 5.84$ mm⁻¹
 $T = 298$ K

 $0.20 \times 0.20 \times 0.17$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Sheldrick, 2004)

 $T_{\text{min}} = 0.388$, $T_{\text{max}} = 0.437$

14106 measured reflections

4664 independent reflections

 3499 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.105$
 $S = 1.06$

4664 reflections

225 parameters

8 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.75$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.85$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Zn1—O1	1.958 (4)	Zn1—I1	2.5627 (9)
Zn1—N1	2.032 (5)	Zn1—I2	2.5768 (18)
Zn1—I2'	2.545 (6)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O2}-\text{H2}\cdots\text{O1}^{\text{i}}$	0.82	1.86	2.640 (6)	158
$\text{N2}-\text{H2A}\cdots\text{O2}$	0.91 (6)	1.81 (6)	2.716 (7)	173 (8)

 Symmetry code: (i) $x + 1, y, z$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2241).

References

- Ali, H. M., Mohamed Mustafa, M. I., Rizal, M. R. & Ng, S. W. (2008). *Acta Cryst.* **E64**, m718–m719.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Biswas, C., Drew, M. G. B. & Ghosh, A. (2008). *Inorg. Chem.* **47**, 4513–4519.
- Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, Z., Morimoto, H., Matsunaga, S. & Shibasaki, M. (2008). *J. Am. Chem. Soc.* **130**, 2170–2171.
- Darensbourg, D. J. & Frantz, E. B. (2007). *Inorg. Chem.* **46**, 5967–5978.

- Habibi, M. H., Askari, E., Chantrapromma, S. & Fun, H.-K. (2007). *Acta Cryst. E* **63**, m2905–m2906.
- Kawamoto, T., Nishiwaki, M., Tsunekawa, Y., Nozaki, K. & Konno, T. (2008). *Inorg. Chem.* **47**, 3095–3104.
- Lipscomb, W. N. & Sträter, N. (1996). *Chem. Rev.* **96**, 2375–2434.
- Qiu, X.-Y. (2006a). *Acta Cryst. E* **62**, m717–m718.
- Qiu, X.-Y. (2006b). *Acta Cryst. E* **62**, m2173–m2174.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tomat, E., Cuesta, L., Lynch, V. M. & Sessler, J. L. (2007). *Inorg. Chem.* **46**, 6224–6226.
- Wei, Y.-J., Wang, F.-W. & Zhu, Q.-Y. (2007). *Acta Cryst. E* **63**, m654–m655.
- Wu, J.-C., Liu, S.-X., Keene, T. D., Neels, A., Mereacre, V., Powell, A. K. & Decurtins, S. (2008). *Inorg. Chem.* **47**, 3452–3459.
- Yuan, M., Zhao, F., Zhang, W., Wang, Z.-M. & Gao, S. (2007). *Inorg. Chem.* **46**, 11235–11242.
- Zhu, X.-W. (2008). *Acta Cryst. E* **64**, m1456–m1457.
- Zhu, Q.-Y., Wei, Y.-J. & Wang, F.-W. (2007). *Acta Cryst. E* **63**, m1431–m1432.
- Zhu, X.-W. & Yang, X.-Z. (2008a). *Acta Cryst. E* **64**, m1090–m1091.
- Zhu, X.-W. & Yang, X.-Z. (2008b). *Acta Cryst. E* **64**, m1092–m1093.
- Zhu, X.-W. & Yang, X.-Z. (2008c). *Acta Cryst. E* **64**, m1094–m1095.

supplementary materials

Acta Cryst. (2009). E65, m1293-m1294 [doi:10.1107/S1600536809038446]

{4-Bromo-2-[3-(diethylammonio)propyliminomethyl]phenolato}diiodidozinc(II) methanol solvate

X.-W. Zhu, Z.-G. Yin, X.-Z. Yang, G.-S. Li and C.-X. Zhang

Comment

Schiff bases have widely been used as versatile ligands in coordination chemistry (Biswas *et al.*, 2008; Wu *et al.*, 2008; Kawamoto *et al.*, 2008; Ali *et al.*, 2008; Habibi *et al.*, 2007), and their metal complexes are of great interest in many fields (Chen *et al.*, 2008; Yuan *et al.*, 2007; Tomat *et al.*, 2007; Darensbourg & Frantz, 2007). Zinc(II) is an important element in biological systems and functions as the active site of hydrolytic enzymes, such as carboxypeptidase and carbonic anhydrase where it is in a hard-donor coordination environment of nitrogen and oxygen ligands (Lipscomb & Sträter, 1996). Recently, we have reported a few Schiff base zinc complexes (Zhu, 2008; Zhu & Yang, 2008a,b,c). In this paper, the title new zinc(II) complex, Fig. 1, is reported.

The complex consists of a mononuclear zinc(II) complex molecule and a methanol molecule. The Zn^{II} atom is four-coordinated by the imine N and phenolate O atoms of the zwitterionic form of the Schiff base ligand, and by two I⁻ ions, in a distorted tetrahedral coordination. The coordinate bond lengths (Table 1) are typical and comparable to the corresponding values observed in the Schiff base zinc complexes we reported previously and other similar Schiff base zinc complexes (Zhu *et al.*, 2007; Wei *et al.*, 2007; Qiu, 2006a,b). I2 atom is disordered over two positions [0.702(19)/0.298 (19)].

In the crystal structure, the methanol molecules are linked to the Schiff base molecules through O—H \cdots O and N—H \cdots O hydrogen bonds generating a graph-set motif C²₂(10) chain along [100] direction (Table 2, Fig. 2). (Bernstein *et al.*, 1995)

Experimental

The Schiff base compound was prepared by the condensation of equimolar amounts of 5-bromosalicylaldehyde with *N,N*-diethylpropane-1,3-diamine in a methanol solution. The complex was prepared by the following method. To an anhydrous methanol solution (5 ml) of ZnI₂ (31.9 mg, 0.1 mmol) was added a methanol solution (10 ml) of the Schiff base compound (31.3 mg, 0.1 mmol) with stirring. The mixture was stirred for 30 min at room temperature and filtered. Upon keeping the filtrate in air for a few days, colorless block-shaped crystals were formed.

Refinement

H2A was located from a difference Fourier map and refined isotropically, with N—H distance restrained to 0.90 (1) Å. Other H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances in the range 0.93–0.97 Å, O—H distance of 0.82 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{methyl C and O})$. The I2 atom is disordered over two distinct sites with occupancies of 0.702 (2) and 0.298 (2), respectively.

Figures




Fig. 1. The molecular structure of the title complex, with ellipsoids drawn at the 30% probability level.




Fig. 2. The crystal packing of the title complex.

{4-Bromo-2-[3-(diethylammonio)propyliminomethyl]phenolato}diiodidozinc(II) methanol solvate

Crystal data

[ZnI₂(C₁₄H₂₁BrN₂O)]·CH₄O

M_r = 664.45

Monoclinic, *P*2₁/*n*

Hall symbol: -*P* 2₁ *n*

a = 10.869 (2) Å

b = 17.562 (3) Å

c = 11.377 (2) Å

β = 94.358 (3)°

V = 2165.4 (7) Å³

Z = 4

*F*₀₀₀ = 1264

D_x = 2.038 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 3253 reflections

θ = 2.2–24.5°

μ = 5.84 mm⁻¹

T = 298 K

Block, colorless

0.20 × 0.20 × 0.17 mm

Data collection

Bruker APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 298 K

ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 2004)

T_{min} = 0.388, *T_{max}* = 0.437

14106 measured reflections

4664 independent reflections

3499 reflections with *I* > 2σ(*I*)

R_{int} = 0.041

θ_{max} = 27.0°

θ_{min} = 2.1°

h = -13→13

k = -22→22

l = -14→13

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.048$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.105$	$w = 1/[\sigma^2(F_o^2) + (0.0298P)^2 + 7.4538P]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
4664 reflections	$(\Delta/\sigma)_{\max} = 0.001$
225 parameters	$\Delta\rho_{\max} = 0.75 \text{ e } \text{\AA}^{-3}$
8 restraints	$\Delta\rho_{\min} = -0.85 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.17105 (6)	0.26943 (4)	-0.06556 (7)	0.0380 (2)	
I1	0.13989 (4)	0.14098 (3)	-0.17051 (4)	0.05239 (16)	
I2	0.1766 (2)	0.39343 (11)	-0.1861 (2)	0.0541 (8)	0.702 (19)
I2'	0.1771 (8)	0.3793 (9)	-0.2109 (16)	0.118 (2)	0.298 (19)
Br1	0.19879 (8)	0.49547 (5)	0.46762 (7)	0.0630 (2)	
O1	0.0560 (4)	0.2904 (3)	0.0548 (4)	0.0443 (11)	
O2	0.8239 (4)	0.2628 (3)	-0.0180 (5)	0.0546 (13)	
H2	0.8982	0.2714	-0.0142	0.082*	
N1	0.3188 (4)	0.2626 (3)	0.0547 (4)	0.0311 (11)	
N2	0.6505 (5)	0.3665 (3)	-0.0945 (5)	0.0383 (12)	
C1	0.2134 (5)	0.3368 (3)	0.1971 (5)	0.0301 (13)	
C2	0.0899 (5)	0.3330 (4)	0.1466 (5)	0.0338 (14)	
C3	0.0011 (6)	0.3764 (4)	0.2014 (6)	0.0414 (16)	
H3	-0.0810	0.3733	0.1723	0.050*	
C4	0.0311 (6)	0.4231 (4)	0.2955 (6)	0.0431 (16)	
H4	-0.0297	0.4516	0.3284	0.052*	

supplementary materials

C5	0.1523 (6)	0.4276 (4)	0.3417 (6)	0.0400 (15)
C6	0.2417 (6)	0.3841 (3)	0.2958 (5)	0.0355 (14)
H6	0.3220	0.3857	0.3302	0.043*
C7	0.3161 (5)	0.2950 (3)	0.1550 (5)	0.0340 (14)
H7	0.3870	0.2914	0.2059	0.041*
C8	0.4322 (5)	0.2223 (3)	0.0264 (6)	0.0369 (14)
H8A	0.4115	0.1707	0.0018	0.044*
H8B	0.4888	0.2195	0.0965	0.044*
C9	0.4942 (5)	0.2625 (3)	-0.0704 (6)	0.0349 (14)
H9A	0.4367	0.2659	-0.1397	0.042*
H9B	0.5644	0.2326	-0.0911	0.042*
C10	0.5383 (5)	0.3429 (4)	-0.0348 (6)	0.0395 (15)
H10A	0.4722	0.3789	-0.0547	0.047*
H10B	0.5568	0.3446	0.0499	0.047*
C11	0.6305 (6)	0.3689 (4)	-0.2268 (6)	0.0506 (18)
H11A	0.5991	0.3198	-0.2544	0.061*
H11B	0.7096	0.3766	-0.2591	0.061*
C12	0.5423 (8)	0.4303 (5)	-0.2747 (8)	0.074 (3)
H12A	0.4639	0.4237	-0.2425	0.111*
H12B	0.5320	0.4265	-0.3590	0.111*
H12C	0.5753	0.4794	-0.2528	0.111*
C13	0.7034 (6)	0.4385 (4)	-0.0402 (7)	0.0545 (19)
H13A	0.7097	0.4334	0.0450	0.065*
H13B	0.6478	0.4804	-0.0608	0.065*
C14	0.8299 (7)	0.4567 (5)	-0.0810 (8)	0.064 (2)
H14A	0.8834	0.4137	-0.0666	0.096*
H14B	0.8637	0.5000	-0.0384	0.096*
H14C	0.8225	0.4679	-0.1638	0.096*
C15	0.7989 (8)	0.2122 (6)	0.0666 (8)	0.075 (3)
H15A	0.8652	0.1762	0.0773	0.113*
H15B	0.7236	0.1858	0.0433	0.113*
H15C	0.7900	0.2387	0.1393	0.113*
H2A	0.705 (6)	0.329 (3)	-0.073 (7)	0.080*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0302 (4)	0.0493 (4)	0.0342 (4)	0.0016 (3)	0.0001 (3)	-0.0028 (3)
I1	0.0523 (3)	0.0513 (3)	0.0513 (3)	0.0029 (2)	-0.0108 (2)	-0.0073 (2)
I2	0.0678 (10)	0.0487 (8)	0.0474 (9)	0.0168 (6)	0.0150 (8)	0.0091 (5)
I2'	0.130 (4)	0.100 (4)	0.115 (5)	-0.016 (3)	-0.036 (3)	0.035 (4)
Br1	0.0832 (6)	0.0568 (5)	0.0503 (5)	0.0002 (4)	0.0141 (4)	-0.0192 (4)
O1	0.026 (2)	0.065 (3)	0.042 (3)	-0.005 (2)	0.0035 (19)	-0.009 (2)
O2	0.032 (2)	0.059 (3)	0.071 (4)	-0.006 (2)	-0.006 (2)	0.006 (3)
N1	0.024 (2)	0.035 (3)	0.035 (3)	0.0020 (19)	0.005 (2)	-0.002 (2)
N2	0.034 (3)	0.040 (3)	0.042 (3)	-0.003 (2)	0.009 (2)	0.002 (3)
C1	0.025 (3)	0.032 (3)	0.033 (3)	-0.003 (2)	0.009 (2)	0.003 (3)
C2	0.029 (3)	0.041 (4)	0.032 (3)	-0.002 (2)	0.010 (3)	0.005 (3)

C3	0.030 (3)	0.048 (4)	0.047 (4)	0.005 (3)	0.010 (3)	0.009 (3)
C4	0.051 (4)	0.040 (4)	0.041 (4)	0.008 (3)	0.020 (3)	0.008 (3)
C5	0.055 (4)	0.034 (3)	0.032 (4)	0.000 (3)	0.011 (3)	-0.003 (3)
C6	0.038 (3)	0.042 (4)	0.027 (3)	-0.003 (3)	0.006 (3)	-0.002 (3)
C7	0.024 (3)	0.041 (3)	0.036 (4)	-0.003 (2)	-0.003 (3)	0.005 (3)
C8	0.030 (3)	0.032 (3)	0.049 (4)	0.004 (2)	0.006 (3)	-0.003 (3)
C9	0.024 (3)	0.038 (3)	0.043 (4)	0.003 (2)	0.007 (3)	-0.011 (3)
C10	0.035 (3)	0.045 (4)	0.040 (4)	0.000 (3)	0.013 (3)	-0.006 (3)
C11	0.049 (4)	0.058 (5)	0.045 (4)	-0.006 (3)	0.011 (3)	0.003 (4)
C12	0.070 (5)	0.076 (6)	0.074 (6)	-0.002 (4)	-0.003 (5)	0.026 (5)
C13	0.051 (4)	0.043 (4)	0.069 (5)	-0.009 (3)	0.005 (4)	-0.007 (4)
C14	0.054 (4)	0.061 (5)	0.077 (6)	-0.019 (4)	0.008 (4)	0.011 (4)
C15	0.055 (5)	0.098 (7)	0.071 (6)	-0.005 (5)	-0.002 (4)	0.008 (5)

Geometric parameters (Å, °)

Zn1—O1	1.958 (4)	C7—H7	0.9300
Zn1—N1	2.032 (5)	C8—C9	1.510 (9)
Zn1—I2'	2.545 (6)	C8—H8A	0.9700
Zn1—I1	2.5627 (9)	C8—H8B	0.9700
Zn1—I2	2.5768 (18)	C9—C10	1.536 (8)
Br1—C5	1.902 (6)	C9—H9A	0.9700
O1—C2	1.315 (7)	C9—H9B	0.9700
O2—C15	1.353 (9)	C10—H10A	0.9700
O2—H2	0.8200	C10—H10B	0.9700
N1—C7	1.278 (7)	C11—C12	1.516 (10)
N1—C8	1.478 (7)	C11—H11A	0.9700
N2—C10	1.499 (8)	C11—H11B	0.9700
N2—C13	1.503 (8)	C12—H12A	0.9600
N2—C11	1.506 (9)	C12—H12B	0.9600
N2—H2A	0.91 (6)	C12—H12C	0.9600
C1—C6	1.412 (8)	C13—C14	1.519 (9)
C1—C2	1.421 (8)	C13—H13A	0.9700
C1—C7	1.447 (8)	C13—H13B	0.9700
C2—C3	1.411 (8)	C14—H14A	0.9600
C3—C4	1.368 (9)	C14—H14B	0.9600
C3—H3	0.9300	C14—H14C	0.9600
C4—C5	1.383 (9)	C15—H15A	0.9600
C4—H4	0.9300	C15—H15B	0.9600
C5—C6	1.370 (8)	C15—H15C	0.9600
C6—H6	0.9300		
O1—Zn1—N1	93.11 (18)	N1—C8—H8B	109.4
O1—Zn1—I2'	111.2 (5)	C9—C8—H8B	109.4
N1—Zn1—I2'	115.1 (3)	H8A—C8—H8B	108.0
O1—Zn1—I1	115.00 (13)	C8—C9—C10	112.7 (5)
N1—Zn1—I1	109.36 (14)	C8—C9—H9A	109.1
I2'—Zn1—I1	111.9 (5)	C10—C9—H9A	109.1
O1—Zn1—I2	104.79 (15)	C8—C9—H9B	109.1
N1—Zn1—I2	111.03 (15)	C10—C9—H9B	109.1

supplementary materials

I2'—Zn1—I2	8.4 (5)	H9A—C9—H9B	107.8
I1—Zn1—I2	120.23 (7)	N2—C10—C9	112.5 (5)
C2—O1—Zn1	120.5 (4)	N2—C10—H10A	109.1
C15—O2—H2	109.5	C9—C10—H10A	109.1
C7—N1—C8	118.9 (5)	N2—C10—H10B	109.1
C7—N1—Zn1	120.3 (4)	C9—C10—H10B	109.1
C8—N1—Zn1	120.7 (4)	H10A—C10—H10B	107.8
C10—N2—C13	110.2 (5)	N2—C11—C12	114.7 (6)
C10—N2—C11	113.6 (5)	N2—C11—H11A	108.6
C13—N2—C11	114.2 (5)	C12—C11—H11A	108.6
C10—N2—H2A	103 (5)	N2—C11—H11B	108.6
C13—N2—H2A	106 (5)	C12—C11—H11B	108.6
C11—N2—H2A	109 (5)	H11A—C11—H11B	107.6
C6—C1—C2	119.4 (5)	C11—C12—H12A	109.5
C6—C1—C7	115.8 (5)	C11—C12—H12B	109.5
C2—C1—C7	124.9 (5)	H12A—C12—H12B	109.5
O1—C2—C3	119.9 (5)	C11—C12—H12C	109.5
O1—C2—C1	123.1 (5)	H12A—C12—H12C	109.5
C3—C2—C1	116.9 (6)	H12B—C12—H12C	109.5
C4—C3—C2	122.6 (6)	N2—C13—C14	112.2 (6)
C4—C3—H3	118.7	N2—C13—H13A	109.2
C2—C3—H3	118.7	C14—C13—H13A	109.2
C3—C4—C5	119.7 (6)	N2—C13—H13B	109.2
C3—C4—H4	120.2	C14—C13—H13B	109.2
C5—C4—H4	120.2	H13A—C13—H13B	107.9
C6—C5—C4	120.4 (6)	C13—C14—H14A	109.5
C6—C5—Br1	118.8 (5)	C13—C14—H14B	109.5
C4—C5—Br1	120.8 (5)	H14A—C14—H14B	109.5
C5—C6—C1	120.9 (6)	C13—C14—H14C	109.5
C5—C6—H6	119.5	H14A—C14—H14C	109.5
C1—C6—H6	119.5	H14B—C14—H14C	109.5
N1—C7—C1	126.3 (5)	O2—C15—H15A	109.5
N1—C7—H7	116.9	O2—C15—H15B	109.5
C1—C7—H7	116.9	H15A—C15—H15B	109.5
N1—C8—C9	111.1 (5)	O2—C15—H15C	109.5
N1—C8—H8A	109.4	H15A—C15—H15C	109.5
C9—C8—H8A	109.4	H15B—C15—H15C	109.5

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$O2-H2\cdots O1^i$	0.82	1.86	2.640 (6)	158
$N2-H2A\cdots O2$	0.91 (6)	1.81 (6)	2.716 (7)	173 (8)

Symmetry codes: (i) $x+1, y, z$.

Fig. 1




Fig. 2

