

1-Benzyl-1*H*-benzimidazole

Gang Lei* and Li Zhou

School of Chemistry and Chemical Engineering, China West Normal University,
 Nanchong 637002, People's Republic of China
 Correspondence e-mail: leigang307@yahoo.com.cn


Received 25 September 2009; accepted 25 September 2009

Key indicators: single-crystal X-ray study; $T = 93$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.038; wR factor = 0.093; data-to-parameter ratio = 16.6.

In the title molecule, $C_{14}H_{12}N_2$, the benzimidazole ring system is essentially planar (r.m.s. deviation = 0.024 Å). The dihedral angle between the imidazole ring and the benzyl ring is $85.77(4)^\circ$. In the crystal, molecules are linked into chains along the a axis by $C-H \cdots N$ hydrogen bonds. In addition, the packing is stabilized by $C-H \cdots \pi$ interactions involving both six-membered rings.

Related literature

For general background to benzimidazole derivatives, see: Ansari & Lal (2009). For the synthesis, see: Hayat *et al.* (2001).



Experimental

Crystal data

$C_{14}H_{12}N_2$
 $M_r = 208.26$
 Monoclinic, $P2_1/n$

$a = 6.2265(10)$ Å
 $b = 8.1740(13)$ Å
 $c = 20.975(4)$ Å

$\beta = 97.839(2)^\circ$
 $V = 1057.5(3)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.08$ mm⁻¹
 $T = 93$ K
 $0.57 \times 0.50 \times 0.37$ mm

Data collection

Rigaku SPIDER diffractometer
 Absorption correction: none
 8358 measured reflections

2412 independent reflections
 2198 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.093$
 $S = 1.00$
 2412 reflections

145 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.21$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.25$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
C10—H10B···N2 ⁱ	0.99	2.50	3.4890 (14)	173
C7—H7···Cg1 ⁱⁱ	0.95	2.66	3.5220 (1)	151
C13—H13···Cg2 ⁱⁱⁱ	0.95	2.80	3.5660 (3)	139

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x + 1, -y - 1, -z + 1$; (iii) $x, y + 1, z$. Cg1 and Cg2 are the centroids of the C11–C16 and C4–C9 rings, respectively.

Data collection: *RAPID-AUTO* (Rigaku, 2004); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank the Centre for Testing and Analysis, Cheng Du Branch of the Chinese Academy of Sciences, for analytical support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2926).

References

- Ansari, K. F. & Lal, C. (2009). *Eur. J. Med. Chem.* **44**, 4028–4033.
- Hayat, S., Atta-ur-Rahman, Choudhary, M. I., Khan, K. M., Schumann, W. & Bayer, E. (2001). *Tetrahedron*, **57**, 9951–9957.
- Rigaku (2004). *RAPID-AUTO*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supporting information

Acta Cryst. (2009). E65, o2613 [https://doi.org/10.1107/S1600536809039051]

1-Benzyl-1*H*-benzimidazole

Gang Lei and Li Zhou

S1. Comment

Benzimidazole derivatives are a class of important compounds which exhibit antimicrobial activity (Ansari & Lal, 2009). Here, we report the crystal structure of the title compound.

Bond lengths and angles in the title molecule are normal. The benzimidazole ring system is planar, with a maximum deviation of 0.035 (2) Å for atom C4. The imidazole ring and benzene ring in benzyl group are almost mutually perpendicular, with a dihedral angle of 85.77 (4)% (Fig. 1). The crystal packing is stabilized by C—H···N hydrogen bonds and C—H···π interactions ($Cg1$ is the centroid of the C11–C16 ring and $Cg2$ is the centroid of the C4—C9 ring) (Table 1).

S2. Experimental

The title compound was synthesized according to the method reported in the literature (Hayat *et al.*, 2001). Colourless single crystals suitable for X-ray diffraction were obtained by slow evaporation of a methanol solution.

S3. Refinement

All H atoms were placed in calculated positions, with C–H = 0.95 or 0.99 Å, and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atomic numbering.

1-Benzyl-1*H*-benzimidazole*Crystal data*

C₁₄H₁₂N₂
 $M_r = 208.26$
 Monoclinic, $P2_1/n$
 Hall symbol: -P 2yn
 $a = 6.2265 (10)$ Å
 $b = 8.1740 (13)$ Å
 $c = 20.975 (4)$ Å
 $\beta = 97.839 (2)^\circ$
 $V = 1057.5 (3)$ Å³
 $Z = 4$

$F(000) = 440$
 $D_x = 1.308 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 3307 reflections
 $\theta = 3.2\text{--}27.5^\circ$
 $\mu = 0.08 \text{ mm}^{-1}$
 $T = 93$ K
 Block, colourless
 $0.57 \times 0.50 \times 0.37$ mm

Data collection

Rigaku SPIDER
 diffractometer
 Radiation source: Rotating Anode
 Graphite monochromator
 ω scans
 8358 measured reflections
 2412 independent reflections

2198 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$
 $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.2^\circ$
 $h = -8 \rightarrow 8$
 $k = -10 \rightarrow 10$
 $l = -27 \rightarrow 27$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.093$
 $S = 1.00$
 2412 reflections
 145 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0482P)^2 + 0.296P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.21 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.25 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.41468 (13)	0.11825 (10)	0.58340 (4)	0.0151 (2)
N2	0.08985 (14)	0.00680 (11)	0.59213 (4)	0.0184 (2)
C2	0.21067 (16)	0.08189 (12)	0.55458 (5)	0.0164 (2)
H2	0.1602	0.1086	0.5110	0.020*
C4	0.22460 (16)	-0.00841 (12)	0.65071 (5)	0.0164 (2)

C5	0.18809 (18)	-0.08552 (13)	0.70781 (5)	0.0206 (2)
H5	0.0516	-0.1329	0.7122	0.025*
C6	0.35775 (19)	-0.09021 (14)	0.75762 (5)	0.0234 (3)
H6	0.3374	-0.1434	0.7966	0.028*
C7	0.55957 (19)	-0.01845 (14)	0.75220 (5)	0.0230 (3)
H7	0.6720	-0.0241	0.7876	0.028*
C8	0.59841 (17)	0.06012 (13)	0.69652 (5)	0.0190 (2)
H8	0.7338	0.1103	0.6929	0.023*
C9	0.42797 (16)	0.06166 (12)	0.64598 (5)	0.0153 (2)
C10	0.58686 (16)	0.19667 (13)	0.55403 (5)	0.0163 (2)
H10A	0.5488	0.1927	0.5067	0.020*
H10B	0.7223	0.1334	0.5653	0.020*
C11	0.62877 (16)	0.37280 (13)	0.57428 (5)	0.0148 (2)
C12	0.47606 (16)	0.46721 (13)	0.59998 (5)	0.0175 (2)
H12	0.3414	0.4201	0.6066	0.021*
C13	0.51869 (18)	0.63021 (13)	0.61606 (5)	0.0198 (2)
H13	0.4138	0.6936	0.6339	0.024*
C14	0.71437 (17)	0.70020 (13)	0.60598 (5)	0.0187 (2)
H14	0.7429	0.8119	0.6165	0.022*
C15	0.86811 (17)	0.60677 (13)	0.58054 (5)	0.0185 (2)
H15	1.0023	0.6544	0.5738	0.022*
C16	0.82598 (16)	0.44380 (13)	0.56497 (5)	0.0171 (2)
H16	0.9322	0.3801	0.5478	0.021*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0142 (4)	0.0141 (4)	0.0168 (4)	-0.0009 (3)	0.0017 (3)	-0.0012 (3)
N2	0.0160 (4)	0.0162 (4)	0.0226 (5)	-0.0005 (3)	0.0016 (3)	0.0004 (4)
C2	0.0155 (5)	0.0136 (5)	0.0195 (5)	0.0007 (4)	-0.0002 (4)	-0.0013 (4)
C4	0.0162 (5)	0.0131 (5)	0.0201 (5)	0.0014 (4)	0.0030 (4)	-0.0028 (4)
C5	0.0211 (5)	0.0181 (5)	0.0239 (5)	-0.0002 (4)	0.0084 (4)	-0.0008 (4)
C6	0.0305 (6)	0.0229 (6)	0.0178 (5)	0.0026 (5)	0.0074 (4)	0.0005 (4)
C7	0.0256 (6)	0.0248 (6)	0.0176 (5)	0.0029 (5)	-0.0006 (4)	-0.0033 (4)
C8	0.0171 (5)	0.0193 (5)	0.0205 (5)	-0.0003 (4)	0.0015 (4)	-0.0040 (4)
C9	0.0174 (5)	0.0119 (5)	0.0172 (5)	0.0017 (4)	0.0037 (4)	-0.0028 (4)
C10	0.0140 (5)	0.0167 (5)	0.0188 (5)	-0.0008 (4)	0.0043 (4)	-0.0016 (4)
C11	0.0154 (5)	0.0159 (5)	0.0129 (5)	0.0001 (4)	0.0009 (4)	0.0012 (4)
C12	0.0143 (5)	0.0187 (5)	0.0201 (5)	-0.0008 (4)	0.0039 (4)	0.0000 (4)
C13	0.0198 (5)	0.0173 (5)	0.0227 (5)	0.0031 (4)	0.0048 (4)	-0.0009 (4)
C14	0.0223 (5)	0.0143 (5)	0.0189 (5)	-0.0009 (4)	0.0010 (4)	0.0005 (4)
C15	0.0165 (5)	0.0198 (5)	0.0192 (5)	-0.0036 (4)	0.0027 (4)	0.0019 (4)
C16	0.0154 (5)	0.0196 (5)	0.0170 (5)	0.0009 (4)	0.0044 (4)	-0.0001 (4)

Geometric parameters (\AA , $^\circ$)

N1—C2	1.3630 (13)	C8—H8	0.95
N1—C9	1.3835 (13)	C10—C11	1.5137 (15)

N1—C10	1.4560 (13)	C10—H10A	0.99
N2—C2	1.3132 (14)	C10—H10B	0.99
N2—C4	1.3956 (13)	C11—C12	1.3892 (14)
C2—H2	0.95	C11—C16	1.3958 (14)
C4—C5	1.3992 (15)	C12—C13	1.3910 (15)
C4—C9	1.4055 (14)	C12—H12	0.95
C5—C6	1.3812 (16)	C13—C14	1.3881 (15)
C5—H5	0.95	C13—H13	0.95
C6—C7	1.4053 (17)	C14—C15	1.3870 (15)
C6—H6	0.95	C14—H14	0.95
C7—C8	1.3826 (16)	C15—C16	1.3881 (16)
C7—H7	0.95	C15—H15	0.95
C8—C9	1.3940 (14)	C16—H16	0.95
C2—N1—C9	106.23 (9)	N1—C10—C11	114.12 (8)
C2—N1—C10	127.14 (9)	N1—C10—H10A	108.7
C9—N1—C10	126.61 (8)	C11—C10—H10A	108.7
C2—N2—C4	104.17 (9)	N1—C10—H10B	108.7
N2—C2—N1	114.29 (9)	C11—C10—H10B	108.7
N2—C2—H2	122.9	H10A—C10—H10B	107.6
N1—C2—H2	122.9	C12—C11—C16	119.02 (10)
N2—C4—C5	130.19 (10)	C12—C11—C10	122.43 (9)
N2—C4—C9	109.93 (9)	C16—C11—C10	118.52 (9)
C5—C4—C9	119.80 (10)	C11—C12—C13	120.53 (10)
C6—C5—C4	117.59 (10)	C11—C12—H12	119.7
C6—C5—H5	121.2	C13—C12—H12	119.7
C4—C5—H5	121.2	C14—C13—C12	120.02 (10)
C5—C6—C7	121.82 (10)	C14—C13—H13	120.0
C5—C6—H6	119.1	C12—C13—H13	120.0
C7—C6—H6	119.1	C15—C14—C13	119.86 (10)
C8—C7—C6	121.57 (10)	C15—C14—H14	120.1
C8—C7—H7	119.2	C13—C14—H14	120.1
C6—C7—H7	119.2	C14—C15—C16	120.05 (10)
C7—C8—C9	116.34 (10)	C14—C15—H15	120.0
C7—C8—H8	121.8	C16—C15—H15	120.0
C9—C8—H8	121.8	C15—C16—C11	120.51 (10)
N1—C9—C8	131.71 (10)	C15—C16—H16	119.7
N1—C9—C4	105.38 (9)	C11—C16—H16	119.7
C8—C9—C4	122.85 (10)	 	
C4—N2—C2—N1	-0.14 (12)	N2—C4—C9—N1	-0.51 (11)
C9—N1—C2—N2	-0.17 (12)	C5—C4—C9—N1	176.66 (9)
C10—N1—C2—N2	178.24 (9)	N2—C4—C9—C8	-177.95 (9)
C2—N2—C4—C5	-176.38 (11)	C5—C4—C9—C8	-0.78 (15)
C2—N2—C4—C9	0.40 (11)	C2—N1—C10—C11	105.93 (11)
N2—C4—C5—C6	176.02 (10)	C9—N1—C10—C11	-75.97 (12)
C9—C4—C5—C6	-0.49 (15)	N1—C10—C11—C12	-20.55 (13)
C4—C5—C6—C7	0.97 (16)	N1—C10—C11—C16	161.34 (9)

C5—C6—C7—C8	−0.20 (17)	C16—C11—C12—C13	0.16 (15)
C6—C7—C8—C9	−1.03 (16)	C10—C11—C12—C13	−177.95 (10)
C2—N1—C9—C8	177.52 (11)	C11—C12—C13—C14	0.52 (16)
C10—N1—C9—C8	−0.91 (17)	C12—C13—C14—C15	−0.72 (16)
C2—N1—C9—C4	0.41 (11)	C13—C14—C15—C16	0.24 (16)
C10—N1—C9—C4	−178.02 (9)	C14—C15—C16—C11	0.45 (15)
C7—C8—C9—N1	−175.17 (10)	C12—C11—C16—C15	−0.64 (15)
C7—C8—C9—C4	1.52 (15)	C10—C11—C16—C15	177.54 (9)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C10—H10B···N2 ⁱ	0.99	2.50	3.4890 (14)	173
C7—H7···Cg1 ⁱⁱ	0.95	2.66	3.5220 (1)	151
C13—H13···Cg2 ⁱⁱⁱ	0.95	2.80	3.5660 (3)	139

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+1, -y-1, -z+1$; (iii) $x, y+1, z$.