

[μ -4-Benzoyl-1-(1-oxido-2-naphthyl-carbonyl)thiosemicarbazidato(4-)]-bis[pyridinecopper(II)]

Wen Zhang, Jin-Ping Gao, Xue-Feng Shi and Da-Cheng Li*

College of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China

Correspondence e-mail: lidacheng@lcu.edu.cn


Received 12 September 2009; accepted 7 October 2009

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; R factor = 0.046; wR factor = 0.161; data-to-parameter ratio = 12.5.

In the title dinuclear complex, $[\text{Cu}_2(\text{C}_{19}\text{H}_{11}\text{N}_3\text{O}_3\text{S})(\text{C}_5\text{H}_5\text{N})_2]$, the two Cu^{II} centers have different coordination environments, *viz.* N_2OS and N_2O_2 , each exhibiting a distorted square-planar geometry. π - π interactions between the aromatic rings of neighbouring complexes [centroid-centroid distance = 3.856 (5) Å] link pairs of molecules into centrosymmetric dimers, which are further packed into stacks along the b axis with relatively short $\text{Cu} \cdots \text{Cu}$ separations of 3.482 (1) Å. Weak intermolecular $\text{C}-\text{H} \cdots \text{N}$ hydrogen bonds help to consolidate the crystal packing.

Related literature

For details of the synthesis, see: Chen *et al.* (2007). For pharmacological properties of complexes of acylthiosemicarbazides, see: Wei *et al.* (1995).



Experimental

Crystal data

 $[\text{Cu}_2(\text{C}_{19}\text{H}_{11}\text{N}_3\text{O}_3\text{S})(\text{C}_5\text{H}_5\text{N})_2]$
 $M_r = 646.65$

 Monoclinic, $P2_1/n$
 $a = 12.7621$ (14) Å

 $b = 9.2609$ (11) Å

 $c = 21.683$ (2) Å

 $\beta = 92.168$ (2)°

 $V = 2560.9$ (5) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 1.79$ mm⁻¹
 $T = 298$ K

 $0.48 \times 0.42 \times 0.21$ mm

Data collection

Bruker SMART 1000

diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\text{min}} = 0.481$, $T_{\text{max}} = 0.706$

12450 measured reflections

4504 independent reflections

 2532 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.064$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.161$
 $S = 1.00$

4504 reflections

361 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.48$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.42$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{C23}-\text{H23} \cdots \text{N3}^i$	0.93	2.53	3.451 (8)	173

 Symmetry code: (i) $-x + 1, -y + 1, -z + 2$.

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We acknowledge the National Natural Science Foundation of China for financial support (grant No. 20671048).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2616).

References

- Chen, Y.-T., Dou, J.-M., Li, D.-C., Wang, D.-Q. & Zhu, Y.-H. (2007). *Acta Cryst.* **E63**, m2503–m2504.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). SMART and SAINT. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Wei, T. B., Chen, J. C., Yang, S. Y. & Wang, X. C. (1995). *Chem. Res. Appl.* **7**, 194–198.

supplementary materials

Acta Cryst. (2009). E65, m1363 [doi:10.1107/S1600536809040951]

[μ -4-Benzoyl-1-(1-oxido-2-naphthylcarbonyl)thiosemicarbazidato(4-)]bis[pyridinecopper(II)]

W. Zhang, J.-P. Gao, X.-F. Shi and D.-C. Li

Comment

Acylthiosemicarbazides possess strong coordination ability, and their complexes exhibit various biological activities (Wei *et al.*, 1995). Herewith we present the crystal structure of the title compound (I) - new dinuclear complex of naphthalene-carbothiosemicarbazide.

In (I) (Fig. 1), two Cu centers have different coordination environments - N₂OS and N₂O₂, respectively, exhibiting distorted square-planar geometry each. The Cu1 atom is coordinated with one phenolate oxygen, one nitrogen atom of pyridine, one sulfur atom and one hydrazide nitrogen atom, forming five-membered and six-membered chelating rings. The Cu2 atom is coordinated with two carbonyl oxygen atoms, one nitrogen atom of pyridine and one hydrazide nitrogen atom, forming five-membered and six-membered chelating rings too. π - π interactions between the aromatic rings of neighbouring complexes [centroid-to centroid distance = 3.856 (5) Å] link two molecules into centrosymmetric dimers, which are further packed into stacks along *b* axis with relatively short Cu...Cu separation of 3.482 (1) Å. Weak intermolecular C—H...N hydrogen bonds (Table 1) help to consolidate the crystal packing.

Experimental

1.406 g (10 mmol) of benzoyl chloride was added dropwise to a stirred solution of acetonitrile (50 ml) containing 0.981 g (10 mmol) of potassium thiocyanate. Stirring was continued for 1 h, and the solution was slowly warmed to ambient temperature. Then 1.68 g (10 mmol) of 1-hydroxy-2-naphthalenecarbohydrazide was added to the mixture, with stirring being continued for 5 h. After staying for overnight at refrigerator, the resulting yellow precipitate was filtered and rinsed with diethyl ether, then dried *in vacuo*, 87% yield. m.p. 225–227 C. The solution of CuNO₃(0.04 g, 0.2 mmol) in pyridine(10 ml) was added to the mixture of *N*-benzoyl-1-hydroxy-2-naphthalenecarbothiosemicarbazide (0.073 g, 0.2 mmol)and sodium methylate(0.0324 g,0.6 mmol)in DMSO(10 ml). A green solution was obtained after refluxing for 3 h. After filtrated,dimethyl ether was slowly diffused into the filtrate, then crystals suitable for X-ray diffraction were obtained after two weeks (m.p. >400 K) (Chen *et al.* 2007) Elemental analysis calculated for Cu₂C₂₉H₂₁N₅O₃S₁: C, 53.64; H, 3.29; N, 9.43. Found (%): C, 53.86; H, 3.27; N, 10.83.

Refinement

The C-bound H atoms were geometrically positioned (C—H = 0.93 Å), and were refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids. H atoms have been omitted for clarity.

[μ -4-Benzoyl-1-(1-oxido-2-naphthylcarbonyl)thiosemicarbazidato(4-)]bis[pyridinecopper(II)]

Crystal data

[Cu₂(C₁₉H₁₁N₃O₃S)(C₅H₅N₁)₂]

$M_r = 646.65$

Monoclinic, $P2_1/n$

$a = 12.7621$ (14) Å

$b = 9.2609$ (11) Å

$c = 21.683$ (2) Å

$\beta = 92.168$ (2)°

$V = 2560.9$ (5) Å³

$Z = 4$

$F_{000} = 1312$

$D_x = 1.677$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2465 reflections

$\theta = 2.4$ – 22.0 °

$\mu = 1.79$ mm⁻¹

$T = 298$ K

Block, green

$0.48 \times 0.42 \times 0.21$ mm

Data collection

Bruker SMART 1000
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298$ K

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.481$, $T_{\max} = 0.706$

12450 measured reflections

4504 independent reflections

2532 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.064$

$\theta_{\text{max}} = 25.0$ °

$\theta_{\text{min}} = 1.8$ °

$h = -15$ → 15

$k = -9$ → 11

$l = -22$ → 25

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.046$

$wR(F^2) = 0.161$

$S = 1.00$

4504 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0835P)^2]$,

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.001$

$\Delta\rho_{\text{max}} = 0.48$ e Å⁻³

361 parameters

$$\Delta\rho_{\min} = -0.42 \text{ e } \text{\AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.28704 (5)	0.20605 (8)	1.04261 (3)	0.0436 (3)
Cu2	-0.01899 (6)	0.36338 (9)	0.94557 (3)	0.0496 (3)
S1	0.33285 (12)	0.34418 (19)	0.96271 (7)	0.0508 (5)
N1	0.1252 (4)	0.3405 (5)	0.9676 (2)	0.0431 (12)
N2	0.1458 (4)	0.2438 (5)	1.0159 (2)	0.0416 (12)
N3	0.1971 (4)	0.4823 (5)	0.8891 (2)	0.0418 (12)
N4	0.4361 (4)	0.1750 (5)	1.0747 (2)	0.0432 (12)
N5	-0.1717 (4)	0.3765 (6)	0.9268 (2)	0.0469 (13)
O1	0.0142 (3)	0.4972 (5)	0.88313 (19)	0.0540 (12)
O2	-0.0331 (3)	0.2309 (5)	1.0121 (2)	0.0585 (12)
O3	0.2433 (3)	0.0804 (5)	1.10597 (19)	0.0526 (11)
C1	0.2062 (5)	0.3912 (6)	0.9385 (3)	0.0415 (15)
C2	0.1062 (5)	0.5270 (7)	0.8657 (3)	0.0446 (15)
C3	0.1091 (5)	0.6286 (6)	0.8125 (3)	0.0407 (14)
C4	0.0197 (5)	0.6999 (8)	0.7931 (3)	0.0595 (19)
H4	-0.0429	0.6815	0.8122	0.071*
C5	0.0213 (6)	0.7982 (8)	0.7456 (3)	0.067 (2)
H5	-0.0396	0.8474	0.7337	0.080*
C6	0.1120 (5)	0.8241 (7)	0.7158 (3)	0.0542 (18)
H6	0.1135	0.8910	0.6839	0.065*
C7	0.2003 (5)	0.7498 (8)	0.7339 (3)	0.0561 (18)
H7	0.2620	0.7652	0.7134	0.067*
C8	0.1998 (5)	0.6532 (7)	0.7815 (3)	0.0527 (17)
H8	0.2609	0.6038	0.7931	0.063*
C9	0.0565 (5)	0.1886 (7)	1.0356 (3)	0.0488 (16)
C10	0.1489 (5)	0.0355 (7)	1.1163 (3)	0.0459 (16)
C11	0.0576 (5)	0.0804 (7)	1.0840 (3)	0.0466 (16)
C12	-0.0399 (5)	0.0174 (8)	1.0983 (3)	0.066 (2)
H12	-0.0999	0.0467	1.0760	0.079*
C13	-0.0496 (6)	-0.0831 (9)	1.1429 (3)	0.070 (2)
H13	-0.1151	-0.1215	1.1507	0.084*
C14	0.0390 (5)	-0.1290 (7)	1.1771 (3)	0.0552 (18)
C15	0.1381 (5)	-0.0719 (7)	1.1649 (3)	0.0460 (16)
C16	0.2265 (5)	-0.1219 (7)	1.1990 (3)	0.0511 (17)
H16	0.2925	-0.0856	1.1908	0.061*
C17	0.2165 (6)	-0.2236 (8)	1.2443 (3)	0.0607 (19)
H17	0.2755	-0.2570	1.2665	0.073*
C18	0.1178 (6)	-0.2766 (8)	1.2570 (3)	0.064 (2)
H18	0.1110	-0.3431	1.2888	0.077*
C19	0.0320 (6)	-0.2337 (8)	1.2244 (3)	0.0615 (19)
H19	-0.0328	-0.2733	1.2329	0.074*

supplementary materials

C20	0.4648 (6)	0.0479 (8)	1.0975 (3)	0.0620 (19)
H20	0.4152	-0.0257	1.0978	0.074*
C21	0.5641 (6)	0.0195 (9)	1.1207 (4)	0.075 (2)
H21	0.5816	-0.0723	1.1351	0.090*
C22	0.6363 (6)	0.1260 (9)	1.1223 (3)	0.071 (2)
H22	0.7040	0.1089	1.1380	0.085*
C23	0.6084 (5)	0.2589 (8)	1.1005 (3)	0.0593 (19)
H23	0.6562	0.3347	1.1019	0.071*
C24	0.5086 (5)	0.2789 (7)	1.0765 (3)	0.0497 (16)
H24	0.4905	0.3693	1.0606	0.060*
C25	-0.2383 (5)	0.2922 (8)	0.9581 (3)	0.0553 (18)
H25	-0.2114	0.2330	0.9896	0.066*
C26	-0.3436 (5)	0.2916 (8)	0.9450 (3)	0.060 (2)
H26	-0.3874	0.2313	0.9667	0.072*
C27	-0.3842 (5)	0.3799 (9)	0.8998 (3)	0.063 (2)
H27	-0.4559	0.3805	0.8904	0.075*
C28	-0.3189 (5)	0.4669 (9)	0.8687 (3)	0.065 (2)
H28	-0.3447	0.5290	0.8380	0.078*
C29	-0.2128 (5)	0.4602 (8)	0.8839 (3)	0.0571 (18)
H29	-0.1679	0.5190	0.8623	0.069*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0386 (4)	0.0450 (5)	0.0473 (5)	-0.0080 (4)	0.0043 (3)	0.0035 (4)
Cu2	0.0350 (4)	0.0602 (6)	0.0536 (5)	-0.0116 (4)	0.0012 (4)	0.0081 (4)
S1	0.0370 (9)	0.0626 (12)	0.0528 (10)	-0.0031 (8)	0.0045 (7)	0.0147 (8)
N1	0.038 (3)	0.044 (3)	0.047 (3)	-0.012 (2)	-0.001 (2)	0.008 (2)
N2	0.039 (3)	0.051 (3)	0.035 (3)	-0.010 (2)	0.000 (2)	0.011 (2)
N3	0.036 (3)	0.046 (3)	0.043 (3)	-0.005 (2)	0.000 (2)	0.009 (2)
N4	0.043 (3)	0.036 (3)	0.050 (3)	-0.003 (2)	0.002 (2)	0.005 (2)
N5	0.038 (3)	0.056 (4)	0.046 (3)	-0.006 (3)	-0.004 (3)	-0.005 (3)
O1	0.033 (2)	0.065 (3)	0.063 (3)	-0.012 (2)	-0.001 (2)	0.015 (2)
O2	0.042 (3)	0.075 (3)	0.059 (3)	-0.012 (2)	0.001 (2)	0.017 (2)
O3	0.038 (2)	0.061 (3)	0.059 (3)	-0.009 (2)	0.003 (2)	0.014 (2)
C1	0.043 (4)	0.038 (4)	0.043 (4)	-0.004 (3)	0.000 (3)	-0.005 (3)
C2	0.039 (4)	0.048 (4)	0.047 (4)	-0.009 (3)	0.001 (3)	-0.004 (3)
C3	0.038 (3)	0.042 (4)	0.041 (3)	-0.006 (3)	-0.005 (3)	0.004 (3)
C4	0.041 (4)	0.080 (5)	0.058 (4)	-0.003 (4)	0.006 (3)	0.011 (4)
C5	0.050 (4)	0.082 (6)	0.069 (5)	0.011 (4)	-0.003 (4)	0.019 (4)
C6	0.060 (5)	0.055 (5)	0.048 (4)	-0.010 (4)	-0.002 (3)	0.011 (3)
C7	0.044 (4)	0.068 (5)	0.056 (4)	-0.001 (4)	0.007 (3)	0.009 (4)
C8	0.046 (4)	0.059 (5)	0.053 (4)	0.001 (3)	-0.001 (3)	0.006 (3)
C9	0.042 (4)	0.060 (5)	0.045 (4)	-0.012 (3)	0.006 (3)	-0.003 (3)
C10	0.044 (4)	0.052 (4)	0.042 (4)	-0.013 (3)	0.003 (3)	-0.005 (3)
C11	0.044 (4)	0.053 (4)	0.043 (4)	-0.015 (3)	0.009 (3)	0.005 (3)
C12	0.044 (4)	0.087 (6)	0.067 (5)	-0.021 (4)	0.003 (3)	0.024 (4)
C13	0.052 (5)	0.091 (6)	0.069 (5)	-0.032 (4)	0.015 (4)	0.020 (4)

C14	0.054 (4)	0.053 (5)	0.059 (4)	-0.016 (4)	0.008 (4)	0.010 (3)
C15	0.060 (4)	0.042 (4)	0.037 (4)	-0.007 (3)	0.013 (3)	-0.003 (3)
C16	0.048 (4)	0.056 (5)	0.050 (4)	-0.005 (3)	0.008 (3)	0.005 (3)
C17	0.070 (5)	0.058 (5)	0.054 (4)	0.012 (4)	0.006 (4)	0.002 (4)
C18	0.079 (6)	0.057 (5)	0.057 (5)	0.000 (4)	0.026 (4)	0.012 (4)
C19	0.060 (5)	0.061 (5)	0.065 (5)	-0.016 (4)	0.011 (4)	0.008 (4)
C20	0.061 (5)	0.049 (5)	0.075 (5)	-0.008 (4)	-0.001 (4)	0.008 (4)
C21	0.067 (5)	0.050 (5)	0.107 (6)	0.010 (4)	-0.005 (5)	0.021 (4)
C22	0.047 (4)	0.079 (6)	0.087 (6)	0.007 (4)	-0.005 (4)	0.006 (5)
C23	0.048 (4)	0.061 (5)	0.068 (5)	-0.016 (4)	-0.005 (4)	0.006 (4)
C24	0.045 (4)	0.046 (4)	0.059 (4)	-0.003 (3)	0.011 (3)	0.011 (3)
C25	0.045 (4)	0.072 (5)	0.050 (4)	-0.013 (4)	0.011 (3)	-0.003 (4)
C26	0.048 (4)	0.081 (6)	0.053 (4)	-0.022 (4)	0.016 (3)	-0.011 (4)
C27	0.039 (4)	0.088 (6)	0.062 (5)	-0.014 (4)	0.006 (4)	-0.022 (4)
C28	0.036 (4)	0.093 (6)	0.067 (5)	-0.010 (4)	-0.005 (4)	-0.009 (4)
C29	0.039 (4)	0.073 (5)	0.060 (4)	-0.010 (4)	0.005 (3)	-0.002 (4)

Geometric parameters (Å, °)

Cu1—O3	1.900 (4)	C10—C15	1.460 (8)
Cu1—N2	1.905 (5)	C11—C12	1.420 (8)
Cu1—N4	2.022 (5)	C12—C13	1.350 (9)
Cu1—S1	2.2487 (17)	C12—H12	0.9300
Cu2—O1	1.895 (4)	C13—C14	1.395 (9)
Cu2—N1	1.896 (5)	C13—H13	0.9300
Cu2—O2	1.908 (4)	C14—C15	1.405 (8)
Cu2—N5	1.980 (5)	C14—C19	1.415 (9)
S1—C1	1.736 (6)	C15—C16	1.404 (8)
N1—C1	1.317 (7)	C16—C17	1.370 (9)
N1—N2	1.395 (6)	C16—H16	0.9300
N2—C9	1.333 (7)	C17—C18	1.390 (9)
N3—C2	1.314 (7)	C17—H17	0.9300
N3—C1	1.366 (7)	C18—C19	1.341 (9)
N4—C20	1.322 (8)	C18—H18	0.9300
N4—C24	1.335 (7)	C19—H19	0.9300
N5—C29	1.305 (8)	C20—C21	1.372 (9)
N5—C25	1.355 (8)	C20—H20	0.9300
O1—C2	1.278 (7)	C21—C22	1.349 (10)
O2—C9	1.295 (7)	C21—H21	0.9300
O3—C10	1.302 (7)	C22—C23	1.361 (10)
C2—C3	1.490 (8)	C22—H22	0.9300
C3—C4	1.372 (8)	C23—C24	1.371 (8)
C3—C8	1.379 (8)	C23—H23	0.9300
C4—C5	1.375 (9)	C24—H24	0.9300
C4—H4	0.9300	C25—C26	1.363 (9)
C5—C6	1.368 (9)	C25—H25	0.9300
C5—H5	0.9300	C26—C27	1.364 (10)
C6—C7	1.365 (8)	C26—H26	0.9300
C6—H6	0.9300	C27—C28	1.356 (9)

supplementary materials

C7—C8	1.367 (9)	C27—H27	0.9300
C7—H7	0.9300	C28—C29	1.384 (8)
C8—H8	0.9300	C28—H28	0.9300
C9—C11	1.451 (8)	C29—H29	0.9300
C10—C11	1.400 (8)		
O3—Cu1—N2	91.84 (18)	C11—C10—C15	117.6 (5)
O3—Cu1—N4	87.81 (18)	C10—C11—C12	119.3 (6)
N2—Cu1—N4	176.7 (2)	C10—C11—C9	123.5 (5)
O3—Cu1—S1	175.88 (14)	C12—C11—C9	117.2 (6)
N2—Cu1—S1	86.09 (14)	C13—C12—C11	122.9 (6)
N4—Cu1—S1	94.46 (14)	C13—C12—H12	118.5
O1—Cu2—N1	90.70 (19)	C11—C12—H12	118.5
O1—Cu2—O2	172.29 (18)	C12—C13—C14	119.8 (6)
N1—Cu2—O2	81.73 (19)	C12—C13—H13	120.1
O1—Cu2—N5	93.3 (2)	C14—C13—H13	120.1
N1—Cu2—N5	176.0 (2)	C13—C14—C15	120.1 (6)
O2—Cu2—N5	94.3 (2)	C13—C14—C19	121.5 (6)
C1—S1—Cu1	96.3 (2)	C15—C14—C19	118.4 (6)
C1—N1—N2	117.4 (5)	C16—C15—C14	119.2 (6)
C1—N1—Cu2	127.7 (4)	C16—C15—C10	120.6 (6)
N2—N1—Cu2	114.4 (3)	C14—C15—C10	120.2 (6)
C9—N2—N1	110.4 (5)	C17—C16—C15	120.6 (6)
C9—N2—Cu1	129.9 (4)	C17—C16—H16	119.7
N1—N2—Cu1	119.7 (3)	C15—C16—H16	119.7
C2—N3—C1	123.0 (5)	C16—C17—C18	119.6 (7)
C20—N4—C24	116.7 (6)	C16—C17—H17	120.2
C20—N4—Cu1	119.9 (4)	C18—C17—H17	120.2
C24—N4—Cu1	123.3 (4)	C19—C18—C17	121.2 (7)
C29—N5—C25	117.2 (6)	C19—C18—H18	119.4
C29—N5—Cu2	123.2 (4)	C17—C18—H18	119.4
C25—N5—Cu2	119.5 (5)	C18—C19—C14	120.9 (7)
C2—O1—Cu2	125.8 (4)	C18—C19—H19	119.6
C9—O2—Cu2	112.6 (4)	C14—C19—H19	119.6
C10—O3—Cu1	128.3 (4)	N4—C20—C21	123.1 (7)
N1—C1—N3	123.4 (5)	N4—C20—H20	118.4
N1—C1—S1	120.3 (5)	C21—C20—H20	118.4
N3—C1—S1	116.3 (4)	C22—C21—C20	119.4 (7)
O1—C2—N3	128.8 (6)	C22—C21—H21	120.3
O1—C2—C3	114.5 (5)	C20—C21—H21	120.3
N3—C2—C3	116.7 (5)	C21—C22—C23	118.8 (7)
C4—C3—C8	118.3 (6)	C21—C22—H22	120.6
C4—C3—C2	119.8 (6)	C23—C22—H22	120.6
C8—C3—C2	121.8 (6)	C22—C23—C24	118.8 (7)
C3—C4—C5	120.9 (6)	C22—C23—H23	120.6
C3—C4—H4	119.6	C24—C23—H23	120.6
C5—C4—H4	119.6	N4—C24—C23	123.1 (6)
C6—C5—C4	120.5 (7)	N4—C24—H24	118.4
C6—C5—H5	119.8	C23—C24—H24	118.4
C4—C5—H5	119.8	N5—C25—C26	122.0 (7)

C7—C6—C5	118.7 (6)	N5—C25—H25	119.0
C7—C6—H6	120.7	C26—C25—H25	119.0
C5—C6—H6	120.7	C25—C26—C27	119.5 (7)
C6—C7—C8	121.4 (6)	C25—C26—H26	120.2
C6—C7—H7	119.3	C27—C26—H26	120.2
C8—C7—H7	119.3	C28—C27—C26	119.3 (7)
C7—C8—C3	120.3 (6)	C28—C27—H27	120.3
C7—C8—H8	119.9	C26—C27—H27	120.3
C3—C8—H8	119.9	C27—C28—C29	117.9 (7)
O2—C9—N2	120.7 (6)	C27—C28—H28	121.0
O2—C9—C11	118.5 (5)	C29—C28—H28	121.0
N2—C9—C11	120.8 (6)	N5—C29—C28	124.0 (7)
O3—C10—C11	125.2 (6)	N5—C29—H29	118.0
O3—C10—C15	117.1 (6)	C28—C29—H29	118.0
O3—Cu1—S1—C1	63.1 (19)	C3—C4—C5—C6	1.7 (11)
N2—Cu1—S1—C1	3.2 (2)	C4—C5—C6—C7	0.4 (11)
N4—Cu1—S1—C1	-173.5 (2)	C5—C6—C7—C8	-1.3 (11)
O1—Cu2—N1—C1	-8.4 (5)	C6—C7—C8—C3	0.0 (11)
O2—Cu2—N1—C1	173.0 (6)	C4—C3—C8—C7	2.0 (10)
N5—Cu2—N1—C1	167 (3)	C2—C3—C8—C7	-178.4 (6)
O1—Cu2—N1—N2	-179.8 (4)	Cu2—O2—C9—N2	4.1 (8)
O2—Cu2—N1—N2	1.6 (4)	Cu2—O2—C9—C11	-176.1 (5)
N5—Cu2—N1—N2	-5(3)	N1—N2—C9—O2	-2.8 (8)
C1—N1—N2—C9	-172.4 (5)	Cu1—N2—C9—O2	178.5 (4)
Cu2—N1—N2—C9	0.0 (6)	N1—N2—C9—C11	177.5 (5)
C1—N1—N2—Cu1	6.5 (7)	Cu1—N2—C9—C11	-1.2 (9)
Cu2—N1—N2—Cu1	178.9 (2)	Cu1—O3—C10—C11	-4.5 (9)
O3—Cu1—N2—C9	-3.5 (6)	Cu1—O3—C10—C15	174.4 (4)
N4—Cu1—N2—C9	-87 (3)	O3—C10—C11—C12	177.1 (6)
S1—Cu1—N2—C9	172.9 (6)	C15—C10—C11—C12	-1.9 (9)
O3—Cu1—N2—N1	177.9 (4)	O3—C10—C11—C9	-2.4 (11)
N4—Cu1—N2—N1	94 (3)	C15—C10—C11—C9	178.6 (6)
S1—Cu1—N2—N1	-5.7 (4)	O2—C9—C11—C10	-174.5 (6)
O3—Cu1—N4—C20	34.8 (5)	N2—C9—C11—C10	5.3 (10)
N2—Cu1—N4—C20	119 (3)	O2—C9—C11—C12	6.0 (9)
S1—Cu1—N4—C20	-141.7 (5)	N2—C9—C11—C12	-174.2 (6)
O3—Cu1—N4—C24	-142.7 (5)	C10—C11—C12—C13	1.3 (12)
N2—Cu1—N4—C24	-59 (3)	C9—C11—C12—C13	-179.2 (7)
S1—Cu1—N4—C24	40.7 (5)	C11—C12—C13—C14	0.0 (12)
O1—Cu2—N5—C29	-0.9 (5)	C12—C13—C14—C15	-0.6 (12)
N1—Cu2—N5—C29	-176 (3)	C12—C13—C14—C19	179.8 (7)
O2—Cu2—N5—C29	177.8 (5)	C13—C14—C15—C16	-178.6 (7)
O1—Cu2—N5—C25	177.0 (5)	C19—C14—C15—C16	1.0 (10)
N1—Cu2—N5—C25	2(3)	C13—C14—C15—C10	-0.1 (10)
O2—Cu2—N5—C25	-4.3 (5)	C19—C14—C15—C10	179.6 (6)
N1—Cu2—O1—C2	7.9 (5)	O3—C10—C15—C16	0.9 (9)
O2—Cu2—O1—C2	18.4 (18)	C11—C10—C15—C16	179.9 (6)
N5—Cu2—O1—C2	-171.8 (5)	O3—C10—C15—C14	-177.7 (6)
O1—Cu2—O2—C9	-13.7 (18)	C11—C10—C15—C14	1.3 (9)

supplementary materials

N1—Cu2—O2—C9	-3.0 (4)	C14—C15—C16—C17	-0.9 (10)
N5—Cu2—O2—C9	176.6 (4)	C10—C15—C16—C17	-179.5 (6)
N2—Cu1—O3—C10	6.4 (5)	C15—C16—C17—C18	-0.7 (10)
N4—Cu1—O3—C10	-176.9 (5)	C16—C17—C18—C19	2.3 (11)
S1—Cu1—O3—C10	-53 (2)	C17—C18—C19—C14	-2.2 (11)
N2—N1—C1—N3	177.5 (5)	C13—C14—C19—C18	-179.8 (7)
Cu2—N1—C1—N3	6.3 (9)	C15—C14—C19—C18	0.5 (11)
N2—N1—C1—S1	-3.0 (8)	C24—N4—C20—C21	-1.6 (10)
Cu2—N1—C1—S1	-174.2 (3)	Cu1—N4—C20—C21	-179.3 (6)
C2—N3—C1—N1	-0.4 (9)	N4—C20—C21—C22	1.9 (12)
C2—N3—C1—S1	180.0 (5)	C20—C21—C22—C23	-0.3 (12)
Cu1—S1—C1—N1	-1.1 (5)	C21—C22—C23—C24	-1.4 (11)
Cu1—S1—C1—N3	178.5 (4)	C20—N4—C24—C23	-0.2 (9)
Cu2—O1—C2—N3	-5.5 (10)	Cu1—N4—C24—C23	177.4 (5)
Cu2—O1—C2—C3	175.8 (4)	C22—C23—C24—N4	1.7 (10)
C1—N3—C2—O1	0.1 (10)	C29—N5—C25—C26	1.4 (10)
C1—N3—C2—C3	178.8 (5)	Cu2—N5—C25—C26	-176.7 (5)
O1—C2—C3—C4	10.6 (9)	N5—C25—C26—C27	-1.2 (11)
N3—C2—C3—C4	-168.2 (6)	C25—C26—C27—C28	0.0 (11)
O1—C2—C3—C8	-168.9 (6)	C26—C27—C28—C29	0.9 (11)
N3—C2—C3—C8	12.2 (9)	C25—N5—C29—C28	-0.4 (10)
C8—C3—C4—C5	-2.9 (10)	Cu2—N5—C29—C28	177.6 (5)
C2—C3—C4—C5	177.5 (6)	C27—C28—C29—N5	-0.7 (11)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C23—H23 \cdots N3 ⁱ	0.93	2.53	3.451 (8)	173

Symmetry codes: (i) $-x+1, -y+1, -z+2$.

Fig. 1

