

trans-Dichloridotetrakis[1-(2-hydroxyethyl)-1*H*-tetrazole- κ N⁴]cobalt(II)

Alexander S. Lyakhov, Anastasiya P. Mosalkova, Mikhail M. Degtyarik, Ludmila S. Ivashkevich* and Pavel N. Gaponik

Research Institute for Physico-Chemical Problems, Belarusian State University, Leningradskaya Str. 14, Minsk 220030, Belarus
Correspondence e-mail: iva@bsu.by


Received 12 October 2009; accepted 14 October 2009

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.038; wR factor = 0.097; data-to-parameter ratio = 16.9.

The title cobalt(II) complex, $[\text{CoCl}_2(\text{C}_3\text{H}_6\text{N}_4\text{O})_4]$, was obtained from metallic cobalt by direct synthesis. There are two Co atoms in the asymmetric unit, each lying on an inversion centre and adopting a distorted octahedral coordination. Classical and non-classical hydrogen bonds are responsible for formation of a three-dimensional polymeric network in the crystal.

Related literature

For a review of complexes of 1-substituted tetrazoles, see: Gaponik *et al.* (2006). For the crystal structure of a related Co(II) complex, see: Shvedenkov *et al.* (2003). For a description of the Cambridge Structural Database, see: Allen (2002).



Experimental

Crystal data

$[\text{CoCl}_2(\text{C}_3\text{H}_6\text{N}_4\text{O})_4]$	$\gamma = 69.481$ (17) $^\circ$
$M_r = 586.30$	$V = 1180.4$ (5) Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 6.8971$ (19) Å	Mo $K\alpha$ radiation
$b = 9.4602$ (17) Å	$\mu = 1.01$ mm ⁻¹
$c = 19.761$ (4) Å	$T = 294$ K
$\alpha = 77.870$ (14) $^\circ$	$0.24 \times 0.16 \times 0.15$ mm
$\beta = 86.721$ (19) $^\circ$	

Data collection

Nicolet R3m four-circle diffractometer	5449 independent reflections
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	3747 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.794$, $T_{\max} = 0.863$	$R_{\text{int}} = 0.024$
5904 measured reflections	2 standard reflections every 100 reflections
	intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	323 parameters
$wR(F^2) = 0.097$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.58$ e Å ⁻³
5449 reflections	$\Delta\rho_{\text{min}} = -0.32$ e Å ⁻³

Table 1

Selected bond lengths (Å).

Co1—N24	2.144 (2)	Co2—N44	2.165 (2)
Co1—N14	2.191 (2)	Co2—N34	2.168 (2)
Co1—Cl1	2.4372 (10)	Co2—Cl2	2.4333 (10)

Table 2

Hydrogen-bond geometry (Å, $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots O3 ⁱ	0.82	1.94	2.741 (3)	165
O2—H2 \cdots Cl1 ⁱⁱ	0.82	2.32	3.104 (2)	160
O3—H3 \cdots Cl2 ⁱⁱⁱ	0.82	2.28	3.098 (3)	173
O4—H4 \cdots O2 ^{iv}	0.82	1.94	2.757 (3)	175
C15—H15 \cdots Cl1 ⁱⁱ	0.93	2.76	3.482 (3)	135
C25—H25 \cdots O1 ^v	0.93	2.56	3.303 (4)	137
C35—H35 \cdots O4 ^{vi}	0.93	2.38	3.156 (3)	141
C45—H45 \cdots Cl2 ⁱⁱⁱ	0.93	2.54	3.405 (3)	155

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $x + 1, y, z$; (iii) $x - 1, y, z$; (iv) $-x + 1, -y, -z$; (v) $-x, -y + 1, -z + 1$; (vi) $-x, -y, -z$.

Data collection: *R3m Software* (Nicolet, 1980); cell refinement: *R3m Software*; data reduction: *OMNIBUS* (Gałdecka, 2002); program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97* and *PLATON*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2499).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Galdecka, E. (2002). *J. Appl. Cryst.* **35**, 641–643.
- Gaponik, P. N., Voitekhovich, S. V. & Ivashkevich, O. A. (2006). *Russ. Chem. Rev.* **75**, 507–539.
- Nicolet (1980). *R3m Software*. Nicolet XRD Corporation, Cupertino, USA.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shvedenkov, Y. G., Virovets, A. V. & Lavrenova, L. G. (2003). *Russ. Chem. Bull.* **52**, 1353–1357.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, m1397-m1398 [doi:10.1107/S1600536809042263]

***trans*-Dichloridotetrakis[1-(2-hydroxyethyl)-1*H*-tetrazole- κ N⁴]cobalt(II)**

A. S. Lyakhov, A. P. Mosalkova, M. M. Degtyarik, L. S. Ivashkevich and P. N. Gaponik

Comment

Coordination compounds of 1-substituted tetrazoles have been the subject of many investigations (Gaponik *et al.*, 2006). Among reported metal(II) halide complexes containing 1-alkyltetrazoles overwhelming majority present copper(II) complexes CuL_2X_2 , where $L = 1\text{-alkyltetrazole}$, $X = \text{Cl}$ or Br . Until now, only one cobalt(II) chloride complex with 1-allyltetrazole of composition CoL_2Cl_2 have been structurally characterized (Shvedenkov *et al.*, 2003). Here we present novel cobalt(II) chloride complex, namely CoL_4Cl_2 where L is 1-(2-hydroxyethyl)tetrazole, obtained by dissolving metallic cobalt in a methanol solution of 1-(2-hydroxyethyl)tetrazole in presence of hydrochloric acid in air. This is the first complex of such composition among metal(II) halide with 1-alkyltetrazoles obtained by now.

The title compound, (I), presents molecular complex, with two Co atoms in the asymmetric unit, both lying on inversion centres (Fig. 1). Co atoms adopt rather distorted octahedral coordination composed of two Cl atoms in axial positions and four tetrazole ring N⁴ atoms in equatorial sites (Table 1). So, the complex molecules present *trans*-isomers.

The tetrazole ring geometry of ligand molecules is usual for 1-substituted tetrazoles (Cambridge Structural Database, Version 5.30 of November 2008; Allen, 2002).

The crystal packing of (I) is stabilized by a series of intermolecular hydrogen bonds (Table 2). Classic HB, O—H \cdots O and O—H \cdots Cl, link complex molecules to polymeric layers parallel to the *ac* plane (Fig. 2). Non-classic HB C^{Tz}—H \cdots Cl, formed by the tetrazole ring C—H groups, are additional within the layers. The above layers are connected *via* non-classic C^{Tz}—H \cdots O hydrogen bonds to give three-dimensional polymeric network (Fig. 3).

Experimental

A mixture, containing cobalt powder (0.06 g, 0.001 mol), 1-(2-hydroxyethyl)tetrazole (0.47 g, 0.0041 mol), 5 ml of methanol and 0.2 ml of concentrated solution of HCl, was heated at 325 K with stirring on air until the metal was fully dissolved (6 h). Pink crystals of the title complex were grown by slow evaporation of the reaction mixture in air at room temperature during two weeks. The crystals were filtered off, washed with diethyl ether and dried in air (0.45 g, yield 85%; m.p. 383 K; decomp. 468 K). Calc. (%): Cu 10.1, Cl 12.1. Found (%): Cu 10.2, Cl 12.8. IR (cm^{-1}): 3391 (*s*), 3091 (*s*), 3134 (*s*), 2896 (*m*), 2946 (*m*), 2975 (*s*), 1623 (*s*), 1497 (*s*), 1437 (*s*), 1380 (*w*), 1359 (*m*), 1275 (*m*), 1247 (*w*), 1171 (*s*), 1099 (*s*), 1062 (*s*), 998 (*s*), 259 (*s*), 200 (*m*), 176(*m*).

Refinement

H atoms were placed in calculated positions and refined using riding model, with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$ for the methylene and the tetrazole ring CH group, and $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}(\text{O})$ for the hydroxyl groups.

Figures




Fig. 1. Two complex molecules in the crystal structure of (I), with atom numbering for the asymmetric unit. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as spheres of arbitrary radii.




Fig. 2. A layer in the structure of (I) parallel with the *ac* plane formed by classic hydrogen bonds O—H...O and O—H...Cl (dashed lines). Only H atoms, participating in classic hydrogen bonds, are shown.




Fig. 3. The crystal structure of (I) viewed along the *a* axis. Dashed lines show hydrogen bonds.

trans-Dichloridotetrakis[1-(2-hydroxyethyl)-1*H*-tetrazole- κN^4]cobalt(II)

Crystal data

[CoCl₂(C₃H₆N₄O)₄]

M_r = 586.30

Triclinic, *P* $\bar{1}$

Hall symbol: -P 1

a = 6.8971 (19) Å

b = 9.4602 (17) Å

c = 19.761 (4) Å

α = 77.870 (14)°

β = 86.721 (19)°

γ = 69.481 (17)°

V = 1180.4 (5) Å³

Z = 2

*F*₀₀₀ = 602

D_x = 1.650 Mg m⁻³

Melting point: 383 K

Mo *K* α radiation, λ = 0.71073 Å

Cell parameters from 25 reflections

θ = 13.4–19.5°

μ = 1.01 mm⁻¹

T = 294 K

Prism, pink

0.24 × 0.16 × 0.15 mm

Data collection

Nicolet R3m four-circle diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 294 K

$\omega/2\theta$ scans

Absorption correction: ψ scan (North *et al.*, 1968)

*T*_{min} = 0.794, *T*_{max} = 0.863

*R*_{int} = 0.024

θ_{\max} = 27.6°

θ_{\min} = 1.1°

h = 0→8

k = -11→12

l = -25→25

2 standard reflections

5904 measured reflections
 5449 independent reflections
 3747 reflections with $I > 2\sigma(I)$

every 100 reflections
 intensity decay: none

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.097$
 $S = 1.04$
 5449 reflections
 323 parameters
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.031P)^2 + 0.9939P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.58 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.32 \text{ e } \text{\AA}^{-3}$
 Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.0000	0.0000	0.5000	0.02622 (12)
Cl1	-0.33214 (10)	0.17425 (8)	0.44772 (3)	0.03638 (16)
N11	0.1924 (3)	0.3322 (2)	0.56697 (11)	0.0329 (5)
N12	-0.0116 (4)	0.3971 (3)	0.57773 (15)	0.0471 (6)
N13	-0.1006 (4)	0.3080 (3)	0.56183 (14)	0.0450 (6)
N14	0.0441 (3)	0.1851 (2)	0.54069 (11)	0.0326 (5)
C15	0.2219 (4)	0.2040 (3)	0.54426 (14)	0.0342 (6)
H15	0.3491	0.1380	0.5327	0.041*
C16	0.3466 (4)	0.3983 (3)	0.58033 (15)	0.0389 (6)
H16A	0.2933	0.5089	0.5631	0.047*
H16B	0.4718	0.3556	0.5555	0.047*
C17	0.3979 (5)	0.3655 (3)	0.65626 (16)	0.0445 (7)
H17A	0.4320	0.2561	0.6750	0.053*
H17B	0.5183	0.3923	0.6624	0.053*
O1	0.2309 (4)	0.4496 (2)	0.69294 (12)	0.0564 (6)

supplementary materials

H1	0.1933	0.3896	0.7222	0.085*
N21	0.2420 (3)	0.1961 (3)	0.31580 (11)	0.0320 (5)
N22	0.3564 (4)	0.0462 (3)	0.32073 (13)	0.0455 (6)
N23	0.2998 (4)	-0.0278 (3)	0.37622 (13)	0.0417 (6)
N24	0.1491 (3)	0.0717 (2)	0.40773 (11)	0.0312 (5)
C25	0.1170 (4)	0.2095 (3)	0.36908 (14)	0.0369 (6)
H25	0.0219	0.3013	0.3778	0.044*
C26	0.2763 (5)	0.3161 (3)	0.26103 (14)	0.0417 (7)
H26A	0.1492	0.4045	0.2521	0.050*
H26B	0.3132	0.2770	0.2187	0.050*
C27	0.4452 (5)	0.3655 (3)	0.28125 (16)	0.0458 (7)
H27A	0.4527	0.4530	0.2468	0.055*
H27B	0.4124	0.3981	0.3252	0.055*
O2	0.6405 (3)	0.2449 (3)	0.28720 (11)	0.0480 (5)
H2	0.6670	0.2039	0.3281	0.072*
Co2	0.5000	0.0000	0.0000	0.02837 (12)
Cl2	0.84297 (10)	-0.16973 (8)	0.04419 (4)	0.03764 (16)
N31	0.1868 (4)	-0.1224 (3)	0.18430 (12)	0.0368 (5)
N32	0.3021 (4)	-0.2628 (3)	0.17307 (14)	0.0507 (7)
N33	0.4073 (4)	-0.2413 (3)	0.11812 (14)	0.0472 (7)
N34	0.3617 (3)	-0.0878 (2)	0.09318 (11)	0.0334 (5)
C35	0.2248 (4)	-0.0173 (3)	0.13501 (14)	0.0337 (6)
H35	0.1646	0.0889	0.1308	0.040*
C36	0.0403 (5)	-0.1038 (4)	0.24140 (15)	0.0464 (7)
H36A	0.0348	-0.0141	0.2589	0.056*
H36B	0.0895	-0.1933	0.2787	0.056*
C37	-0.1743 (5)	-0.0850 (4)	0.21924 (17)	0.0511 (8)
H37A	-0.2701	-0.0556	0.2559	0.061*
H37B	-0.2174	-0.0028	0.1786	0.061*
O3	-0.1834 (5)	-0.2206 (3)	0.20404 (12)	0.0682 (7)
H3	-0.1787	-0.2147	0.1620	0.102*
N41	0.3421 (3)	-0.3452 (2)	-0.06627 (11)	0.0318 (5)
N42	0.5298 (4)	-0.3745 (3)	-0.09471 (13)	0.0443 (6)
N43	0.6031 (4)	-0.2741 (3)	-0.08241 (13)	0.0406 (6)
N44	0.4653 (3)	-0.1786 (2)	-0.04598 (11)	0.0326 (5)
C45	0.3046 (4)	-0.2248 (3)	-0.03711 (14)	0.0346 (6)
H45	0.1844	-0.1801	-0.0141	0.041*
C46	0.2061 (5)	-0.4333 (3)	-0.07074 (14)	0.0377 (6)
H46A	0.0768	-0.3897	-0.0481	0.045*
H46B	0.2717	-0.5391	-0.0466	0.045*
C47	0.1617 (5)	-0.4311 (3)	-0.14470 (14)	0.0386 (6)
H47A	0.2876	-0.4887	-0.1655	0.046*
H47B	0.0607	-0.4807	-0.1459	0.046*
O4	0.0844 (3)	-0.2775 (2)	-0.18342 (11)	0.0432 (5)
H4	0.1605	-0.2673	-0.2161	0.065*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0253 (2)	0.0280 (2)	0.0256 (2)	-0.0109 (2)	0.00245 (19)	-0.00379 (19)
Cl1	0.0302 (3)	0.0385 (4)	0.0346 (3)	-0.0074 (3)	-0.0020 (3)	-0.0023 (3)
N11	0.0332 (12)	0.0315 (11)	0.0356 (12)	-0.0121 (10)	0.0050 (9)	-0.0098 (9)
N12	0.0347 (13)	0.0423 (14)	0.0682 (18)	-0.0114 (11)	0.0058 (12)	-0.0246 (13)
N13	0.0314 (13)	0.0444 (14)	0.0629 (17)	-0.0121 (11)	0.0078 (12)	-0.0223 (13)
N14	0.0310 (12)	0.0320 (12)	0.0363 (12)	-0.0118 (9)	0.0034 (9)	-0.0093 (9)
C15	0.0302 (14)	0.0343 (14)	0.0410 (15)	-0.0115 (11)	0.0062 (11)	-0.0147 (12)
C16	0.0406 (16)	0.0359 (15)	0.0484 (17)	-0.0213 (13)	0.0058 (13)	-0.0136 (13)
C17	0.0465 (18)	0.0400 (16)	0.0512 (18)	-0.0150 (14)	-0.0017 (14)	-0.0178 (14)
O1	0.0714 (16)	0.0409 (12)	0.0480 (13)	-0.0071 (11)	0.0100 (11)	-0.0142 (10)
N21	0.0338 (12)	0.0337 (12)	0.0284 (11)	-0.0146 (10)	0.0032 (9)	-0.0016 (9)
N22	0.0558 (16)	0.0366 (13)	0.0438 (14)	-0.0174 (12)	0.0207 (12)	-0.0105 (11)
N23	0.0475 (14)	0.0316 (12)	0.0446 (14)	-0.0139 (11)	0.0174 (11)	-0.0092 (10)
N24	0.0310 (11)	0.0307 (11)	0.0305 (11)	-0.0102 (9)	0.0078 (9)	-0.0057 (9)
C25	0.0343 (14)	0.0309 (14)	0.0384 (15)	-0.0074 (11)	0.0066 (12)	-0.0005 (11)
C26	0.0446 (17)	0.0445 (16)	0.0321 (14)	-0.0195 (13)	0.0014 (12)	0.0074 (12)
C27	0.057 (2)	0.0396 (16)	0.0419 (17)	-0.0244 (15)	0.0012 (14)	0.0018 (13)
O2	0.0466 (12)	0.0555 (13)	0.0414 (12)	-0.0228 (11)	0.0020 (10)	0.0001 (10)
Co2	0.0272 (3)	0.0275 (3)	0.0326 (3)	-0.0113 (2)	0.0062 (2)	-0.0092 (2)
Cl2	0.0288 (3)	0.0396 (4)	0.0415 (4)	-0.0086 (3)	0.0040 (3)	-0.0082 (3)
N31	0.0374 (13)	0.0353 (12)	0.0340 (12)	-0.0121 (10)	0.0061 (10)	-0.0018 (10)
N32	0.0539 (16)	0.0317 (13)	0.0546 (16)	-0.0084 (12)	0.0135 (13)	0.0024 (11)
N33	0.0461 (15)	0.0292 (12)	0.0563 (16)	-0.0068 (11)	0.0165 (12)	-0.0026 (11)
N34	0.0327 (12)	0.0285 (11)	0.0373 (12)	-0.0105 (9)	0.0061 (9)	-0.0044 (9)
C35	0.0362 (14)	0.0294 (13)	0.0345 (14)	-0.0120 (11)	0.0069 (11)	-0.0054 (11)
C36	0.0562 (19)	0.0509 (18)	0.0338 (15)	-0.0231 (15)	0.0149 (14)	-0.0082 (13)
C37	0.0527 (19)	0.057 (2)	0.0509 (19)	-0.0274 (16)	0.0196 (15)	-0.0163 (16)
O3	0.106 (2)	0.0774 (17)	0.0483 (14)	-0.0632 (17)	0.0223 (15)	-0.0207 (13)
N41	0.0370 (12)	0.0334 (12)	0.0285 (11)	-0.0157 (10)	0.0021 (9)	-0.0080 (9)
N42	0.0465 (15)	0.0453 (14)	0.0502 (15)	-0.0205 (12)	0.0122 (12)	-0.0242 (12)
N43	0.0400 (13)	0.0427 (13)	0.0458 (14)	-0.0188 (11)	0.0124 (11)	-0.0189 (11)
N44	0.0311 (12)	0.0354 (12)	0.0348 (12)	-0.0129 (10)	0.0043 (9)	-0.0130 (10)
C45	0.0324 (14)	0.0365 (14)	0.0391 (15)	-0.0133 (12)	0.0056 (11)	-0.0162 (12)
C46	0.0491 (17)	0.0352 (14)	0.0354 (14)	-0.0233 (13)	0.0021 (12)	-0.0066 (11)
C47	0.0484 (17)	0.0319 (14)	0.0384 (15)	-0.0169 (13)	-0.0031 (13)	-0.0070 (12)
O4	0.0442 (12)	0.0343 (10)	0.0431 (12)	-0.0087 (9)	0.0006 (9)	0.0006 (9)

Geometric parameters (\AA , $^\circ$)

Co1—N24	2.144 (2)	Co2—N44 ⁱⁱ	2.165 (2)
Co1—N24 ⁱ	2.144 (2)	Co2—N44	2.165 (2)
Co1—N14 ⁱ	2.191 (2)	Co2—N34 ⁱⁱ	2.168 (2)
Co1—N14	2.191 (2)	Co2—N34	2.168 (2)
Co1—Cl1 ⁱ	2.4372 (10)	Co2—Cl2 ⁱⁱ	2.4333 (10)

supplementary materials

Co1—C11	2.4372 (10)	Co2—C12	2.4333 (10)
N11—C15	1.326 (3)	N31—C35	1.325 (3)
N11—N12	1.348 (3)	N31—N32	1.345 (3)
N11—C16	1.469 (3)	N31—C36	1.467 (3)
N12—N13	1.297 (3)	N32—N33	1.292 (3)
N13—N14	1.366 (3)	N33—N34	1.359 (3)
N14—C15	1.308 (3)	N34—C35	1.312 (3)
C15—H15	0.9300	C35—H35	0.9300
C16—C17	1.504 (4)	C36—C37	1.507 (4)
C16—H16A	0.9700	C36—H36A	0.9700
C16—H16B	0.9700	C36—H36B	0.9700
C17—O1	1.410 (3)	C37—O3	1.400 (4)
C17—H17A	0.9700	C37—H37A	0.9700
C17—H17B	0.9700	C37—H37B	0.9700
O1—H1	0.8200	O3—H3	0.8200
N21—C25	1.320 (3)	N41—C45	1.322 (3)
N21—N22	1.346 (3)	N41—N42	1.342 (3)
N21—C26	1.469 (3)	N41—C46	1.474 (3)
N22—N23	1.290 (3)	N42—N43	1.290 (3)
N23—N24	1.356 (3)	N43—N44	1.353 (3)
N24—C25	1.316 (3)	N44—C45	1.318 (3)
C25—H25	0.9300	C45—H45	0.9300
C26—C27	1.500 (4)	C46—C47	1.505 (4)
C26—H26A	0.9700	C46—H46A	0.9700
C26—H26B	0.9700	C46—H46B	0.9700
C27—O2	1.419 (4)	C47—O4	1.421 (3)
C27—H27A	0.9700	C47—H47A	0.9700
C27—H27B	0.9700	C47—H47B	0.9700
O2—H2	0.8200	O4—H4	0.8200
N24—Co1—N24 ⁱ	180.00 (10)	N44 ⁱⁱ —Co2—N44	180.000 (1)
N24—Co1—N14 ⁱ	92.41 (8)	N44 ⁱⁱ —Co2—N34 ⁱⁱ	88.52 (8)
N24 ⁱ —Co1—N14 ⁱ	87.59 (8)	N44—Co2—N34 ⁱⁱ	91.48 (8)
N24—Co1—N14	87.59 (8)	N44 ⁱⁱ —Co2—N34	91.48 (8)
N24 ⁱ —Co1—N14	92.41 (8)	N44—Co2—N34	88.52 (8)
N14 ⁱ —Co1—N14	180.0	N34 ⁱⁱ —Co2—N34	180.00 (8)
N24—Co1—C11 ⁱ	91.04 (6)	N44 ⁱⁱ —Co2—C12 ⁱⁱ	90.56 (6)
N24 ⁱ —Co1—C11 ⁱ	88.96 (6)	N44—Co2—C12 ⁱⁱ	89.45 (6)
N14 ⁱ —Co1—C11 ⁱ	91.03 (6)	N34 ⁱⁱ —Co2—C12 ⁱⁱ	90.42 (6)
N14—Co1—C11 ⁱ	88.97 (6)	N34—Co2—C12 ⁱⁱ	89.58 (6)
N24—Co1—C11	88.96 (6)	N44 ⁱⁱ —Co2—C12	89.45 (6)
N24 ⁱ —Co1—C11	91.04 (6)	N44—Co2—C12	90.55 (6)
N14 ⁱ —Co1—C11	88.97 (6)	N34 ⁱⁱ —Co2—C12	89.58 (6)
N14—Co1—C11	91.03 (6)	N34—Co2—C12	90.42 (6)
C11 ⁱ —Co1—C11	180.00 (2)	C12 ⁱⁱ —Co2—C12	180.0
C15—N11—N12	108.2 (2)	C35—N31—N32	108.4 (2)
C15—N11—C16	128.7 (2)	C35—N31—C36	130.1 (2)

N12—N11—C16	123.1 (2)	N32—N31—C36	121.4 (2)
N13—N12—N11	106.7 (2)	N33—N32—N31	106.8 (2)
N12—N13—N14	109.9 (2)	N32—N33—N34	109.9 (2)
C15—N14—N13	105.9 (2)	C35—N34—N33	106.3 (2)
C15—N14—Co1	124.65 (18)	C35—N34—Co2	131.63 (18)
N13—N14—Co1	129.21 (17)	N33—N34—Co2	122.09 (17)
N14—C15—N11	109.3 (2)	N34—C35—N31	108.7 (2)
N14—C15—H15	125.4	N34—C35—H35	125.7
N11—C15—H15	125.4	N31—C35—H35	125.7
N11—C16—C17	111.7 (2)	N31—C36—C37	112.1 (3)
N11—C16—H16A	109.3	N31—C36—H36A	109.2
C17—C16—H16A	109.3	C37—C36—H36A	109.2
N11—C16—H16B	109.3	N31—C36—H36B	109.2
C17—C16—H16B	109.3	C37—C36—H36B	109.2
H16A—C16—H16B	107.9	H36A—C36—H36B	107.9
O1—C17—C16	111.5 (3)	O3—C37—C36	112.0 (3)
O1—C17—H17A	109.3	O3—C37—H37A	109.2
C16—C17—H17A	109.3	C36—C37—H37A	109.2
O1—C17—H17B	109.3	O3—C37—H37B	109.2
C16—C17—H17B	109.3	C36—C37—H37B	109.2
H17A—C17—H17B	108.0	H37A—C37—H37B	107.9
C17—O1—H1	109.5	C37—O3—H3	109.5
C25—N21—N22	108.5 (2)	C45—N41—N42	108.1 (2)
C25—N21—C26	129.8 (2)	C45—N41—C46	128.3 (2)
N22—N21—C26	121.5 (2)	N42—N41—C46	123.6 (2)
N23—N22—N21	106.7 (2)	N43—N42—N41	107.2 (2)
N22—N23—N24	110.2 (2)	N42—N43—N44	109.9 (2)
C25—N24—N23	105.9 (2)	C45—N44—N43	106.0 (2)
C25—N24—Co1	130.62 (18)	C45—N44—Co2	124.45 (17)
N23—N24—Co1	123.40 (16)	N43—N44—Co2	129.29 (17)
N24—C25—N21	108.8 (2)	N44—C45—N41	108.9 (2)
N24—C25—H25	125.6	N44—C45—H45	125.6
N21—C25—H25	125.6	N41—C45—H45	125.6
N21—C26—C27	111.3 (2)	N41—C46—C47	111.7 (2)
N21—C26—H26A	109.4	N41—C46—H46A	109.3
C27—C26—H26A	109.4	C47—C46—H46A	109.3
N21—C26—H26B	109.4	N41—C46—H46B	109.3
C27—C26—H26B	109.4	C47—C46—H46B	109.3
H26A—C26—H26B	108.0	H46A—C46—H46B	108.0
O2—C27—C26	111.9 (3)	O4—C47—C46	110.8 (2)
O2—C27—H27A	109.2	O4—C47—H47A	109.5
C26—C27—H27A	109.2	C46—C47—H47A	109.5
O2—C27—H27B	109.2	O4—C47—H47B	109.5
C26—C27—H27B	109.2	C46—C47—H47B	109.5
H27A—C27—H27B	107.9	H47A—C47—H47B	108.1
C27—O2—H2	109.5	C47—O4—H4	109.5
C15—N11—N12—N13	0.2 (3)	C35—N31—N32—N33	0.2 (3)
C16—N11—N12—N13	-179.0 (2)	C36—N31—N32—N33	177.3 (3)
N11—N12—N13—N14	0.0 (3)	N31—N32—N33—N34	-0.1 (4)

supplementary materials

N12—N13—N14—C15	-0.2 (3)	N32—N33—N34—C35	-0.1 (3)
N12—N13—N14—Co1	-175.40 (19)	N32—N33—N34—Co2	-178.8 (2)
N24—Co1—N14—C15	-53.8 (2)	N44 ⁱⁱ —Co2—N34—C35	41.8 (3)
N24 ⁱ —Co1—N14—C15	126.2 (2)	N44—Co2—N34—C35	-138.2 (3)
Cl1 ⁱ —Co1—N14—C15	37.3 (2)	Cl2 ⁱⁱ —Co2—N34—C35	-48.7 (3)
Cl1—Co1—N14—C15	-142.7 (2)	Cl2—Co2—N34—C35	131.3 (3)
N24—Co1—N14—N13	120.6 (2)	N44 ⁱⁱ —Co2—N34—N33	-139.8 (2)
N24 ⁱ —Co1—N14—N13	-59.4 (2)	N44—Co2—N34—N33	40.2 (2)
Cl1 ⁱ —Co1—N14—N13	-148.4 (2)	Cl2 ⁱⁱ —Co2—N34—N33	129.6 (2)
Cl1—Co1—N14—N13	31.6 (2)	Cl2—Co2—N34—N33	-50.4 (2)
N13—N14—C15—N11	0.4 (3)	N33—N34—C35—N31	0.2 (3)
Co1—N14—C15—N11	175.83 (16)	Co2—N34—C35—N31	178.73 (19)
N12—N11—C15—N14	-0.4 (3)	N32—N31—C35—N34	-0.2 (3)
C16—N11—C15—N14	178.8 (2)	C36—N31—C35—N34	-177.0 (3)
C15—N11—C16—C17	-101.2 (3)	C35—N31—C36—C37	84.0 (4)
N12—N11—C16—C17	77.8 (3)	N32—N31—C36—C37	-92.4 (3)
N11—C16—C17—O1	-70.2 (3)	N31—C36—C37—O3	68.6 (3)
C25—N21—N22—N23	-0.1 (3)	C45—N41—N42—N43	0.3 (3)
C26—N21—N22—N23	175.4 (3)	C46—N41—N42—N43	178.5 (2)
N21—N22—N23—N24	0.2 (3)	N41—N42—N43—N44	0.0 (3)
N22—N23—N24—C25	-0.2 (3)	N42—N43—N44—C45	-0.4 (3)
N22—N23—N24—Co1	177.84 (19)	N42—N43—N44—Co2	173.47 (19)
N14 ⁱ —Co1—N24—C25	131.3 (3)	N34 ⁱⁱ —Co2—N44—C45	-141.5 (2)
N14—Co1—N24—C25	-48.7 (3)	N34—Co2—N44—C45	38.5 (2)
Cl1 ⁱ —Co1—N24—C25	-137.6 (3)	Cl2 ⁱⁱ —Co2—N44—C45	-51.1 (2)
Cl1—Co1—N24—C25	42.4 (3)	Cl2—Co2—N44—C45	128.9 (2)
N14 ⁱ —Co1—N24—N23	-46.2 (2)	N34 ⁱⁱ —Co2—N44—N43	45.6 (2)
N14—Co1—N24—N23	133.8 (2)	N34—Co2—N44—N43	-134.4 (2)
Cl1 ⁱ —Co1—N24—N23	44.8 (2)	Cl2 ⁱⁱ —Co2—N44—N43	136.0 (2)
Cl1—Co1—N24—N23	-135.2 (2)	Cl2—Co2—N44—N43	-44.0 (2)
N23—N24—C25—N21	0.2 (3)	N43—N44—C45—N41	0.6 (3)
Co1—N24—C25—N21	-177.68 (18)	Co2—N44—C45—N41	-173.65 (17)
N22—N21—C25—N24	-0.1 (3)	N42—N41—C45—N44	-0.6 (3)
C26—N21—C25—N24	-175.1 (3)	C46—N41—C45—N44	-178.6 (2)
C25—N21—C26—C27	89.6 (4)	C45—N41—C46—C47	120.9 (3)
N22—N21—C26—C27	-84.9 (3)	N42—N41—C46—C47	-56.9 (3)
N21—C26—C27—O2	65.8 (3)	N41—C46—C47—O4	-53.2 (3)

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x+1, -y, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots O3 ⁱ	0.82	1.94	2.741 (3)	165
O2—H2 \cdots Cl1 ⁱⁱⁱ	0.82	2.32	3.104 (2)	160
O3—H3 \cdots Cl2 ^{iv}	0.82	2.28	3.098 (3)	173
O4—H4 \cdots O2 ⁱⁱ	0.82	1.94	2.757 (3)	175

C15—H15...C11 ⁱⁱⁱ	0.93	2.76	3.482 (3)	135
C25—H25...O1 ^v	0.93	2.56	3.303 (4)	137
C35—H35...O4 ^{vi}	0.93	2.38	3.156 (3)	141
C45—H45...C12 ^{iv}	0.93	2.54	3.405 (3)	155

Symmetry codes: (i) $-x, -y, -z+1$; (iii) $x+1, y, z$; (iv) $x-1, y, z$; (ii) $-x+1, -y, -z$; (v) $-x, -y+1, -z+1$; (vi) $-x, -y, -z$.

Fig. 1




Fig. 2




Fig. 3

