

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

A new Schiff base nickel(II) complex: [5,5'-dihydroxy-2,2'-[o-phenylenebis-(nitrilomethylidene)]diphenolato}-nickel(II) methanol disolvate

Meiju Niu,* Guihua Liu, Daqi Wang and Jianmin Dou

College of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China

Correspondence e-mail: niumeiju@163.com


Received 7 September 2009; accepted 6 October 2009

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.040; wR factor = 0.081; data-to-parameter ratio = 12.7.

The monomeric title nickel(II) complex of a salicylaldehyde, $[\text{Ni}(\text{C}_{20}\text{H}_{14}\text{N}_2\text{O}_4)] \cdot 2\text{CH}_3\text{OH}$, was obtained by the reaction of 2,4-dihydroxybenzaldehyde and 1,2-phenylenediamine with nickel(II) acetate. The Ni^{II} atom is coordinated by two N atoms [$\text{Ni}-\text{N} = 1.839$ (2) Å] and two O atoms [$\text{Ni}-\text{O} = 1.8253$ (19) Å] in an approximately square-planar geometry. In the crystal structure, intermolecular $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds link the molecules into a chain along [001].

Related literature

For related structures, see: Amirnasr *et al.* (2006); Shi *et al.* (2004); Chen *et al.* (2009); Hermindez-Molina *et al.* (1997); Zhang *et al.* (2009).



Experimental

Crystal data

 $[\text{Ni}(\text{C}_{20}\text{H}_{14}\text{N}_2\text{O}_4)] \cdot 2\text{CH}_4\text{O}$
 $M_r = 469.13$

 Monoclinic, $C2/c$
 $a = 15.673$ (3) Å
 $b = 15.090$ (2) Å
 $c = 8.8680$ (2) Å
 $\beta = 104.593$ (3)°
 $V = 2029.7$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.00$ mm⁻¹
 $T = 298$ K
 $0.31 \times 0.14 \times 0.13$ mm

Data collection

 Siemens SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.747$, $T_{\text{max}} = 0.881$

 5206 measured reflections
 1788 independent reflections
 1333 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.081$
 $S = 1.00$
 1788 reflections

 141 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.35$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.35$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

| $D-H \cdots A$ | $D-H$ | $H \cdots A$ | $D \cdots A$ | $D-H \cdots A$ |
|--|-------|--------------|--------------|----------------|
| $\text{O2}-\text{H2} \cdots \text{O3}^{\text{i}}$ | 0.82 | 1.97 | 2.734 (3) | 154 |
| $\text{O3}-\text{H3} \cdots \text{O1}^{\text{ii}}$ | 0.82 | 1.98 | 2.797 (3) | 172 |

 Symmetry codes: (i) $x, -y + 1, z - \frac{1}{2}$; (ii) $x, y, z + 1$.

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We acknowledge the financial support of the National Natural Science Foundation of China (20671048).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DS2006).

References

- Amirnasr, M., Schenk, K. J., Meghdadi, S. & Morshedi, M. (2006). *Polyhedron*, **25**, 671–677.
- Chen, X. H., Liang, Z. Y., Zhan, C. R. & Yang, M. X. (2009). *J. Fujian Branch Fujian Normal Univ.* **2**, 7–11.
- Hermindez-Molina, R., Mederos, A., Gill, P., Domfnguez, S., Ndfiez, P., Germain, G. & Debaerdemaeker, T. (1997). *Inorg. Chim. Acta*, **256**, 319–325.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shi, Y. C., Shen, W. B., Yang, H. M., Song, H. B. & Hu, X. Y. (2004). *Polyhedron*, **23**, 749–754.
- Siemens (1996). *SMART* and *SAINTE*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Zhang, Q. F., Jiang, F. L., Huang, Y. G., Wei, W., Gao, Q., Yang, M., Xiong, K. C. & Hong, M. C. (2009). *Dalton Trans.* pp. 2673–2676.

supplementary materials

Acta Cryst. (2009). E65, m1357 [doi:10.1107/S160053680904063X]

A new Schiff base nickel(II) complex: {5,5'-dihydroxy-2,2'-[*o*-phenylenebis(nitrilomethyldyne)]diphenolato}nickel(II) methanol disolvate

M. Niu, G. Liu, D. Wang and J. Dou

Comment

Nickel complexes have attracted intensive interest in the past decade because they play important roles in bioinorganic chemistry and redox enzyme systems (Amirnasr *et al.*, 2006). In a continuation of a study of Schiff base ligands and their nickel(II) complexes, we report here the title complex (Fig. 1), in which the main plane being formed by the three phenyl and the N₂O₂. The angles O1—Ni1—N1A and O1A—Ni1—N1 (177.30 (10)°) indicate that the coordination geometry of the nickel atom is four-coordinate in an approximately square planar, which acts as a tetradentate ligand through its *o*-phenylenediamine N atoms and its deprotonated phenol O atoms. This square planar geometry is the most usual for Ni^{II} complexes (Shi *et al.*, 2004) in the N₂O₂ donor set with Schiff base ligands. The Ni—O distances of 1.8253 (19) Å are very close to the corresponding values in related structures (1.820 Å, Chen *et al.*, 2009). However, the Ni—N distances of 1.8392 (2) Å are significantly shorter than that for a related complex (1.859 Å, Hemindez-Molina *et al.*, 1997). As shown in Fig. 2, intermolecular O—H...O hydrogen bonds (Table 1) link the molecules into a one-dimensional chain along [0 0 1] direction (Zhang *et al.*, 2009).

Experimental

o-Phenylenediamine (1 mmol, 108.22 mg) was dissolved in hot methanol (20 ml) and added dropwise to a methanol solution (10 ml) of 2,4-dihydroxybenzaldehyde (2 mmol, 276.2 mg). The mixture was then stirred at 323 K for 4 h. The triethylamine solution (3 ml) of nickel (II) acetate (1.5 mmol, 292.2 mg) was then added dropwise and the mixture stirred for another 4 h, at which point a red precipitate collected by suction filtration and washed with ethanol and ether. Crystals of the title compound suitable for X-ray analysis were from the methanol and dimethylsulfoxide solution after about one week.

Refinement

All H atoms were placed geometrically and treated as riding on their parent atoms with C—H = 0.96 Å (methylene) or 0.93 Å (aromatic), 0.82 Å (hydroxyl) and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the compound, with atom labels and 30% probability displacement ellipsoids for non-H atoms.




Fig. 2. Crystal packing of the compound, showing a one-dimensional chain linked by O—H...O hydrogen bonds (dashed lines).

{5,5'-dihydroxy-2,2'-[o-phenylenebis(nitrilomethylidyne)]diphenolato}nickel(II) methanol disolvate

Crystal data

[Ni(C₂₀H₁₄N₂O₄)]·2CH₄O

M_r = 469.13

Monoclinic, *C*2/*c*

Hall symbol: -*C* 2yc

a = 15.673 (3) Å

b = 15.090 (2) Å

c = 8.8680 (2) Å

β = 104.593 (3)°

V = 2029.7 (5) Å³

Z = 4

*F*₀₀₀ = 976

D_x = 1.535 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 1287 reflections

θ = 2.7–23.9°

μ = 1.00 mm⁻¹

T = 298 K

Block, red

0.31 × 0.14 × 0.13 mm

Data collection

Siemens SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 298 K

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

*T*_{min} = 0.747, *T*_{max} = 0.881

5206 measured reflections

1788 independent reflections

1333 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.049

θ_{max} = 25.0°

θ_{min} = 1.9°

h = -15→18

k = -17→16

l = -9→10

Refinement

Refinement on *F*²

Least-squares matrix: full

R [*F*² > 2σ(*F*²)] = 0.040

wR(*F*²) = 0.081

S = 1.00

1788 reflections

141 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F*_o²) + (0.0294*P*)²]

where *P* = (*F*_o² + 2*F*_c²)/3

(Δ/σ)_{max} < 0.001

Δρ_{max} = 0.35 e Å⁻³

Δρ_{min} = -0.35 e Å⁻³

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|--------------|---------------|-------------|----------------------------------|
| Ni1 | 0.0000 | 0.21153 (3) | -0.2500 | 0.03428 (19) |
| N1 | 0.06329 (15) | 0.12289 (14) | -0.1294 (3) | 0.0322 (6) |
| O1 | 0.06161 (13) | 0.30290 (12) | -0.1384 (2) | 0.0407 (5) |
| O2 | 0.23991 (13) | 0.45598 (13) | 0.2855 (2) | 0.0538 (6) |
| H2 | 0.2136 | 0.4976 | 0.2348 | 0.081* |
| O3 | 0.11150 (15) | 0.42336 (14) | 0.6598 (3) | 0.0615 (7) |
| H3 | 0.0921 | 0.3895 | 0.7154 | 0.092* |
| C1 | 0.03571 (17) | 0.03697 (17) | -0.1846 (3) | 0.0340 (7) |
| C2 | 0.0716 (2) | -0.04279 (19) | -0.1201 (4) | 0.0438 (8) |
| H2A | 0.1194 | -0.0433 | -0.0332 | 0.053* |
| C3 | 0.0357 (2) | -0.12082 (19) | -0.1861 (4) | 0.0476 (9) |
| H3A | 0.0598 | -0.1744 | -0.1441 | 0.057* |
| C4 | 0.12589 (19) | 0.1350 (2) | -0.0019 (4) | 0.0384 (8) |
| H4 | 0.1529 | 0.0847 | 0.0494 | 0.046* |
| C5 | 0.15581 (18) | 0.21777 (19) | 0.0639 (3) | 0.0330 (7) |
| C6 | 0.12162 (18) | 0.2986 (2) | -0.0053 (3) | 0.0352 (7) |
| C7 | 0.1515 (2) | 0.3782 (2) | 0.0694 (4) | 0.0417 (8) |
| H7 | 0.1292 | 0.4315 | 0.0235 | 0.050* |
| C8 | 0.21309 (19) | 0.3788 (2) | 0.2092 (4) | 0.0385 (7) |
| C9 | 0.2494 (2) | 0.3000 (2) | 0.2786 (4) | 0.0461 (8) |
| H9 | 0.2925 | 0.3007 | 0.3723 | 0.055* |
| C10 | 0.22071 (19) | 0.2219 (2) | 0.2066 (4) | 0.0434 (8) |
| H10 | 0.2448 | 0.1693 | 0.2533 | 0.052* |
| C11 | 0.0943 (3) | 0.3892 (2) | 0.5095 (5) | 0.0717 (11) |
| H11A | 0.0431 | 0.4176 | 0.4454 | 0.107* |
| H11B | 0.0839 | 0.3266 | 0.5123 | 0.107* |
| H11C | 0.1440 | 0.3997 | 0.4671 | 0.107* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|------------|------------|------------|----------|------------|----------|
| Ni1 | 0.0358 (3) | 0.0298 (3) | 0.0330 (3) | 0.000 | 0.0007 (2) | 0.000 |

supplementary materials

| | | | | | | |
|-----|-------------|-------------|-------------|--------------|--------------|--------------|
| N1 | 0.0334 (14) | 0.0260 (13) | 0.0370 (15) | 0.0003 (11) | 0.0083 (12) | 0.0010 (12) |
| O1 | 0.0474 (13) | 0.0288 (11) | 0.0357 (12) | -0.0001 (10) | -0.0088 (10) | 0.0008 (10) |
| O2 | 0.0578 (14) | 0.0451 (13) | 0.0460 (15) | -0.0056 (11) | -0.0101 (11) | -0.0067 (12) |
| O3 | 0.0716 (17) | 0.0518 (15) | 0.0575 (17) | -0.0104 (12) | 0.0094 (13) | 0.0108 (13) |
| C1 | 0.0379 (18) | 0.0288 (16) | 0.0376 (19) | 0.0023 (13) | 0.0138 (13) | -0.0020 (14) |
| C2 | 0.047 (2) | 0.0360 (18) | 0.046 (2) | 0.0054 (16) | 0.0088 (16) | 0.0085 (16) |
| C3 | 0.061 (2) | 0.0294 (16) | 0.054 (2) | 0.0048 (15) | 0.0172 (16) | 0.0030 (15) |
| C4 | 0.0375 (18) | 0.0372 (18) | 0.040 (2) | 0.0054 (15) | 0.0081 (15) | 0.0062 (16) |
| C5 | 0.0325 (16) | 0.0320 (16) | 0.0330 (16) | 0.0017 (15) | 0.0053 (13) | -0.0008 (15) |
| C6 | 0.0319 (16) | 0.0385 (18) | 0.0326 (17) | -0.0002 (15) | 0.0035 (13) | -0.0006 (15) |
| C7 | 0.046 (2) | 0.0350 (17) | 0.038 (2) | -0.0007 (15) | -0.0007 (15) | -0.0017 (15) |
| C8 | 0.0373 (18) | 0.0395 (18) | 0.0364 (19) | -0.0043 (15) | 0.0051 (15) | -0.0070 (16) |
| C9 | 0.0395 (18) | 0.054 (2) | 0.0352 (19) | 0.0037 (17) | -0.0088 (14) | -0.0008 (17) |
| C10 | 0.0420 (19) | 0.0410 (19) | 0.0409 (19) | 0.0068 (16) | -0.0014 (15) | 0.0055 (17) |
| C11 | 0.081 (3) | 0.064 (3) | 0.072 (3) | -0.010 (2) | 0.025 (2) | 0.000 (2) |

Geometric parameters (Å, °)

| | | | |
|--------------------------------------|-------------|-----------|-----------|
| Ni1—O1 | 1.8253 (19) | C3—H3A | 0.9300 |
| Ni1—O1 ⁱ | 1.8253 (19) | C4—C5 | 1.408 (4) |
| Ni1—N1 ⁱ | 1.839 (2) | C4—H4 | 0.9300 |
| Ni1—N1 | 1.839 (2) | C5—C6 | 1.409 (4) |
| N1—C4 | 1.310 (4) | C5—C10 | 1.411 (4) |
| N1—C1 | 1.414 (3) | C6—C7 | 1.395 (4) |
| O1—C6 | 1.312 (3) | C7—C8 | 1.366 (4) |
| O2—C8 | 1.360 (3) | C7—H7 | 0.9300 |
| O2—H2 | 0.8200 | C8—C9 | 1.393 (4) |
| O3—C11 | 1.391 (4) | C9—C10 | 1.361 (4) |
| O3—H3 | 0.8200 | C9—H9 | 0.9300 |
| C1—C2 | 1.389 (3) | C10—H10 | 0.9300 |
| C1—C1 ⁱ | 1.394 (6) | C11—H11A | 0.9600 |
| C2—C3 | 1.371 (4) | C11—H11B | 0.9600 |
| C2—H2A | 0.9300 | C11—H11C | 0.9600 |
| C3—C3 ⁱ | 1.377 (6) | | |
| O1—Ni1—O1 ⁱ | 81.88 (12) | C4—C5—C6 | 122.5 (3) |
| O1—Ni1—N1 ⁱ | 177.30 (10) | C4—C5—C10 | 120.1 (3) |
| O1 ⁱ —Ni1—N1 ⁱ | 95.74 (9) | C6—C5—C10 | 117.5 (3) |
| O1—Ni1—N1 | 95.74 (9) | O1—C6—C7 | 117.6 (3) |
| O1 ⁱ —Ni1—N1 | 177.30 (10) | O1—C6—C5 | 122.9 (3) |
| N1 ⁱ —Ni1—N1 | 86.67 (15) | C7—C6—C5 | 119.6 (3) |
| C4—N1—C1 | 121.6 (3) | C8—C7—C6 | 120.8 (3) |
| C4—N1—Ni1 | 125.3 (2) | C8—C7—H7 | 119.6 |
| C1—N1—Ni1 | 113.11 (19) | C6—C7—H7 | 119.6 |
| C6—O1—Ni1 | 127.69 (19) | O2—C8—C7 | 121.1 (3) |
| C8—O2—H2 | 109.5 | O2—C8—C9 | 118.1 (3) |
| C11—O3—H3 | 109.5 | C7—C8—C9 | 120.8 (3) |
| C2—C1—C1 ⁱ | 119.95 (18) | C10—C9—C8 | 118.8 (3) |

| | | | |
|-------------------------|-------------|---------------|-----------|
| C2—C1—N1 | 126.5 (3) | C10—C9—H9 | 120.6 |
| C1 ⁱ —C1—N1 | 113.53 (15) | C8—C9—H9 | 120.6 |
| C3—C2—C1 | 119.2 (3) | C9—C10—C5 | 122.5 (3) |
| C3—C2—H2A | 120.4 | C9—C10—H10 | 118.7 |
| C1—C2—H2A | 120.4 | C5—C10—H10 | 118.7 |
| C2—C3—C3 ⁱ | 120.81 (18) | O3—C11—H11A | 109.5 |
| C2—C3—H3A | 119.6 | O3—C11—H11B | 109.5 |
| C3 ⁱ —C3—H3A | 119.6 | H11A—C11—H11B | 109.5 |
| N1—C4—C5 | 125.5 (3) | O3—C11—H11C | 109.5 |
| N1—C4—H4 | 117.2 | H11A—C11—H11C | 109.5 |
| C5—C4—H4 | 117.2 | H11B—C11—H11C | 109.5 |

Symmetry codes: (i) $-x, y, -z-1/2$.

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|----------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| O2—H2 \cdots O3 ⁱⁱ | 0.82 | 1.97 | 2.734 (3) | 154 |
| O3—H3 \cdots O1 ⁱⁱⁱ | 0.82 | 1.98 | 2.797 (3) | 172 |

Symmetry codes: (ii) $x, -y+1, z-1/2$; (iii) $x, y, z+1$.

Fig. 1




Fig. 2

