

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Poly[[bis(μ_2 -6-methylpyrazin-2-carboxylato- $\kappa^3 N^1, O: N^4$)copper(II)] dihydrate]

Chuan-gang Fan,^a Xin-Ting Wei^{b*} and Linwei Li^b

^aDepartment of Chemistry and Chemical Engineering, Binzhou University, Shandong 256603, People's Republic of China, and ^bCollege of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China
Correspondence e-mail: xintingwei@cn.yahoo.com


Received 29 September 2009; accepted 11 October 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.008$ Å; disorder in main residue; R factor = 0.049; wR factor = 0.160; data-to-parameter ratio = 11.0.

In the title compound, $\{[Cu(C_6H_5N_2O_2)_2] \cdot 2H_2O\}_n$, the Cu^{II} ion (site symmetry $\bar{1}$) is coordinated by two N,O -bidentate ligands and two N -monodentate ligands in a distorted CuO_2N_4 octahedral geometry. Each anion acts as a bridge between two cations, thus forming a two-dimensional polymeric network parallel to the ab plane. The packing is consolidated by $O-H \cdots O$ hydrogen bonds. One of the O atoms of the ligand and both water molecules are disordered.

Related literature

For a related structure, see: Yigit *et al.* (2006). For background to coordination networks, see: Kesanli & Lin (2003); Barnett & Champness (2003).



Experimental

Crystal data

$[Cu(C_6H_5N_2O_2)_2] \cdot 2H_2O$
 $M_r = 373.81$
Monoclinic, $P2_1/c$

$a = 8.371$ (1) Å
 $b = 9.7901$ (11) Å
 $c = 10.3849$ (13) Å

$\beta = 112.277$ (1)°
 $V = 787.55$ (16) Å³
 $Z = 2$
Mo $K\alpha$ radiation

$\mu = 1.42$ mm⁻¹
 $T = 298$ K
 $0.34 \times 0.32 \times 0.30$ mm

Data collection

Bruker SMART CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2003)
 $T_{min} = 0.644, T_{max} = 0.675$

3821 measured reflections
1388 independent reflections
1094 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.160$
 $S = 1.06$
1388 reflections

126 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 1.23$ e Å⁻³
 $\Delta\rho_{min} = -0.47$ e Å⁻³

Table 1
Selected bond lengths (Å).

Cu1—O1	1.949 (3)	Cu1—N1	2.354 (4)
Cu1—N2 ⁱ	2.064 (4)		

Symmetry code: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Table 2
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O3—H3C ⁱ ···O2 ⁱⁱ	0.85	1.78	2.624 (3)	175
O3—H3C ⁱ ···O2 ⁱⁱⁱ	0.85	2.36	3.181 (3)	163
O3—H3D ⁱ ···O2 ^{iv}	0.85	2.23	3.074 (2)	175
O4—H4D ⁱ ···O3 ⁱⁱⁱ	0.85	1.97	2.73 (3)	147
O4—H4E ⁱ ···O3 ^{iv}	0.85	2.01	2.77 (2)	150

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x + 1, y, z$; (iii) $x, y, z + 1$; (iv) $-x + 1, -y + 1, -z + 1$.

Data collection: SMART (Bruker, 2003); cell refinement: SAINT (Bruker, 2003); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5123).

References

Barnett, S. A. & Champness, N. R. (2003). *Coord. Chem. Rev.* **246**, 145–168.
Bruker (2003). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Kesanli, B. & Lin, W. B. (2003). *Coord. Chem. Rev.* **246**, 305–326.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Yigit, M. V., Wang, Y., Moulton, B. & MacDonald, J. C. (2006). *Cryst. Growth Des.* **6**, 829–832.

supplementary materials

Acta Cryst. (2009). E65, m1412 [doi:10.1107/S1600536809041440]

Poly[[bis(μ_2 -6-methylpyrazin-2-carboxylato- $\kappa^3 N^1, O: N^4$)copper(II)] dihydrate]

C. Fan, X.-T. Wei and L. Li

Comment

In recent years, the construction of metal-organic coordination polymers (MOCPs) by metal-directed self-assembly is of great interest not only for their potential applications as functional materials in ion exchange, catalysis, hydrogen storage, and magnetic devices, but also for their aesthetic structural and host-guest chemistry associated with large central cavities. (Kesanli *et al.*, 2003; Barnett *et al.*, 2003). As an extension of this work, the 6-methyl-2-pyrazinecarboxylic acid was chosen due to its chelating coordinated effect leading to a linear metal center.

As shown in Fig. 1. X-ray structural analyses of complex (1) reveal the core structure of (1) is the symmetric dinuclear unit of $[\text{Cu}(L)_2(\text{H}_2\text{O})_2]$ (L = anion of 6-methyl-2-pyrazinecarboxylic acid). Each Cu^{II} atom is coordinated to two ligands as well as to the two N atoms of the other two ligands, forming an octahedral coordination conformation.

The ligand forms a coordination polymer with tridentate and monodentate binding of Cu^{II} ions at opposite ends of the ligand bridge neighboring units and yield a two-dimensional arrangement running in the ab plane (Fig. 2). Yigit *et al.* (2006) recently reported a coordination polymer featuring a similar head-to-tail arrangement of pyrazine-2,3,5,6-tetracarboxylic acid. There is intermolecular H-bonds ($\text{O}3\text{---H}3\cdots\text{O}4$) showed as strong H bond, which link the molecular and water to form host-guest model.

Experimental

4.00 g Potassium permanganate was to be dissolved in 30 ml pure water in a beaker and 1.0 ml 98% H_2SO_4 was added to the solution. After stirring 10 min, and then added 2,6-dimethylpyrazine to the mixture. The reaction was keeping at room temperature for 24 h. The resulting solution was filtered, and the filtrate was left in a beaker, then 2.00 g copper(II) sulfate pentahydrate was added to the filtrate. After stirring for 20 min, the copper(II) sulfate pentahydrate was solved completely. The blue solution was kept at the room temperature for two weeks and blue blocks of (I) were obtained. Yield: 86 percent, m.p. 551 K. Anal. Calc. for $\text{C}_{12}\text{H}_{14}\text{CuN}_4\text{O}_6$: C: 38.5567; H: 3.7750; N: 14.9881; Found: C: 37.67; H: 3.86; N: 14.32%. Selected IR (KBr, cm^{-1}) 3439(w), 2889 (w), 1638(s), 1596 (s), 1536(w), 1409(s), 1364(m), 1275(m), 1149(s), 1151(s), 1034(s), 941(s), 820(m), 800(s), 712(w), 531(m), 471(w).

Refinement

All H atoms were placed geometrically and treated as riding on their parent atoms with C—H 0.93(pyrazine), C—H 0.97 (methylene) Å [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$] and O—H 0.82 Å (hydroxyl) [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$].

Figures



Fig. 1. The molecular structure of (I) showing 30% probability displacement ellipsoids.




Fig. 2. The crystal packing of (I), viewed approximately along the *c* axis.

Poly[[bis(μ_2 -6-methylpyrazin-2-carboxylato- $\kappa^3N^1,O:N^4$)copper(II)] dihydrate]

Crystal data

[Cu(C₆H₅N₂O₂)₂] \cdot 2H₂O

M_r = 373.81

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

a = 8.371 (1) Å

b = 9.7901 (11) Å

c = 10.3849 (13) Å

β = 112.277 (1)°

V = 787.55 (16) Å³

Z = 2

F_{000} = 382

D_x = 1.576 Mg m⁻³

Mo $K\alpha$ radiation, λ = 0.71073 Å

Cell parameters from 1598 reflections

θ = 2.6–26.2°

μ = 1.42 mm⁻¹

T = 298 K

Block, blue

0.34 × 0.32 × 0.30 mm

Data collection

Bruker SMART CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 298 K

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2003)

T_{\min} = 0.644, T_{\max} = 0.675

3821 measured reflections

1388 independent reflections

1094 reflections with $I > 2\sigma(I)$

R_{int} = 0.027

θ_{max} = 25.0°

θ_{min} = 2.6°

h = -9→9

k = -11→11

l = -12→9

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.049$$

$$wR(F^2) = 0.160$$

$$S = 1.06$$

1388 reflections

126 parameters

Primary atom site location: structure-invariant direct methods

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0987P)^2 + 1.8186P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 1.23 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.47 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cu1	0.5000	0.5000	0.5000	0.0210 (3)	
N1	0.5565 (5)	0.3429 (4)	0.3509 (4)	0.0273 (9)	
N2	0.5246 (5)	0.1511 (4)	0.1450 (4)	0.0270 (9)	
O1	0.2704 (4)	0.4333 (3)	0.3861 (3)	0.0259 (7)	
O2	0.1014 (18)	0.343 (6)	0.184 (5)	0.049 (10)	0.46 (9)
O2'	0.1110 (16)	0.279 (5)	0.235 (4)	0.049 (9)	0.54 (9)
O3	0.8445 (18)	0.5171 (12)	0.0914 (16)	0.097 (4)	0.50
H3C	0.9307	0.4639	0.1248	0.116*	0.50
H3D	0.8631	0.5877	0.1426	0.116*	0.50
O4	0.499 (3)	0.4908 (16)	0.9398 (18)	0.126 (6)	0.50
H4D	0.5904	0.5025	1.0121	0.152*	0.50
H4E	0.4122	0.4929	0.9628	0.152*	0.50
C1	0.2469 (6)	0.3484 (6)	0.2899 (6)	0.0411 (14)	
C2	0.4033 (6)	0.2942 (5)	0.2678 (5)	0.0302 (11)	
C3	0.3863 (6)	0.1987 (6)	0.1653 (5)	0.0337 (12)	
H3	0.2771	0.1672	0.1096	0.040*	
C4	0.6768 (6)	0.1996 (5)	0.2266 (5)	0.0319 (11)	
H4A	0.7750	0.1682	0.2146	0.038*	
C5	0.6955 (6)	0.2969 (5)	0.3308 (5)	0.0301 (11)	
C6	0.8676 (7)	0.3506 (7)	0.4211 (6)	0.0496 (16)	
H6A	0.8532	0.4283	0.4721	0.074*	
H6B	0.9300	0.3775	0.3644	0.074*	

supplementary materials

H6C 0.9309 0.2809 0.4850 0.074*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0214 (5)	0.0226 (5)	0.0193 (5)	-0.0019 (3)	0.0079 (3)	0.0006 (3)
N1	0.023 (2)	0.033 (2)	0.026 (2)	-0.0023 (17)	0.0093 (16)	-0.0047 (17)
N2	0.025 (2)	0.031 (2)	0.027 (2)	-0.0020 (18)	0.0120 (16)	-0.0056 (17)
O1	0.0234 (16)	0.0301 (18)	0.0267 (17)	-0.0018 (14)	0.0124 (13)	-0.0062 (15)
O2	0.025 (5)	0.065 (19)	0.054 (13)	-0.004 (6)	0.012 (5)	-0.030 (15)
O2'	0.025 (4)	0.065 (17)	0.054 (11)	-0.004 (5)	0.012 (5)	-0.030 (12)
O3	0.074 (8)	0.092 (10)	0.115 (11)	0.018 (6)	0.026 (8)	0.013 (7)
O4	0.120 (14)	0.133 (15)	0.129 (17)	-0.002 (9)	0.050 (14)	0.000 (11)
C1	0.019 (3)	0.058 (4)	0.045 (3)	-0.004 (2)	0.010 (2)	-0.019 (3)
C2	0.025 (3)	0.033 (3)	0.032 (3)	-0.004 (2)	0.010 (2)	-0.010 (2)
C3	0.025 (3)	0.041 (3)	0.035 (3)	-0.004 (2)	0.011 (2)	-0.012 (2)
C4	0.023 (3)	0.040 (3)	0.034 (3)	0.002 (2)	0.012 (2)	-0.007 (2)
C5	0.026 (3)	0.032 (3)	0.033 (3)	-0.002 (2)	0.012 (2)	-0.006 (2)
C6	0.024 (3)	0.068 (4)	0.054 (4)	-0.001 (3)	0.011 (3)	-0.027 (3)

Geometric parameters (\AA , $^\circ$)

Cu1—O1 ⁱ	1.949 (3)	O3—H3C	0.8500
Cu1—O1	1.949 (3)	O3—H3D	0.8500
Cu1—N2 ⁱⁱ	2.064 (4)	O4—O4 ^v	1.26 (3)
Cu1—N2 ⁱⁱⁱ	2.064 (4)	O4—H4D	0.8500
Cu1—N1 ⁱ	2.354 (4)	O4—H4E	0.8500
Cu1—N1	2.354 (4)	C1—C2	1.509 (7)
N1—C2	1.333 (6)	C2—C3	1.383 (7)
N1—C5	1.336 (6)	C3—H3	0.9300
N2—C4	1.322 (6)	C4—C5	1.405 (7)
N2—C3	1.337 (6)	C4—H4A	0.9300
N2—Cu1 ^{iv}	2.065 (4)	C5—C6	1.486 (7)
O1—C1	1.256 (6)	C6—H6A	0.9600
O2—C1	1.30 (3)	C6—H6B	0.9600
O2'—C1	1.26 (2)	C6—H6C	0.9600
O1 ⁱ —Cu1—O1	180.0	H4D—O4—H4E	109.0
O1 ⁱ —Cu1—N2 ⁱⁱ	89.70 (14)	O1—C1—O2'	124.0 (8)
O1—Cu1—N2 ⁱⁱ	90.30 (14)	O1—C1—O2	121.0 (11)
O1 ⁱ —Cu1—N2 ⁱⁱⁱ	90.30 (14)	O2'—C1—O2	36.7 (6)
O1—Cu1—N2 ⁱⁱⁱ	89.70 (14)	O1—C1—C2	118.1 (4)
N2 ⁱⁱ —Cu1—N2 ⁱⁱⁱ	180.0	O2'—C1—C2	115.3 (8)
O1 ⁱ —Cu1—N1 ⁱ	77.27 (13)	O2—C1—C2	116.5 (8)
O1—Cu1—N1 ⁱ	102.73 (13)	N1—C2—C3	122.2 (5)
N2 ⁱⁱ —Cu1—N1 ⁱ	88.74 (15)	N1—C2—C1	116.9 (4)
N2 ⁱⁱⁱ —Cu1—N1 ⁱ	91.26 (15)	C3—C2—C1	120.9 (4)

O1 ⁱ —Cu1—N1	102.73 (13)	N2—C3—C2	121.0 (4)
O1—Cu1—N1	77.27 (13)	N2—C3—H3	119.5
N2 ⁱⁱ —Cu1—N1	91.26 (15)	C2—C3—H3	119.5
N2 ⁱⁱⁱ —Cu1—N1	88.74 (15)	N2—C4—C5	122.4 (4)
N1 ⁱ —Cu1—N1	180.0	N2—C4—H4A	118.8
C2—N1—C5	117.4 (4)	C5—C4—H4A	118.8
C2—N1—Cu1	106.0 (3)	N1—C5—C4	120.0 (4)
C5—N1—Cu1	136.6 (3)	N1—C5—C6	118.4 (4)
C4—N2—C3	117.1 (4)	C4—C5—C6	121.6 (4)
C4—N2—Cu1 ^{iv}	122.0 (3)	C5—C6—H6A	109.5
C3—N2—Cu1 ^{iv}	120.8 (3)	C5—C6—H6B	109.5
C1—O1—Cu1	121.7 (3)	H6A—C6—H6B	109.5
H3C—O3—H3D	108.5	C5—C6—H6C	109.5
O4 ^v —O4—H4D	55.8	H6A—C6—H6C	109.5
O4 ^v —O4—H4E	53.3	H6B—C6—H6C	109.5
O1 ⁱ —Cu1—N1—C2	-179.1 (3)	C5—N1—C2—C1	-179.1 (5)
O1—Cu1—N1—C2	0.9 (3)	Cu1—N1—C2—C1	0.0 (5)
N2 ⁱⁱ —Cu1—N1—C2	90.9 (3)	O1—C1—C2—N1	-1.4 (8)
N2 ⁱⁱⁱ —Cu1—N1—C2	-89.1 (3)	O2'—C1—C2—N1	-164 (3)
N1 ⁱ —Cu1—N1—C2	168 (100)	O2—C1—C2—N1	155 (4)
O1 ⁱ —Cu1—N1—C5	-0.4 (5)	O1—C1—C2—C3	178.9 (5)
O1—Cu1—N1—C5	179.6 (5)	O2'—C1—C2—C3	17 (3)
N2 ⁱⁱ —Cu1—N1—C5	-90.4 (5)	O2—C1—C2—C3	-24 (4)
N2 ⁱⁱⁱ —Cu1—N1—C5	89.6 (5)	C4—N2—C3—C2	-0.2 (8)
N1 ⁱ —Cu1—N1—C5	-13 (100)	Cu1 ^{iv} —N2—C3—C2	176.2 (4)
O1 ⁱ —Cu1—O1—C1	5(100)	N1—C2—C3—N2	-0.1 (8)
N2 ⁱⁱ —Cu1—O1—C1	-93.0 (4)	C1—C2—C3—N2	179.6 (5)
N2 ⁱⁱⁱ —Cu1—O1—C1	87.0 (4)	C3—N2—C4—C5	0.0 (8)
N1 ⁱ —Cu1—O1—C1	178.2 (4)	Cu1 ^{iv} —N2—C4—C5	-176.3 (4)
N1—Cu1—O1—C1	-1.8 (4)	C2—N1—C5—C4	-0.8 (7)
Cu1—O1—C1—O2'	163 (3)	Cu1—N1—C5—C4	-179.5 (4)
Cu1—O1—C1—O2	-153 (4)	C2—N1—C5—C6	179.4 (5)
Cu1—O1—C1—C2	2.4 (7)	Cu1—N1—C5—C6	0.7 (8)
C5—N1—C2—C3	0.6 (7)	N2—C4—C5—N1	0.5 (8)
Cu1—N1—C2—C3	179.7 (4)	N2—C4—C5—C6	-179.7 (5)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x, -y+1/2, z+1/2$; (iii) $-x+1, y+1/2, -z+1/2$; (iv) $-x+1, y-1/2, -z+1/2$; (v) $-x+1, -y+1, -z+2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O3—H3C \cdots O2 ^{vi}	0.85	1.78	2.624 (3)	175
O3—H3C \cdots O2' ^{vi}	0.85	2.36	3.181 (3)	163
O3—H3D \cdots O2 ⁱⁱⁱ	0.85	2.23	3.074 (2)	175

supplementary materials

O4—H4D···O3 ^{vii}	0.85	1.97	2.73 (3)	147
O4—H4E···O3 ⁱ	0.85	2.01	2.77 (2)	150

Symmetry codes: (vi) $x+1, y, z$; (iii) $-x+1, y+1/2, -z+1/2$; (vii) $x, y, z+1$; (i) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

