

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

**(E)-3-(9-Anthryl)-1-(4-fluorophenyl)-2-(1H-1,2,4-triazol-1-yl)prop-2-en-1-one**Guang-Zhou Wang,<sup>a</sup> Min Su<sup>b</sup> and Cheng-He Zhou<sup>a\*</sup>

<sup>a</sup>Laboratory of Bioorganic & Medicinal Chemistry, School of Chemistry and Chemical Engineering, Southwest University, Chongqing 400715, People's Republic of China, and <sup>b</sup>College of Pharmacy, The Third Military Medical University, Chongqing 400038, People's Republic of China

Correspondence e-mail: zhouch@swu.edu.cn


Received 26 September 2009; accepted 26 September 2009

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.064;  $wR$  factor = 0.152; data-to-parameter ratio = 15.5.

The  $\text{C}=\text{C}$  double-bond in the title compound,  $\text{C}_{25}\text{H}_{16}\text{FN}_3\text{O}$ , has an *E* configuration. The dihedral angle between the fluorophenyl and triazole rings is  $80.57(2)^\circ$ .

## Related literature

For the synthesis, see: Erhardt *et al.* (1985); Kranz *et al.* (1980). For the pharmacological activity of azoles including imidazole and triazole derivatives, see: Luo *et al.* (2009); Zhou *et al.* (2009). For related structures, see: Lu *et al.* (2009); Wang *et al.* (2009); Yan *et al.* (2009).



## Experimental

## Crystal data

 $\text{C}_{25}\text{H}_{16}\text{FN}_3\text{O}$  $M_r = 393.41$ Orthorhombic, *Pbca* $a = 13.1115(8)$  Å $b = 13.4737(8)$  Å $c = 21.7019(14)$  Å $V = 3833.9(4)$  Å<sup>3</sup> $Z = 8$ Mo  $K\alpha$  radiation $\mu = 0.09$  mm<sup>-1</sup> $T = 298$  K $0.20 \times 0.10 \times 0.10$  mm

## Data collection

Bruker SMART diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.982$ ,  $T_{\max} = 0.991$ 

16309 measured reflections

4187 independent reflections

3550 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.041$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.064$  $wR(F^2) = 0.152$  $S = 1.14$ 

4187 reflections

271 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.24$  e Å<sup>-3</sup> $\Delta\rho_{\text{min}} = -0.19$  e Å<sup>-3</sup>

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINTE-Plus* (Bruker, 2001); data reduction: *SAINTE-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We thank Southwest University (SWUB2006018, XSGX0602 and SWUF2007023) and the Natural Science Foundation of Chongqing (2007BB5369) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2652).

## References

- Bruker (2001). *SAINTE-Plus* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Erhardt, H., Mildenerger, H., Handte, R., Sachse, B., Hartz, P. & Bürstell, H. (1985). German Patent No. DE3406908.
- Kranz, E., Krämer, W., Büchel, K.-H., Brandes, W. & Forhberger, P.-E. (1980). German Patent No. DE2832233.
- Lu, Y.-H., Wang, G.-Z., Zhou, C.-H. & Zhang, Y.-Y. (2009). *Acta Cryst.* **E65**, o1396.
- Luo, Y., Lu, Y.-H., Gan, L.-L., Zhou, C.-H., Wu, J., Geng, R.-X. & Zhang, Y.-Y. (2009). *Arch. Pharm. Chem. Life Sci.* **342**, 386–393.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, G., Lu, Y., Zhou, C. & Zhang, Y. (2009). *Acta Cryst.* **E65**, o1113.
- Yan, C.-Y., Wang, G.-Z. & Zhou, C.-H. (2009). *Acta Cryst.* **E65**, o2054.
- Zhou, C.-H., Gan, L.-L., Zhang, Y.-Y., Zhang, F.-F., Wang, G.-Z., Jin, L. & Geng, R.-X. (2009). *Sci. China Ser. Chem.* **52**, 415–458.

**supplementary materials**

*Acta Cryst.* (2009). E65, o2598 [ doi:10.1107/S1600536809039178 ]

**(E)-3-(9-Anthryl)-1-(4-fluorophenyl)-2-(1H-1,2,4-triazol-1-yl)prop-2-en-1-one**

**G.-Z. Wang, M. Su and C.-H. Zhou**

**Comment**

Chalcones have been investigated for a long time due to important biological activities. We have already synthesized and reported several related structures of chalcones (Lu *et al.*, 2009; Wang *et al.*, 2009; Yan *et al.*, 2009). Azole compounds including imidazole and triazole derivatives are important types of antimicrobial drugs in clinical use, and are excellent ligands with several N-atoms which could coordinate with many kinds of metals (Luo *et al.*, 2009; Zhou *et al.*, 2009). The anthracene moiety is well known for its high absorption co-efficient as well as high fluorescence yields. These interesting properties lead us to develop the title anthryl-triazole chalcone derivatives containing the pharmacophore (triazole) and fluorophore (anthracene), and these compounds exhibit good antimicrobial, antitumor and fluorescent properties. Here we present the title compound (I) crystal structure.

The structure of title compound, C<sub>25</sub>H<sub>16</sub>FN<sub>3</sub>O, has orthorhombic (*Pbca*) symmetry. It is of interest with respect to biological activity. In the structure, the dihedral angle between the benzene and triazole ring is 80.57 (2)°. Weak intermolecular C—H···O and C—H···N interactions contribute to the crystal packing.

**Experimental**

Compound (I) was synthesized according to the procedure of Erhardt *et al.* (1985) and Kranz *et al.* (1980). Single crystals used in X-ray diffraction studies were grown in dichlormethane by slow evaporation at room temperature.

**Refinement**

All the hydrogen atoms were placed at their geometrical positions with C—H = 0.93 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

**Figures**




Fig. 1. The molecular structure of (I), showing 50% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. A partial packing diagram. Hydrogen bonds are shown as dashed lines.

## (E)-3-(9-Anthryl)-1-(4-fluorophenyl)-2-(1H-1,2,4-triazol-1-yl)prop-2-en-1-one

### Crystal data

$C_{25}H_{16}FN_3O$	$F_{000} = 1632$
$M_r = 393.41$	$D_x = 1.363 \text{ Mg m}^{-3}$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2ab	Cell parameters from 5206 reflections
$a = 13.1115 (8) \text{ \AA}$	$\theta = 2.4\text{--}27.6^\circ$
$b = 13.4737 (8) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 21.7019 (14) \text{ \AA}$	$T = 298 \text{ K}$
$V = 3833.9 (4) \text{ \AA}^3$	Block, yellow
$Z = 8$	$0.20 \times 0.10 \times 0.10 \text{ mm}$

### Data collection

Bruker SMART diffractometer	4187 independent reflections
Radiation source: fine-focus sealed tube	3550 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.041$
$T = 298 \text{ K}$	$\theta_{\text{max}} = 27.0^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\text{min}} = 2.4^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -16 \rightarrow 6$
$T_{\text{min}} = 0.982$ , $T_{\text{max}} = 0.991$	$k = -12 \rightarrow 17$
16309 measured reflections	$l = -27 \rightarrow 26$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.064$	H-atom parameters constrained
$wR(F^2) = 0.152$	$w = 1/[\sigma^2(F_o^2) + (0.0581P)^2 + 1.3538P]$
$S = 1.14$	where $P = (F_o^2 + 2F_c^2)/3$
4187 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
271 parameters	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$
	Extinction correction: none

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.33997 (13)	-0.03462 (14)	0.19230 (8)	0.0395 (4)
C2	0.28391 (14)	-0.00708 (17)	0.14089 (9)	0.0471 (5)
H2	0.2586	0.0572	0.1377	0.057*
C3	0.26557 (17)	-0.0751 (2)	0.09435 (10)	0.0613 (6)
H3	0.2296	-0.0569	0.0592	0.074*
C4	0.30156 (18)	-0.1692 (2)	0.10137 (12)	0.0676 (7)
C5	0.35496 (19)	-0.19944 (19)	0.15162 (13)	0.0710 (7)
H5	0.3771	-0.2648	0.1551	0.085*
C6	0.37565 (16)	-0.13104 (16)	0.19739 (11)	0.0544 (5)
H6	0.4136	-0.1498	0.2317	0.065*
C7	0.35581 (14)	0.03737 (14)	0.24300 (8)	0.0385 (4)
C8	0.45641 (13)	0.03523 (13)	0.27672 (8)	0.0360 (4)
C9	0.37480 (17)	0.02786 (19)	0.38130 (9)	0.0587 (6)
H9	0.3100	0.0069	0.3697	0.070*
C10	0.50043 (18)	0.07070 (17)	0.43119 (9)	0.0542 (5)
H10	0.5415	0.0867	0.4647	0.065*
C11	0.54642 (14)	0.03428 (14)	0.24860 (8)	0.0394 (4)
H11	0.6044	0.0310	0.2732	0.047*
C12	0.56121 (13)	0.03800 (14)	0.18104 (8)	0.0376 (4)
C13	0.53007 (13)	0.12230 (14)	0.14735 (8)	0.0392 (4)
C14	0.48685 (16)	0.20863 (15)	0.17516 (9)	0.0475 (5)
H14	0.4783	0.2104	0.2177	0.057*
C15	0.45807 (18)	0.28783 (18)	0.14105 (11)	0.0589 (6)
H15	0.4301	0.3430	0.1605	0.071*
C16	0.46980 (19)	0.28793 (19)	0.07665 (11)	0.0634 (6)
H16	0.4485	0.3424	0.0537	0.076*
C17	0.51180 (17)	0.20924 (18)	0.04815 (10)	0.0576 (6)
H17	0.5198	0.2105	0.0056	0.069*
C18	0.54443 (14)	0.12368 (16)	0.08176 (8)	0.0437 (5)
C19	0.58943 (16)	0.04302 (16)	0.05307 (9)	0.0488 (5)
H19	0.5968	0.0439	0.0105	0.059*
C20	0.62388 (14)	-0.03901 (15)	0.08563 (9)	0.0437 (5)
C21	0.67370 (18)	-0.12021 (18)	0.05638 (10)	0.0583 (6)
H21	0.6821	-0.1195	0.0138	0.070*
C22	0.7087 (2)	-0.19767 (19)	0.08895 (12)	0.0696 (7)
H22	0.7412	-0.2498	0.0689	0.083*
C23	0.6965 (2)	-0.20059 (18)	0.15352 (12)	0.0667 (7)
H23	0.7214	-0.2545	0.1757	0.080*

## supplementary materials

---

C24	0.64870 (17)	-0.12541 (16)	0.18361 (10)	0.0532 (5)
H24	0.6408	-0.1288	0.2261	0.064*
C25	0.61067 (14)	-0.04183 (14)	0.15128 (8)	0.0405 (4)
F1	0.28332 (14)	-0.23601 (15)	0.05633 (9)	0.1089 (7)
N1	0.45059 (11)	0.04595 (11)	0.34187 (7)	0.0386 (4)
N2	0.40276 (16)	0.04324 (16)	0.43811 (8)	0.0624 (5)
N3	0.53431 (13)	0.07357 (14)	0.37481 (7)	0.0507 (4)
O1	0.29069 (10)	0.09647 (12)	0.25734 (7)	0.0580 (4)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0319 (9)	0.0475 (11)	0.0392 (9)	-0.0033 (8)	0.0027 (7)	-0.0032 (8)
C2	0.0384 (10)	0.0588 (12)	0.0442 (11)	-0.0026 (9)	-0.0005 (8)	-0.0006 (9)
C3	0.0460 (11)	0.0943 (19)	0.0437 (11)	-0.0103 (12)	-0.0051 (9)	-0.0090 (12)
C4	0.0497 (12)	0.0877 (19)	0.0653 (15)	-0.0100 (13)	0.0023 (11)	-0.0383 (14)
C5	0.0600 (14)	0.0563 (14)	0.097 (2)	0.0019 (12)	-0.0040 (14)	-0.0270 (14)
C6	0.0499 (12)	0.0502 (12)	0.0632 (13)	0.0028 (10)	-0.0088 (10)	-0.0064 (10)
C7	0.0342 (9)	0.0441 (10)	0.0372 (9)	-0.0024 (8)	0.0023 (7)	-0.0008 (8)
C8	0.0386 (9)	0.0398 (10)	0.0297 (8)	-0.0014 (7)	-0.0009 (7)	-0.0012 (7)
C9	0.0463 (11)	0.0879 (17)	0.0418 (11)	-0.0146 (11)	0.0088 (9)	0.0002 (11)
C10	0.0641 (13)	0.0643 (14)	0.0343 (10)	-0.0054 (11)	-0.0020 (9)	-0.0053 (9)
C11	0.0350 (9)	0.0499 (11)	0.0334 (9)	-0.0006 (8)	-0.0030 (7)	0.0003 (8)
C12	0.0307 (8)	0.0506 (11)	0.0315 (9)	-0.0065 (8)	0.0011 (7)	-0.0026 (8)
C13	0.0324 (8)	0.0494 (11)	0.0358 (9)	-0.0089 (8)	-0.0008 (7)	-0.0002 (8)
C14	0.0499 (11)	0.0516 (12)	0.0411 (10)	-0.0046 (9)	0.0025 (9)	-0.0013 (9)
C15	0.0624 (14)	0.0503 (13)	0.0641 (14)	0.0032 (10)	0.0060 (11)	0.0028 (11)
C16	0.0622 (14)	0.0642 (15)	0.0637 (14)	0.0088 (12)	0.0056 (11)	0.0229 (12)
C17	0.0559 (13)	0.0763 (16)	0.0405 (11)	-0.0025 (12)	0.0027 (9)	0.0155 (11)
C18	0.0378 (9)	0.0573 (12)	0.0360 (10)	-0.0068 (9)	0.0002 (8)	0.0032 (9)
C19	0.0485 (11)	0.0691 (14)	0.0288 (9)	-0.0086 (10)	0.0025 (8)	-0.0007 (9)
C20	0.0389 (10)	0.0542 (12)	0.0381 (10)	-0.0091 (8)	0.0022 (8)	-0.0093 (9)
C21	0.0662 (14)	0.0628 (15)	0.0460 (12)	-0.0052 (12)	0.0095 (10)	-0.0147 (11)
C22	0.0804 (17)	0.0574 (15)	0.0710 (16)	0.0056 (13)	0.0183 (13)	-0.0151 (13)
C23	0.0755 (16)	0.0506 (13)	0.0741 (16)	0.0068 (12)	0.0109 (13)	0.0064 (12)
C24	0.0554 (12)	0.0562 (13)	0.0481 (12)	0.0005 (10)	0.0085 (10)	0.0039 (10)
C25	0.0355 (9)	0.0472 (11)	0.0389 (10)	-0.0084 (8)	0.0027 (7)	-0.0014 (8)
F1	0.0917 (12)	0.1308 (15)	0.1043 (13)	-0.0117 (11)	-0.0058 (10)	-0.0780 (12)
N1	0.0392 (8)	0.0445 (9)	0.0321 (8)	-0.0036 (7)	0.0021 (6)	-0.0009 (6)
N2	0.0652 (12)	0.0856 (14)	0.0365 (9)	-0.0080 (11)	0.0098 (8)	0.0014 (9)
N3	0.0491 (9)	0.0681 (12)	0.0350 (8)	-0.0111 (8)	-0.0023 (7)	-0.0072 (8)
O1	0.0419 (8)	0.0665 (10)	0.0657 (10)	0.0116 (7)	-0.0041 (7)	-0.0224 (8)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C1—C6	1.385 (3)	C12—C25	1.412 (3)
C1—C2	1.387 (3)	C13—C14	1.428 (3)
C1—C7	1.481 (3)	C13—C18	1.436 (3)
C2—C3	1.385 (3)	C14—C15	1.352 (3)

C2—H2	0.9300	C14—H14	0.9300
C3—C4	1.362 (4)	C15—C16	1.406 (3)
C3—H3	0.9300	C15—H15	0.9300
C4—F1	1.350 (3)	C16—C17	1.345 (3)
C4—C5	1.358 (4)	C16—H16	0.9300
C5—C6	1.382 (3)	C17—C18	1.430 (3)
C5—H5	0.9300	C17—H17	0.9300
C6—H6	0.9300	C18—C19	1.385 (3)
C7—O1	1.208 (2)	C19—C20	1.387 (3)
C7—C8	1.509 (2)	C19—H19	0.9300
C8—C11	1.329 (2)	C20—C21	1.424 (3)
C8—N1	1.423 (2)	C20—C25	1.436 (3)
C9—N2	1.303 (3)	C21—C22	1.342 (3)
C9—N1	1.334 (2)	C21—H21	0.9300
C9—H9	0.9300	C22—C23	1.411 (4)
C10—N3	1.302 (3)	C22—H22	0.9300
C10—N2	1.341 (3)	C23—C24	1.358 (3)
C10—H10	0.9300	C23—H23	0.9300
C11—C12	1.480 (2)	C24—C25	1.417 (3)
C11—H11	0.9300	C24—H24	0.9300
C12—C13	1.411 (3)	N1—N3	1.362 (2)
C6—C1—C2	119.61 (18)	C15—C14—H14	119.3
C6—C1—C7	120.50 (17)	C13—C14—H14	119.3
C2—C1—C7	119.78 (18)	C14—C15—C16	120.9 (2)
C3—C2—C1	120.1 (2)	C14—C15—H15	119.5
C3—C2—H2	119.9	C16—C15—H15	119.5
C1—C2—H2	119.9	C17—C16—C15	120.1 (2)
C4—C3—C2	118.3 (2)	C17—C16—H16	120.0
C4—C3—H3	120.8	C15—C16—H16	120.0
C2—C3—H3	120.8	C16—C17—C18	121.6 (2)
F1—C4—C5	118.2 (3)	C16—C17—H17	119.2
F1—C4—C3	118.6 (3)	C18—C17—H17	119.2
C5—C4—C3	123.2 (2)	C19—C18—C17	122.07 (18)
C4—C5—C6	118.6 (2)	C19—C18—C13	119.44 (18)
C4—C5—H5	120.7	C17—C18—C13	118.49 (19)
C6—C5—H5	120.7	C18—C19—C20	122.34 (17)
C5—C6—C1	120.1 (2)	C18—C19—H19	118.8
C5—C6—H6	119.9	C20—C19—H19	118.8
C1—C6—H6	119.9	C19—C20—C21	122.31 (19)
O1—C7—C1	121.59 (17)	C19—C20—C25	119.16 (18)
O1—C7—C8	120.36 (17)	C21—C20—C25	118.53 (19)
C1—C7—C8	118.06 (16)	C22—C21—C20	121.3 (2)
C11—C8—N1	120.33 (16)	C22—C21—H21	119.3
C11—C8—C7	123.64 (16)	C20—C21—H21	119.3
N1—C8—C7	115.66 (15)	C21—C22—C23	120.4 (2)
N2—C9—N1	111.61 (19)	C21—C22—H22	119.8
N2—C9—H9	124.2	C23—C22—H22	119.8
N1—C9—H9	124.2	C24—C23—C22	120.6 (2)
N3—C10—N2	116.04 (19)	C24—C23—H23	119.7

## supplementary materials

---

N3—C10—H10	122.0	C22—C23—H23	119.7
N2—C10—H10	122.0	C23—C24—C25	121.1 (2)
C8—C11—C12	124.83 (16)	C23—C24—H24	119.4
C8—C11—H11	117.6	C25—C24—H24	119.4
C12—C11—H11	117.6	C12—C25—C24	122.70 (17)
C13—C12—C25	120.59 (16)	C12—C25—C20	119.28 (17)
C13—C12—C11	120.18 (17)	C24—C25—C20	118.02 (18)
C25—C12—C11	119.17 (17)	C9—N1—N3	108.28 (15)
C12—C13—C14	123.48 (17)	C9—N1—C8	131.20 (16)
C12—C13—C18	119.09 (17)	N3—N1—C8	120.40 (14)
C14—C13—C18	117.41 (17)	C9—N2—C10	101.93 (17)
C15—C14—C13	121.49 (19)	C10—N3—N1	102.13 (16)
C6—C1—C2—C3	-1.3 (3)	C14—C13—C18—C19	177.72 (18)
C7—C1—C2—C3	-177.61 (18)	C12—C13—C18—C17	179.52 (17)
C1—C2—C3—C4	1.7 (3)	C14—C13—C18—C17	-2.0 (3)
C2—C3—C4—F1	179.6 (2)	C17—C18—C19—C20	178.12 (19)
C2—C3—C4—C5	-0.3 (4)	C13—C18—C19—C20	-1.6 (3)
F1—C4—C5—C6	178.8 (2)	C18—C19—C20—C21	-177.50 (19)
C3—C4—C5—C6	-1.3 (4)	C18—C19—C20—C25	1.4 (3)
C4—C5—C6—C1	1.6 (4)	C19—C20—C21—C22	178.3 (2)
C2—C1—C6—C5	-0.4 (3)	C25—C20—C21—C22	-0.6 (3)
C7—C1—C6—C5	175.9 (2)	C20—C21—C22—C23	0.2 (4)
C6—C1—C7—O1	-141.0 (2)	C21—C22—C23—C24	0.3 (4)
C2—C1—C7—O1	35.2 (3)	C22—C23—C24—C25	-0.5 (4)
C6—C1—C7—C8	39.1 (2)	C13—C12—C25—C24	176.20 (17)
C2—C1—C7—C8	-144.71 (18)	C11—C12—C25—C24	-1.1 (3)
O1—C7—C8—C11	-131.1 (2)	C13—C12—C25—C20	-3.5 (3)
C1—C7—C8—C11	48.8 (3)	C11—C12—C25—C20	179.16 (16)
O1—C7—C8—N1	41.8 (3)	C23—C24—C25—C12	-179.6 (2)
C1—C7—C8—N1	-138.21 (17)	C23—C24—C25—C20	0.2 (3)
N1—C8—C11—C12	-170.82 (17)	C19—C20—C25—C12	1.2 (3)
C7—C8—C11—C12	1.8 (3)	C21—C20—C25—C12	-179.87 (18)
C8—C11—C12—C13	63.7 (3)	C19—C20—C25—C24	-178.53 (18)
C8—C11—C12—C25	-118.9 (2)	C21—C20—C25—C24	0.4 (3)
C25—C12—C13—C14	-175.05 (17)	N2—C9—N1—N3	1.0 (3)
C11—C12—C13—C14	2.2 (3)	N2—C9—N1—C8	176.88 (19)
C25—C12—C13—C18	3.3 (3)	C11—C8—N1—C9	-163.3 (2)
C11—C12—C13—C18	-179.43 (16)	C7—C8—N1—C9	23.5 (3)
C12—C13—C14—C15	179.98 (19)	C11—C8—N1—N3	12.2 (3)
C18—C13—C14—C15	1.6 (3)	C7—C8—N1—N3	-160.99 (17)
C13—C14—C15—C16	0.0 (3)	N1—C9—N2—C10	-0.7 (3)
C14—C15—C16—C17	-1.2 (4)	N3—C10—N2—C9	0.2 (3)
C15—C16—C17—C18	0.7 (4)	N2—C10—N3—N1	0.4 (3)
C16—C17—C18—C19	-178.8 (2)	C9—N1—N3—C10	-0.8 (2)
C16—C17—C18—C13	0.9 (3)	C8—N1—N3—C10	-177.23 (17)
C12—C13—C18—C19	-0.7 (3)		

*Hydrogen-bond geometry* (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C23—H23···O1 <sup>i</sup>	0.93	2.48	3.354 (3)	156
C5—H5···N3 <sup>i</sup>	0.93	2.55	3.434 (3)	158

Symmetry codes: (i)  $-x+1, y-1/2, -z+1/2$ .

Fig. 1




Fig. 2

