

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dibenzoato- $\kappa O; \kappa^2 O, O'$ -(6,6'-dimethyl-2,2'-bipyridine- $\kappa^2 N, N'$)zinc(II)–benzoic acid (1/1)

Li Yao^{a*} and Wen-Juan Li^b

^aSchool of Computer and Information Engineering, Henan University, Kaifeng 475001, Henan, People's Republic of China, and ^bDepartment of Civil and Environmental Engineering, East China Institute of Technology, 56 Xuefu Road, Fuzhou 344000, Jiangxi, People's Republic of China
Correspondence e-mail: yaolihenu@163.com


Received 3 October 2009; accepted 14 October 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.009$ Å; R factor = 0.065; wR factor = 0.136; data-to-parameter ratio = 14.3.

In the crystal structure of the title compound, $[Zn(C_6H_5COO)_2(C_{12}H_{12}N_2)] \cdot C_6H_5COOH$, the Zn atom is pentacoordinated in distorted square-pyramidal geometry by two O atoms of a benzoate anion and two N atoms of a 6,6'-dimethyl-2,2'-bipyridine ligand occupying the basal plane and an O atom of another benzoate anion located at the apical site. In the crystal structure, intermolecular O—H \cdots O and C—H \cdots O hydrogen bonds and C—H $\cdots\pi$ interactions are present.

Related literature

For related structures, see: Alizadeh *et al.* (2009); Cui *et al.* (2005); Hökelek *et al.* (2009a,b); Klausmeyer *et al.* (2007); Phatchimkun *et al.* (2009); Zhang *et al.* (2009).



Experimental

Crystal data

$[Zn(C_6H_5O_2)_2(C_{12}H_{12}N_2)] \cdot C_7H_6O_2$
 $M_r = 613.96$
 Triclinic, $P\bar{1}$
 $a = 9.974$ (6) Å
 $b = 12.546$ (7) Å
 $c = 12.798$ (8) Å
 $\alpha = 96.631$ (11)°
 $\beta = 97.016$ (12)°
 $\gamma = 105.060$ (12)°
 $V = 1516.7$ (16) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.86$ mm⁻¹
 $T = 296$ K
 $0.16 \times 0.14 \times 0.12$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{min} = 0.875$, $T_{max} = 0.904$
 8123 measured reflections
 5295 independent reflections
 2785 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.136$
 $S = 1.04$
 5295 reflections
 370 parameters
 14 restraints
 H-atom parameters constrained
 $\Delta\rho_{max} = 0.64$ e Å⁻³
 $\Delta\rho_{min} = -0.64$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C24—H24 \cdots O5 ⁱ	0.93	2.44	3.340 (8)	162
C33—H33C \cdots O1	0.96	2.52	3.199 (8)	128
C22—H22A \cdots O2	0.96	2.57	3.289 (7)	132
O6—H6 \cdots O4 ⁱⁱ	0.82	1.85	2.655 (5)	166
C31—H31 \cdots Cg1 ⁱⁱⁱ	0.93	2.79	3.708 (7)	170

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $x, y, z + 1$; (iii) $-x + 1, -y + 1, -z$. Cg1 is the centroid of the C9–C14 ring.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2173).

References

- Alizadeh, R., Kalateh, K., Ebadi, A., Ahmadi, R. & Amani, V. (2009). *Acta Cryst.* **E65**, m1250.
 Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cui, G. H., Li, J. R., Gao, D. & Ng, S. W. (2005). *Acta Cryst.* **E61**, m72–m73.
 Hökelek, T., Dal, H., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009a). *Acta Cryst.* **E65**, m1037–m1038.
 Hökelek, T., Dal, H., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009b). *Acta Cryst.* **E65**, m1051–m1052.
 Klausmeyer, K. K., Hung-Low, F. & Renz, A. (2007). *Acta Cryst.* **E63**, m2181.
 Phatchimkun, J., Kongsaree, P., Suchaichit, N. & Chaichit, N. (2009). *Acta Cryst.* **E65**, m1020–m1021.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Zhang, B.-Y., Nie, J.-J. & Xu, D.-J. (2009). *Acta Cryst.* **E65**, m880.

supplementary materials

Acta Cryst. (2009). E65, m1408 [doi:10.1107/S1600536809042093]

Dibenzoato- κO ; $\kappa^2 O, O'$ -(6,6'-dimethyl-2,2'-bipyridine- $\kappa^2 N, N'$)zinc(II)-benzoic acid (1/1)

L. Yao and W. J. Li

Comment

As a contribution to structural characterization of 6,6'-dimethyl-2,2'-bipyridine complexes [Alizadeh *et al.*, (2009); Cui *et al.*, (2005); Hökelek *et al.*, (2009a,b); Klausmeyer *et al.*, (2007); Phatchimkun *et al.* (2009); Zhang *et al.*, (2009)], we present here the molecular structure of the title complex, $ZnL(C_6H_5COO)_2C_6H_5COOH$, where L is 6,6'-dimethyl-2,2'-bipyridine.

The title compound, contains two benzoate anions, one 6,6'-dimethyl-2,2'-bipyridine ligand and one benzoic acid molecules. One of the benzoate anions acts as a bidentate ligand, while the other is monodentate.

The molecular structure of the title complex is shown on Fig. 1. In the crystal structure, the face-to-face separation of 3.783 (4)Å suggests no $\pi \cdots \pi$ stacking between parallel bipyridine ring systems, intermolecular O—H \cdots O and C—H \cdots O hydrogen bonds (Table 1) link the molecules into a one dimensional structure, in which they may be effective in the stabilization of the structure. One weak C—H \cdots π interactions (Table 1) are also found. Cg1 is the centroid of the C9ⁱⁱⁱ-C14ⁱⁱⁱ ring. Symmetry code: (iii) 1-x, 1-y, -z.

Experimental

The title compound was synthesized hydrothermally in a teflon-lined autoclave (25 ml) by heating a mixture of 6,6'-dimethyl-2,2'-bipyridine (0.2 mmol), benzoic acid (0.4 mmol) and $ZnSO_4 \cdot H_2O$ (0.2 mmol) in water (10 ml) at 393 K for 3 d. Crystals suitable for X-ray analysis were obtained.

Refinement

The carboxy H atom was located in a difference Fourier map and refined with a O—H distance of 0.82Å [$U_{iso}(H) = 1.5U_{eq}(O)$]. H atoms bonded to C atoms were positioned geometrically and refined using a riding model with C—H = 0.93Å and 0.96Å for aromatic and methyl H atoms, respectively, and $U_{iso}(H)$ values were calculated at 1.5 $U_{eq}(C)$ for methyl groups and 1.2 $U_{eq}(C)$ for aromatic.

Figures




Fig. 1. The molecular structure of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are presented as small spheres of arbitrary radius.

Dibenzoato- $\kappa O; \kappa^2 O, O'$ -(6,6'-dimethyl-2,2'-bipyridine- $\kappa^2 N, N'$)zinc(II)-benzoic acid (1/1)

Crystal data

$[\text{Zn}(\text{C}_7\text{H}_5\text{O}_2)_2(\text{C}_{12}\text{H}_{12}\text{N}_2)] \cdot \text{C}_7\text{H}_6\text{O}_2$	$Z = 2$
$M_r = 613.96$	$F_{000} = 636$
Triclinic, $P\bar{1}$	$D_x = 1.344 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.974 (6) \text{ \AA}$	Cell parameters from 1683 reflections
$b = 12.546 (7) \text{ \AA}$	$\theta = 2.5\text{--}27.8^\circ$
$c = 12.798 (8) \text{ \AA}$	$\mu = 0.86 \text{ mm}^{-1}$
$\alpha = 96.631 (11)^\circ$	$T = 296 \text{ K}$
$\beta = 97.016 (12)^\circ$	Block, colourless
$\gamma = 105.060 (12)^\circ$	$0.16 \times 0.14 \times 0.12 \text{ mm}$
$V = 1516.7 (16) \text{ \AA}^3$	

Data collection

Bruker APEXII CCD area-detector diffractometer	5295 independent reflections
Radiation source: fine-focus sealed tube	2785 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.046$
$T = 296 \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ - and ω -scans	$\theta_{\text{min}} = 1.6^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -10 \rightarrow 11$
$T_{\text{min}} = 0.875$, $T_{\text{max}} = 0.904$	$k = -14 \rightarrow 13$
8123 measured reflections	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.065$	H-atom parameters constrained
$wR(F^2) = 0.136$	$w = 1/[\sigma^2(F_o^2) + (0.0455P)^2]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
5295 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
370 parameters	$\Delta\rho_{\text{max}} = 0.64 \text{ e \AA}^{-3}$
14 restraints	$\Delta\rho_{\text{min}} = -0.64 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.41555 (6)	0.22239 (5)	0.15235 (4)	0.0561 (2)
N1	0.2722 (4)	0.0734 (3)	0.0835 (3)	0.0509 (10)
N2	0.4792 (4)	0.2028 (3)	0.0057 (3)	0.0461 (10)
O1	0.6236 (4)	0.2621 (4)	0.2360 (3)	0.1060 (12)
O2	0.4751 (4)	0.1472 (3)	0.2994 (3)	0.0945 (13)
O3	0.3506 (4)	0.3454 (3)	0.2070 (3)	0.0708 (10)
O4	0.2026 (4)	0.3185 (3)	0.0566 (2)	0.0672 (10)
O5	0.1962 (4)	0.1688 (3)	0.8042 (3)	0.0760 (11)
O6	0.2806 (4)	0.3497 (3)	0.8688 (3)	0.0717 (10)
H6	0.2520	0.3287	0.9225	0.108*
C1	0.5999 (8)	0.1987 (7)	0.3044 (5)	0.1025 (12)
C2	0.7122 (5)	0.1859 (4)	0.3773 (3)	0.0997 (13)
C3	0.8523 (6)	0.2392 (4)	0.3750 (4)	0.128 (3)
H3	0.8755	0.2882	0.3265	0.153*
C4	0.9579 (4)	0.2192 (5)	0.4450 (5)	0.164 (4)
H4	1.0517	0.2549	0.4434	0.196*
C5	0.9232 (6)	0.1460 (5)	0.5174 (4)	0.172 (5)
H5	0.9938	0.1326	0.5642	0.206*
C6	0.7831 (7)	0.0926 (4)	0.5197 (4)	0.153 (4)
H6A	0.7599	0.0436	0.5681	0.183*
C7	0.6775 (5)	0.1126 (4)	0.4497 (4)	0.111 (3)
H7	0.5837	0.0769	0.4513	0.134*
C8	0.2506 (6)	0.3676 (4)	0.1480 (4)	0.0535 (13)
C9	0.1893 (5)	0.4536 (4)	0.1966 (4)	0.0537 (13)
C10	0.2524 (7)	0.5206 (5)	0.2920 (4)	0.0894 (19)
H10	0.3346	0.5114	0.3277	0.107*
C11	0.1952 (9)	0.6015 (6)	0.3357 (5)	0.116 (2)
H11	0.2383	0.6463	0.4003	0.139*
C12	0.0740 (9)	0.6148 (6)	0.2822 (7)	0.125 (3)
H12	0.0324	0.6668	0.3120	0.150*
C13	0.0151 (7)	0.5522 (6)	0.1864 (7)	0.117 (3)
H13	-0.0654	0.5631	0.1496	0.140*
C14	0.0726 (6)	0.4730 (4)	0.1430 (5)	0.0780 (17)

supplementary materials

H14	0.0320	0.4318	0.0763	0.094*
C15	0.2525 (5)	0.2627 (5)	0.7920 (4)	0.0529 (13)
C16	0.2973 (5)	0.2925 (5)	0.6902 (4)	0.0538 (13)
C17	0.3171 (6)	0.3999 (5)	0.6671 (4)	0.0758 (16)
H17	0.3070	0.4564	0.7167	0.091*
C18	0.3528 (7)	0.4224 (6)	0.5677 (5)	0.097 (2)
H18	0.3660	0.4941	0.5508	0.116*
C19	0.3684 (7)	0.3383 (7)	0.4953 (5)	0.098 (2)
H19	0.3910	0.3533	0.4291	0.118*
C20	0.3510 (6)	0.2331 (6)	0.5199 (5)	0.0866 (19)
H20	0.3653	0.1771	0.4720	0.104*
C21	0.3122 (5)	0.2107 (5)	0.6159 (4)	0.0689 (15)
H21	0.2955	0.1380	0.6308	0.083*
C22	0.1317 (6)	0.0689 (5)	0.2271 (4)	0.0847 (18)
H22A	0.2067	0.0747	0.2840	0.127*
H22B	0.0452	0.0241	0.2431	0.127*
H22C	0.1232	0.1421	0.2192	0.127*
C23	0.1631 (6)	0.0156 (5)	0.1256 (4)	0.0633 (15)
C24	0.0820 (6)	-0.0880 (5)	0.0732 (6)	0.0798 (18)
H24	0.0049	-0.1267	0.1009	0.096*
C25	0.1145 (7)	-0.1335 (5)	-0.0189 (6)	0.087 (2)
H25	0.0622	-0.2045	-0.0524	0.105*
C26	0.2243 (6)	-0.0747 (4)	-0.0621 (4)	0.0719 (16)
H26	0.2461	-0.1040	-0.1259	0.086*
C27	0.3021 (5)	0.0296 (4)	-0.0087 (4)	0.0482 (12)
C28	0.4195 (5)	0.1021 (4)	-0.0519 (4)	0.0471 (12)
C29	0.4627 (6)	0.0693 (5)	-0.1451 (4)	0.0648 (15)
H29	0.4218	-0.0020	-0.1832	0.078*
C30	0.5682 (7)	0.1441 (6)	-0.1813 (4)	0.0785 (17)
H30	0.5997	0.1236	-0.2440	0.094*
C31	0.6250 (6)	0.2471 (5)	-0.1248 (5)	0.0753 (16)
H31	0.6951	0.2982	-0.1492	0.090*
C32	0.5798 (6)	0.2774 (4)	-0.0311 (4)	0.0601 (14)
C33	0.6386 (6)	0.3902 (4)	0.0338 (5)	0.095 (2)
H33A	0.5630	0.4200	0.0507	0.142*
H33B	0.6962	0.4388	-0.0060	0.142*
H33C	0.6944	0.3846	0.0985	0.142*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0556 (4)	0.0578 (4)	0.0511 (4)	0.0185 (3)	-0.0002 (3)	-0.0035 (3)
N1	0.040 (2)	0.057 (3)	0.053 (3)	0.013 (2)	-0.002 (2)	0.013 (2)
N2	0.038 (2)	0.046 (3)	0.054 (2)	0.016 (2)	0.005 (2)	0.002 (2)
O1	0.0884 (18)	0.162 (3)	0.069 (2)	0.063 (2)	-0.0244 (18)	-0.003 (2)
O2	0.093 (3)	0.124 (3)	0.066 (2)	0.061 (3)	-0.034 (2)	-0.0054 (18)
O3	0.081 (3)	0.074 (2)	0.056 (2)	0.037 (2)	-0.008 (2)	-0.0079 (18)
O4	0.078 (3)	0.075 (2)	0.044 (2)	0.0216 (19)	0.0045 (19)	-0.0047 (18)

O5	0.092 (3)	0.073 (3)	0.065 (2)	0.021 (2)	0.019 (2)	0.014 (2)
O6	0.100 (3)	0.068 (2)	0.051 (2)	0.030 (2)	0.016 (2)	0.0073 (19)
C1	0.086 (2)	0.158 (3)	0.066 (2)	0.065 (2)	-0.0219 (19)	-0.006 (2)
C2	0.084 (2)	0.155 (4)	0.064 (2)	0.066 (2)	-0.021 (2)	-0.007 (2)
C3	0.075 (5)	0.201 (8)	0.097 (5)	0.050 (5)	0.003 (4)	-0.027 (5)
C4	0.096 (6)	0.256 (12)	0.121 (7)	0.078 (7)	-0.032 (6)	-0.055 (7)
C5	0.135 (9)	0.205 (11)	0.155 (9)	0.101 (8)	-0.086 (7)	-0.069 (8)
C6	0.211 (10)	0.094 (6)	0.134 (7)	0.068 (6)	-0.083 (7)	-0.003 (5)
C7	0.117 (6)	0.095 (5)	0.100 (5)	0.043 (4)	-0.057 (5)	-0.030 (4)
C8	0.068 (4)	0.049 (3)	0.042 (3)	0.014 (3)	0.014 (3)	0.003 (3)
C9	0.059 (3)	0.053 (3)	0.047 (3)	0.019 (3)	0.006 (3)	-0.002 (3)
C10	0.120 (5)	0.095 (5)	0.063 (4)	0.058 (4)	0.004 (4)	-0.005 (3)
C11	0.160 (8)	0.120 (6)	0.075 (5)	0.071 (6)	0.012 (5)	-0.024 (4)
C12	0.114 (7)	0.120 (6)	0.147 (7)	0.064 (5)	0.025 (6)	-0.030 (6)
C13	0.071 (5)	0.101 (6)	0.166 (8)	0.039 (4)	-0.012 (5)	-0.033 (5)
C14	0.060 (4)	0.068 (4)	0.096 (4)	0.021 (3)	-0.003 (4)	-0.016 (3)
C15	0.048 (3)	0.059 (4)	0.055 (3)	0.022 (3)	0.005 (3)	0.011 (3)
C16	0.049 (3)	0.066 (4)	0.045 (3)	0.016 (3)	0.002 (3)	0.009 (3)
C17	0.075 (4)	0.082 (5)	0.065 (4)	0.012 (3)	0.011 (3)	0.013 (3)
C18	0.103 (5)	0.101 (5)	0.074 (4)	-0.001 (4)	0.018 (4)	0.028 (4)
C19	0.090 (5)	0.141 (7)	0.055 (4)	0.014 (5)	0.017 (4)	0.018 (5)
C20	0.084 (5)	0.117 (6)	0.060 (4)	0.031 (4)	0.015 (3)	0.002 (4)
C21	0.072 (4)	0.082 (4)	0.054 (3)	0.026 (3)	0.009 (3)	0.006 (3)
C22	0.063 (4)	0.116 (5)	0.078 (4)	0.021 (4)	0.014 (3)	0.026 (4)
C23	0.052 (4)	0.069 (4)	0.069 (4)	0.018 (3)	-0.006 (3)	0.025 (3)
C24	0.058 (4)	0.065 (5)	0.110 (5)	0.003 (3)	0.001 (4)	0.034 (4)
C25	0.075 (5)	0.056 (4)	0.114 (6)	0.004 (4)	-0.017 (4)	0.008 (4)
C26	0.069 (4)	0.054 (4)	0.079 (4)	0.011 (3)	-0.012 (3)	-0.004 (3)
C27	0.046 (3)	0.043 (3)	0.051 (3)	0.015 (3)	-0.012 (3)	0.002 (3)
C28	0.047 (3)	0.046 (3)	0.051 (3)	0.025 (3)	-0.003 (3)	0.000 (3)
C29	0.076 (4)	0.061 (4)	0.060 (4)	0.032 (3)	0.005 (3)	-0.008 (3)
C30	0.096 (5)	0.104 (5)	0.055 (4)	0.053 (4)	0.030 (4)	0.016 (4)
C31	0.073 (4)	0.086 (5)	0.074 (4)	0.030 (4)	0.019 (3)	0.017 (4)
C32	0.060 (4)	0.056 (4)	0.064 (4)	0.019 (3)	0.005 (3)	0.009 (3)
C33	0.099 (5)	0.066 (4)	0.099 (4)	-0.012 (3)	0.023 (4)	0.005 (4)

Geometric parameters (Å, °)

Zn1—O3	1.917 (3)	C13—H13	0.9300
Zn1—N1	2.059 (4)	C14—H14	0.9300
Zn1—N2	2.062 (4)	C15—C16	1.484 (7)
Zn1—O1	2.120 (4)	C16—C21	1.368 (6)
Zn1—O2	2.283 (4)	C16—C17	1.381 (7)
N1—C27	1.344 (6)	C17—C18	1.402 (7)
N1—C23	1.350 (6)	C17—H17	0.9300
N2—C28	1.335 (5)	C18—C19	1.375 (8)
N2—C32	1.353 (6)	C18—H18	0.9300
O1—C1	1.252 (8)	C19—C20	1.365 (8)
O2—C1	1.235 (8)	C19—H19	0.9300

supplementary materials

O3—C8	1.285 (5)	C20—C21	1.369 (7)
O4—C8	1.233 (5)	C20—H20	0.9300
O5—C15	1.204 (5)	C21—H21	0.9300
O6—C15	1.327 (5)	C22—C23	1.496 (7)
O6—H6	0.8200	C22—H22A	0.9600
C1—C2	1.423 (7)	C22—H22B	0.9600
C2—C3	1.3900	C22—H22C	0.9600
C2—C7	1.3900	C23—C24	1.384 (7)
C3—C4	1.3900	C24—C25	1.362 (8)
C3—H3	0.9300	C24—H24	0.9300
C4—C5	1.3900	C25—C26	1.368 (8)
C4—H4	0.9300	C25—H25	0.9300
C5—C6	1.3900	C26—C27	1.384 (6)
C5—H5	0.9300	C26—H26	0.9300
C6—C7	1.3900	C27—C28	1.492 (6)
C6—H6A	0.9300	C28—C29	1.372 (6)
C7—H7	0.9300	C29—C30	1.382 (7)
C8—C9	1.485 (7)	C29—H29	0.9300
C9—C14	1.369 (6)	C30—C31	1.349 (7)
C9—C10	1.377 (6)	C30—H30	0.9300
C10—C11	1.384 (8)	C31—C32	1.379 (7)
C10—H10	0.9300	C31—H31	0.9300
C11—C12	1.374 (9)	C32—C33	1.488 (7)
C11—H11	0.9300	C33—H33A	0.9600
C12—C13	1.352 (8)	C33—H33B	0.9600
C12—H12	0.9300	C33—H33C	0.9600
C13—C14	1.369 (8)		
O3—Zn1—N1	119.74 (16)	O5—C15—C16	123.5 (5)
O3—Zn1—N2	124.65 (15)	O6—C15—C16	113.6 (5)
N1—Zn1—N2	80.07 (17)	C21—C16—C17	119.5 (5)
O3—Zn1—O1	102.78 (16)	C21—C16—C15	119.1 (5)
N1—Zn1—O1	132.61 (16)	C17—C16—C15	121.3 (5)
N2—Zn1—O1	93.04 (16)	C16—C17—C18	119.0 (6)
O3—Zn1—O2	104.33 (15)	C16—C17—H17	120.5
N1—Zn1—O2	91.25 (15)	C18—C17—H17	120.5
N2—Zn1—O2	127.89 (14)	C19—C18—C17	120.0 (6)
O1—Zn1—O2	56.64 (17)	C19—C18—H18	120.0
C27—N1—C23	119.8 (4)	C17—C18—H18	120.0
C27—N1—Zn1	113.4 (3)	C20—C19—C18	120.4 (6)
C23—N1—Zn1	126.6 (4)	C20—C19—H19	119.8
C28—N2—C32	119.6 (4)	C18—C19—H19	119.8
C28—N2—Zn1	113.8 (3)	C19—C20—C21	119.4 (6)
C32—N2—Zn1	126.4 (3)	C19—C20—H20	120.3
C1—O1—Zn1	98.0 (4)	C21—C20—H20	120.3
C1—O2—Zn1	90.6 (4)	C20—C21—C16	121.6 (6)
C8—O3—Zn1	117.4 (3)	C20—C21—H21	119.2
C15—O6—H6	109.5	C16—C21—H21	119.2
O2—C1—O1	114.7 (6)	C23—C22—H22A	109.5
O2—C1—C2	124.5 (7)	C23—C22—H22B	109.5

O1—C1—C2	120.8 (7)	H22A—C22—H22B	109.5
C3—C2—C7	120.0	C23—C22—H22C	109.5
C3—C2—C1	122.6 (5)	H22A—C22—H22C	109.5
C7—C2—C1	117.3 (5)	H22B—C22—H22C	109.5
C4—C3—C2	120.0	N1—C23—C24	119.8 (5)
C4—C3—H3	120.0	N1—C23—C22	117.9 (5)
C2—C3—H3	120.0	C24—C23—C22	122.2 (6)
C3—C4—C5	120.0	C25—C24—C23	120.2 (6)
C3—C4—H4	120.0	C25—C24—H24	119.9
C5—C4—H4	120.0	C23—C24—H24	119.9
C4—C5—C6	120.0	C24—C25—C26	120.0 (6)
C4—C5—H5	120.0	C24—C25—H25	120.0
C6—C5—H5	120.0	C26—C25—H25	120.0
C7—C6—C5	120.0	C25—C26—C27	118.4 (6)
C7—C6—H6A	120.0	C25—C26—H26	120.8
C5—C6—H6A	120.0	C27—C26—H26	120.8
C6—C7—C2	120.0	N1—C27—C26	121.7 (5)
C6—C7—H7	120.0	N1—C27—C28	115.9 (4)
C2—C7—H7	120.0	C26—C27—C28	122.4 (5)
O4—C8—O3	123.0 (5)	N2—C28—C29	121.7 (5)
O4—C8—C9	120.3 (5)	N2—C28—C27	115.4 (4)
O3—C8—C9	116.7 (4)	C29—C28—C27	122.9 (5)
C14—C9—C10	118.3 (5)	C28—C29—C30	118.8 (5)
C14—C9—C8	120.0 (5)	C28—C29—H29	120.6
C10—C9—C8	121.6 (5)	C30—C29—H29	120.6
C9—C10—C11	121.0 (6)	C31—C30—C29	119.3 (5)
C9—C10—H10	119.5	C31—C30—H30	120.4
C11—C10—H10	119.5	C29—C30—H30	120.4
C12—C11—C10	119.1 (6)	C30—C31—C32	120.5 (6)
C12—C11—H11	120.5	C30—C31—H31	119.7
C10—C11—H11	120.5	C32—C31—H31	119.7
C13—C12—C11	120.0 (7)	N2—C32—C31	120.0 (5)
C13—C12—H12	120.0	N2—C32—C33	117.5 (5)
C11—C12—H12	120.0	C31—C32—C33	122.5 (5)
C12—C13—C14	120.8 (7)	C32—C33—H33A	109.5
C12—C13—H13	119.6	C32—C33—H33B	109.5
C14—C13—H13	119.6	H33A—C33—H33B	109.5
C9—C14—C13	120.8 (6)	C32—C33—H33C	109.5
C9—C14—H14	119.6	H33A—C33—H33C	109.5
C13—C14—H14	119.6	H33B—C33—H33C	109.5
O5—C15—O6	122.9 (5)		
O3—Zn1—N1—C27	134.4 (3)	C8—C9—C10—C11	179.1 (6)
N2—Zn1—N1—C27	10.0 (3)	C9—C10—C11—C12	-0.2 (11)
O1—Zn1—N1—C27	-74.9 (4)	C10—C11—C12—C13	-2.5 (13)
O2—Zn1—N1—C27	-118.3 (3)	C11—C12—C13—C14	1.9 (13)
O3—Zn1—N1—C23	-50.8 (4)	C10—C9—C14—C13	-4.0 (9)
N2—Zn1—N1—C23	-175.2 (4)	C8—C9—C14—C13	-179.8 (6)
O1—Zn1—N1—C23	99.9 (4)	C12—C13—C14—C9	1.4 (11)
O2—Zn1—N1—C23	56.5 (4)	O5—C15—C16—C21	-18.5 (7)

supplementary materials

O3—Zn1—N2—C28	-130.2 (3)	O6—C15—C16—C21	161.4 (4)
N1—Zn1—N2—C28	-10.8 (3)	O5—C15—C16—C17	158.1 (5)
O1—Zn1—N2—C28	122.0 (3)	O6—C15—C16—C17	-22.0 (6)
O2—Zn1—N2—C28	73.0 (3)	C21—C16—C17—C18	-0.3 (8)
O3—Zn1—N2—C32	55.2 (4)	C15—C16—C17—C18	-176.9 (5)
N1—Zn1—N2—C32	174.6 (4)	C16—C17—C18—C19	-0.4 (9)
O1—Zn1—N2—C32	-52.6 (4)	C17—C18—C19—C20	-0.8 (10)
O2—Zn1—N2—C32	-101.6 (4)	C18—C19—C20—C21	2.6 (10)
O3—Zn1—O1—C1	100.7 (4)	C19—C20—C21—C16	-3.4 (9)
N1—Zn1—O1—C1	-53.5 (5)	C17—C16—C21—C20	2.2 (8)
N2—Zn1—O1—C1	-132.8 (4)	C15—C16—C21—C20	178.9 (5)
O2—Zn1—O1—C1	1.8 (4)	C27—N1—C23—C24	-0.5 (7)
O3—Zn1—O2—C1	-97.8 (4)	Zn1—N1—C23—C24	-175.0 (4)
N1—Zn1—O2—C1	141.0 (4)	C27—N1—C23—C22	-178.6 (4)
N2—Zn1—O2—C1	62.6 (4)	Zn1—N1—C23—C22	6.9 (6)
O1—Zn1—O2—C1	-1.8 (4)	N1—C23—C24—C25	2.1 (8)
N1—Zn1—O3—C8	-45.9 (4)	C22—C23—C24—C25	-179.9 (5)
N2—Zn1—O3—C8	52.9 (4)	C23—C24—C25—C26	-2.7 (9)
O1—Zn1—O3—C8	155.8 (3)	C24—C25—C26—C27	1.6 (8)
O2—Zn1—O3—C8	-145.8 (3)	C23—N1—C27—C26	-0.6 (6)
Zn1—O2—C1—O1	2.8 (6)	Zn1—N1—C27—C26	174.6 (3)
Zn1—O2—C1—C2	-174.3 (6)	C23—N1—C27—C28	176.8 (4)
Zn1—O1—C1—O2	-3.0 (7)	Zn1—N1—C27—C28	-7.9 (5)
Zn1—O1—C1—C2	174.2 (5)	C25—C26—C27—N1	0.1 (7)
O2—C1—C2—C3	175.0 (5)	C25—C26—C27—C28	-177.2 (5)
O1—C1—C2—C3	-1.9 (8)	C32—N2—C28—C29	3.7 (6)
O2—C1—C2—C7	-2.0 (8)	Zn1—N2—C28—C29	-171.4 (3)
O1—C1—C2—C7	-178.9 (5)	C32—N2—C28—C27	-175.3 (4)
C7—C2—C3—C4	0.0	Zn1—N2—C28—C27	9.7 (4)
C1—C2—C3—C4	-176.9 (5)	N1—C27—C28—N2	-1.2 (5)
C2—C3—C4—C5	0.0	C26—C27—C28—N2	176.3 (4)
C3—C4—C5—C6	0.0	N1—C27—C28—C29	179.9 (4)
C4—C5—C6—C7	0.0	C26—C27—C28—C29	-2.7 (7)
C5—C6—C7—C2	0.0	N2—C28—C29—C30	-1.8 (7)
C3—C2—C7—C6	0.0	C27—C28—C29—C30	177.1 (4)
C1—C2—C7—C6	177.1 (4)	C28—C29—C30—C31	-0.5 (8)
Zn1—O3—C8—O4	-5.0 (6)	C29—C30—C31—C32	0.8 (8)
Zn1—O3—C8—C9	172.6 (3)	C28—N2—C32—C31	-3.3 (7)
O4—C8—C9—C14	4.8 (7)	Zn1—N2—C32—C31	171.1 (3)
O3—C8—C9—C14	-172.9 (5)	C28—N2—C32—C33	177.4 (4)
O4—C8—C9—C10	-170.8 (5)	Zn1—N2—C32—C33	-8.2 (6)
O3—C8—C9—C10	11.5 (7)	C30—C31—C32—N2	1.0 (8)
C14—C9—C10—C11	3.4 (9)	C30—C31—C32—C33	-179.7 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C24—H24 \cdots O5 ⁱ	0.93	2.44	3.340 (8)	162
C33—H33C \cdots O1	0.96	2.52	3.199 (8)	128

C22—H22A···O2	0.96	2.57	3.289 (7)	132
O6—H6···O4 ⁱⁱ	0.82	1.85	2.655 (5)	166
C31—H31···Cg1 ⁱⁱⁱ	0.93	2.79	3.708 (7)	170

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $x, y, z+1$; (iii) $-x+1, -y+1, -z$.

