

(1*R*,2*R*)-1,2-Diphenyl-1,2-bis(1*H*-tetrazol-1-yl)ethane

Franz Werner,^{a*}‡ Kurt Mereiter,^b Kenji Tokuno,^a Yuki Inagaki^a and Miki Hasegawa^{a*}

^aAoyama-Gakuin University, College of Science and Engineering, Department of Chemistry and Biological Science, 5-10-1 Fuchinobe, Sagamihara, Kanagawa 229-8558, Japan, and ^bVienna University of Technology, Institute of Chemical Technologies and Analytics, Getreidemarkt 9/164, 1060 Vienna, Austria
Correspondence e-mail: fwerner@mail.zserv.tuwien.ac.at, hasemiki@chem.aoyama.ac.jp


Received 21 September 2009; accepted 7 October 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.035; wR factor = 0.092; data-to-parameter ratio = 11.8.

The title compound, $C_{16}H_{14}N_8$, is a new chiral ligand designed for applications in supramolecular chemistry and Fe^{2+} spin-crossover complexes. The crystal structure shows a herringbone arrangement of the molecules, which are mutually linked via intermolecular C—H···N interactions mainly donated by the alkyl and tetrazole H atoms.

Related literature

For the general synthetic procedure, see: Kamiya & Saito (1973). For the crystal structure of the chiral starting material, see: Jones *et al.* (2003). For studies on the crystal structures and packing of di-tetrazolylalkanes, see: Grunert *et al.* (2005); Absmeier *et al.* (2006). For supramolecular compounds made up of di-tetrazolylalkanes, see: Liu *et al.* (2008, 2009); Yu *et al.* (2008). For Fe^{2+} spin-crossover complexes based on di-tetrazolylalkanes, see: Grunert *et al.* (2004); Quesada *et al.* (2007); Bialonska *et al.* (2008). The absolute structure of the title compound could not be determined from the diffraction data but was known from the chiral precursor compound (1*R*,2*R*)-(+)-1,2-diphenyl-1,2-ethanediamine, see: Jones *et al.* (2003).



‡ Present address: Vienna University of Technology, Institute of Applied Synthetic Chemistry, Getreidemarkt 9/163, 1060 Vienna, Austria.

Experimental

Crystal data

$C_{16}H_{14}N_8$
 $M_r = 318.35$
Orthorhombic, $P2_12_12_1$
 $a = 8.3088$ (4) Å
 $b = 11.2802$ (6) Å
 $c = 16.5187$ (9) Å

$V = 1548.21$ (14) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 100$ K
 $0.65 \times 0.55 \times 0.46$ mm

Data collection

Bruker SMART APEX CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2007)
 $T_{min} = 0.86$, $T_{max} = 0.96$

17102 measured reflections
2551 independent reflections
2475 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.018$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.092$
 $S = 1.10$
2551 reflections

217 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.30$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1···N7 ⁱ	0.95	2.70	3.430 (2)	134
C2—H2···N7 ⁱ	1.00	2.55	3.392 (2)	142
C2—H2···N8 ⁱ	1.00	2.53	3.506 (2)	165
C3—H3···N3 ⁱⁱ	1.00	2.46	3.351 (2)	149
C4—H4···N3 ⁱⁱ	0.95	2.63	3.315 (2)	130
C4—H4···N4 ⁱⁱ	0.95	2.67	3.543 (2)	154
C6—H6···N2	0.95	2.62	3.256 (2)	124
C7—H7···N6 ⁱⁱⁱ	0.95	2.63	3.339 (2)	132
C12—H12···N2 ⁱⁱ	0.95	2.71	3.655 (2)	171
C13—H13···N7 ^{iv}	0.95	2.74	3.379 (2)	125

Symmetry codes: (i) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x - \frac{1}{2}, -y + \frac{3}{2}, -z$; (iii) $-x + \frac{3}{2}, -y + 1, z - \frac{1}{2}$; (iv) $x - 1, y, z$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

FW is grateful to the Japan Society for the Promotion of Science for financial support through a fellowship. MH acknowledges support from a Grant-in-Aid for Young Scientists A (No. 20685011) and a High-Tech Research Center project for private universities with the matching fund subsidy of MEXT in Japan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2147).

References

- Absmeier, A., Bartel, M., Carbonera, C., Jameson, G. N. L., Weinberger, P., Caneschi, A., Mereiter, K., Letard, J.-F. & Linert, W. (2006). *Chem. Eur. J.* **12**, 2235–2243.
Bialonska, A., Bronisz, R. & Weselski, M. (2008). *Inorg. Chem.* **47**, 4436–4438.
Bruker (2007). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.

- Grunert, M., Schweifer, J., Weinberger, P., Linert, W., Mereiter, K., Hilscher, G., Müller, M., Wiesinger, G. & van Koningsbruggen, P. J. (2004). *Inorg. Chem.* **43**, 155–165.
- Grunert, C. M., Weinberger, P., Schweifer, J., Hampel, C., Stassen, A. F., Mereiter, K. & Linert, W. (2005). *J. Mol. Struct.* **733**, 41–52.
- Jones, M. D., Almeida Paz, F. A., Davies, J. E. & Johnson, B. F. G. (2003). *Acta Cryst. E* **59**, o455–o457.
- Kamiya, T. & Saito, Y. (1973). *Offenlegungsschrift* 2147023 (Patent).
- Liu, P.-P., Cheng, A.-L., Yue, Q., Liu, N., Sun, W.-W. & Gao, E.-Q. (2008). *Cryst. Growth Des.* **8**, 1668–1674.
- Liu, P.-P., Wang, Y.-Q., Tian, C.-Y., Peng, H.-Q. & Gao, E.-Q. (2009). *J. Mol. Struct.* **920**, 459–465.
- Quesada, M., Kooijman, H., Gamez, P., Sanchez Costa, J., van Koningsbruggen, P. J., Weinberger, P., Reissner, M., Spek, A. L., Haasnoot, J. G. & Reedijk, J. (2007). *Dalton Trans.* pp. 5434–5440.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yu, J.-H., Mereiter, K., Hassan, N., Feldgitscher, C. & Linert, W. (2008). *Cryst. Growth Des.* **8**, 1535–1540.

supporting information

Acta Cryst. (2009). E65, o2726–o2727 [https://doi.org/10.1107/S160053680904094X]

(*1R,2R*)-1,2-Diphenyl-1,2-bis(1*H*-tetrazol-1-yl)ethane

Franz Werner, Kurt Mereiter, Kenji Tokuno, Yuki Inagaki and Miki Hasegawa

S1. Comment

Bifunctional molecules containing two 1*H*-tetrazol-1-yl groups at both ends of suitable spacer moieties like flexible alkanes or stiff arenes are of growing interest in supramolecular chemistry (Liu *et al.*, 2008, 2009; Yu *et al.*, 2008) and in the construction of new Fe²⁺-based spin-crossover complexes (Grunert *et al.*, 2004; Quesada *et al.*, 2007; Bialonska *et al.*, 2008). In continuation of a previous study on the crystal structures and packing of α - ω -bis(tetrazol-1-yl)-alkanes (Grunert *et al.*, 2005; Absmeier *et al.*, 2006) the title compound, (*1R,2R*)-1,2-diphenyl-1,2-di-(1*H*-tetrazol-1-yl)ethane, became of interest as an example for a chiral bis-tetrazolyl ligand. It was obtained by a standard reaction (Kamiya & Saito, 1973) from the chiral starting material (*1R,2R*)-(+)1,2-diphenyl-1,2-ethanediamine (*cf.* experimental).

The title compound crystallizes in the orthorhombic chiral space group $P2_12_12_1$, with one molecule in the asymmetric unit (Fig. 1). Bond lengths and bond angles in the molecule are normal. The two tetrazole rings are attached to the central ethane group in a (-)-synclinal geometry [$N1—C2—C3—N5 = -53.3 (1)$ °], the two phenyl rings in a (+)-synclinal geometry [$N1—C2—C3—N5 = 60.6 (1)$ °]. The point group symmetry of the free molecule is C_2 . In the solid state it deviates significantly from this symmetry by intermolecular forces, as can be seen particularly from the distinctly differing torsion angles angles [$C2—C3—N5—C4 = 114.3 (1)$ ° and $C3—C2—N1—C1 = 157.8 (1)$ °] involving the tetrazolyl groups. The corresponding angles involving the phenyl rings differ less [$C3—C2—C5—C6 = 76.0 (1)$ ° and $C2—C3—C11—C16 = 52.6 (2)$ °].


In the crystal the molecules are stacked in a typical herring-bone manner (Fig. 2). π – π -stacking is absent but adjacent molecules are linked by weak intermolecular C—H···N interactions (Table 1) between mainly the alkyl CH groups and the tetrazole nitrogen atoms. These are accompanied by somewhat longer C—H···N interactions of the tetrazole CH groups and some phenyl CH groups (Table 1).

S2. Experimental


The title compound was prepared according to the general procedure given by (Kamiya & Saito, 1973). A solution of 2.0 g of (*1R,2R*)-(+)1,2-diphenyl-1,2-ethanediamine (9.42 mmol, Kanto Chemical), 1.41 g of sodium azide (21.7 mmol, Wako, min. 98.0%) and 4.19 g of triethyl orthoformate (28.3 mmol, Sigma-Aldrich, 98%) in 120 ml of glacial acetic acid (Kanto Chemical, 99.5%), was stirred for 2 h at a temperature of 343 – 353 K. After cooling down to rt the solvent was distilled off and 20 ml of distilled water were added, whereupon a yellow solid precipitated. The suspension was stored in the refrigerator overnight, then the product was obtained by suction filtration and was washed with distilled water. Drying under vacuum yielded 0.291 g (9.7%) of the title compound as a colourless microcrystalline powder. Crystals suitable for X-ray diffraction were obtained by recrystallization from methanol. Elemental analysis (Micro Corder JM10, J-Science Lab): C (calculated 60.37%/found 59.96%), H (4.43/4.56), N (35.20/34.80). NMR (Bruker DPX-200): ¹H (DMSO-d₆): δ 7.26–7.37 (*m*, 3 H, Ph—H), 7.50 (*s*, 1 H, CH), 7.66–7.70 (*m*, 2 H, Ph—H), 9.55 (*s*, 1 H, tetrazole). ¹³C (DMSO-d₆): δ 63.8 (CH); 128.5, 129.1, 129.5, 134.0 (Ph); 143.7 (tetrazole).

S3. Refinement

The absolute structure of the title compound could not be determined from the diffraction data but was known from the chiral precursor compound (*1R,2R*)-(+)-1,2-diphenyl-1,2-ethanediamine. In the final cycles of refinement, in the absence of significant anomalous scattering effects, Friedel pairs were merged and $\Delta f''$ set to zero. All the H-atoms were placed in calculated positions and treated as riding: C-H = 0.95 - 1.0 Å, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

**Figure 1**

A view of the molecular structure of the title compound, with displacement ellipsoids drawn at the 50% probability level.

**Figure 2**

Perspective view, along the *a*-axis, of the crystal packing of the title compound. The shortest hydrogen bonds (C3—H3···N3) are represented by cyan dashed lines.

(1*R*,2*R*)-1,2-Diphenyl-1,2-bis(1*H*-tetrazol-1-yl)ethane

Crystal data

$C_{16}H_{14}N_8$
 $M_r = 318.35$
Orthorhombic, $P2_12_12_1$
Hall symbol: P 2ac 2ab
 $a = 8.3088 (4)$ Å
 $b = 11.2802 (6)$ Å
 $c = 16.5187 (9)$ Å
 $V = 1548.21 (14)$ Å³
 $Z = 4$

$F(000) = 664$
 $D_x = 1.366 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 6154 reflections
 $\theta = 2.5\text{--}30.0^\circ$
 $\mu = 0.09 \text{ mm}^{-1}$
 $T = 100$ K
Prism, colourless
 $0.65 \times 0.55 \times 0.46$ mm

Data collection

Bruker SMART APEX CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω and φ scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2007)
 $T_{\min} = 0.86$, $T_{\max} = 0.96$

17102 measured reflections
2551 independent reflections
2475 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.018$
 $\theta_{\max} = 30.0^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -11 \rightarrow 11$
 $k = -15 \rightarrow 14$
 $l = -23 \rightarrow 23$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.092$
 $S = 1.10$
2551 reflections
217 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0636P)^2 + 0.1943P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$
Absolute structure: known from the chirality of
the precursor used; Friedel pairs merged

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.98462 (13)	0.55207 (9)	0.07313 (6)	0.01376 (19)
N2	1.00108 (15)	0.64765 (10)	0.02443 (7)	0.0189 (2)
N3	1.14022 (15)	0.63675 (11)	-0.01065 (7)	0.0210 (2)
N4	1.21530 (15)	0.53546 (11)	0.01269 (7)	0.0220 (2)
N5	0.85774 (13)	0.73023 (9)	0.18013 (6)	0.0135 (2)
N6	0.90650 (14)	0.71246 (10)	0.25717 (6)	0.0176 (2)
N7	0.99083 (15)	0.80496 (10)	0.27687 (7)	0.0192 (2)
N8	0.99906 (15)	0.88385 (10)	0.21407 (7)	0.0200 (2)
C1	1.11653 (16)	0.48451 (12)	0.06448 (8)	0.0189 (2)
H1	1.1356	0.4115	0.0915	0.023*
C2	0.83462 (14)	0.52879 (10)	0.11786 (7)	0.0125 (2)
H2	0.8613	0.4869	0.1695	0.015*
C3	0.74954 (14)	0.64654 (10)	0.13802 (7)	0.0123 (2)
H3	0.7171	0.6840	0.0856	0.015*
C4	0.91482 (17)	0.83523 (11)	0.15502 (8)	0.0173 (2)

H4	0.8974	0.8691	0.1031	0.021*
C5	0.72240 (15)	0.45132 (11)	0.06816 (7)	0.0141 (2)
C6	0.69622 (17)	0.47520 (12)	-0.01377 (7)	0.0175 (2)
H6	0.7531	0.5380	-0.0393	0.021*
C7	0.58752 (18)	0.40772 (13)	-0.05813 (8)	0.0214 (3)
H7	0.5714	0.4236	-0.1140	0.026*
C8	0.50251 (19)	0.31696 (12)	-0.02063 (9)	0.0237 (3)
H8	0.4270	0.2715	-0.0507	0.028*
C9	0.52807 (18)	0.29277 (12)	0.06099 (9)	0.0239 (3)
H9	0.4699	0.2308	0.0867	0.029*
C10	0.63881 (17)	0.35930 (11)	0.10515 (8)	0.0193 (2)
H10	0.6572	0.3417	0.1606	0.023*
C11	0.59769 (15)	0.62608 (11)	0.18692 (7)	0.0144 (2)
C12	0.45409 (17)	0.67425 (12)	0.15886 (8)	0.0188 (3)
H12	0.4529	0.7189	0.1101	0.023*
C13	0.31198 (17)	0.65725 (13)	0.20212 (9)	0.0240 (3)
H13	0.2144	0.6909	0.1830	0.029*
C14	0.31241 (18)	0.59140 (14)	0.27300 (9)	0.0257 (3)
H14	0.2155	0.5804	0.3026	0.031*
C15	0.45537 (19)	0.54143 (13)	0.30064 (9)	0.0249 (3)
H15	0.4555	0.4953	0.3487	0.030*
C16	0.59844 (17)	0.55868 (12)	0.25815 (8)	0.0198 (3)
H16	0.6959	0.5249	0.2774	0.024*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0135 (4)	0.0139 (4)	0.0139 (4)	-0.0001 (4)	0.0009 (4)	0.0007 (3)
N2	0.0194 (5)	0.0180 (5)	0.0192 (5)	-0.0002 (4)	0.0040 (4)	0.0050 (4)
N3	0.0196 (5)	0.0222 (5)	0.0214 (5)	-0.0008 (4)	0.0051 (4)	0.0015 (4)
N4	0.0176 (5)	0.0245 (6)	0.0238 (5)	0.0014 (4)	0.0047 (4)	0.0003 (4)
N5	0.0126 (4)	0.0143 (4)	0.0134 (4)	-0.0007 (4)	-0.0011 (4)	-0.0005 (3)
N6	0.0184 (5)	0.0212 (5)	0.0132 (4)	-0.0027 (4)	-0.0027 (4)	-0.0006 (4)
N7	0.0191 (5)	0.0201 (5)	0.0186 (4)	-0.0014 (4)	-0.0019 (4)	-0.0035 (4)
N8	0.0196 (5)	0.0168 (5)	0.0235 (5)	-0.0016 (4)	-0.0041 (4)	-0.0014 (4)
C1	0.0155 (5)	0.0192 (6)	0.0219 (6)	0.0029 (5)	0.0017 (5)	0.0008 (5)
C2	0.0111 (5)	0.0139 (5)	0.0124 (4)	-0.0007 (4)	0.0011 (4)	0.0006 (4)
C3	0.0110 (5)	0.0135 (5)	0.0124 (4)	-0.0006 (4)	-0.0006 (4)	-0.0006 (4)
C4	0.0172 (6)	0.0143 (5)	0.0204 (5)	-0.0016 (4)	-0.0030 (4)	0.0013 (4)
C5	0.0129 (5)	0.0135 (5)	0.0157 (5)	-0.0004 (4)	-0.0002 (4)	-0.0017 (4)
C6	0.0183 (6)	0.0188 (5)	0.0155 (5)	-0.0001 (5)	0.0003 (4)	-0.0011 (4)
C7	0.0228 (6)	0.0226 (6)	0.0189 (5)	0.0018 (5)	-0.0037 (5)	-0.0057 (5)
C8	0.0221 (6)	0.0177 (6)	0.0314 (7)	0.0001 (5)	-0.0072 (6)	-0.0067 (5)
C9	0.0233 (6)	0.0169 (6)	0.0315 (7)	-0.0056 (5)	-0.0027 (5)	0.0007 (5)
C10	0.0206 (6)	0.0160 (5)	0.0213 (5)	-0.0029 (5)	-0.0018 (5)	0.0019 (5)
C11	0.0130 (5)	0.0152 (5)	0.0151 (5)	-0.0016 (4)	0.0019 (4)	-0.0031 (4)
C12	0.0146 (6)	0.0195 (6)	0.0223 (6)	-0.0002 (4)	0.0003 (4)	-0.0018 (5)
C13	0.0133 (6)	0.0265 (7)	0.0324 (7)	-0.0001 (5)	0.0035 (5)	-0.0054 (6)

C14	0.0197 (6)	0.0262 (7)	0.0310 (7)	-0.0059 (5)	0.0106 (5)	-0.0075 (6)
C15	0.0264 (7)	0.0262 (7)	0.0222 (6)	-0.0056 (5)	0.0090 (5)	0.0002 (5)
C16	0.0190 (6)	0.0222 (6)	0.0183 (5)	-0.0013 (5)	0.0025 (5)	0.0006 (5)

Geometric parameters (\AA , $^{\circ}$)

N1—C1	1.3425 (16)	C6—C7	1.3900 (18)
N1—N2	1.3521 (14)	C6—H6	0.9500
N1—C2	1.4725 (15)	C7—C8	1.389 (2)
N2—N3	1.2991 (16)	C7—H7	0.9500
N3—N4	1.3577 (17)	C8—C9	1.392 (2)
N4—C1	1.3176 (17)	C8—H8	0.9500
N5—C4	1.3416 (16)	C9—C10	1.3935 (18)
N5—N6	1.3504 (14)	C9—H9	0.9500
N5—C3	1.4777 (15)	C10—H10	0.9500
N6—N7	1.2984 (16)	C11—C12	1.3906 (18)
N7—N8	1.3685 (15)	C11—C16	1.4008 (17)
N8—C4	1.3199 (17)	C12—C13	1.3934 (19)
C1—H1	0.9500	C12—H12	0.9500
C2—C5	1.5188 (16)	C13—C14	1.387 (2)
C2—C3	1.5410 (16)	C13—H13	0.9500
C2—H2	1.0000	C14—C15	1.392 (2)
C3—C11	1.5158 (17)	C14—H14	0.9500
C3—H3	1.0000	C15—C16	1.3941 (19)
C4—H4	0.9500	C15—H15	0.9500
C5—C10	1.3903 (17)	C16—H16	0.9500
C5—C6	1.3971 (16)		
C1—N1—N2	107.87 (11)	C7—C6—C5	120.41 (12)
C1—N1—C2	130.02 (10)	C7—C6—H6	119.8
N2—N1—C2	121.76 (10)	C5—C6—H6	119.8
N3—N2—N1	106.26 (11)	C8—C7—C6	119.93 (12)
N2—N3—N4	111.21 (11)	C8—C7—H7	120.0
C1—N4—N3	105.39 (11)	C6—C7—H7	120.0
C4—N5—N6	108.45 (10)	C7—C8—C9	119.91 (13)
C4—N5—C3	129.31 (10)	C7—C8—H8	120.0
N6—N5—C3	122.09 (10)	C9—C8—H8	120.0
N7—N6—N5	106.19 (10)	C8—C9—C10	120.14 (13)
N6—N7—N8	111.07 (10)	C8—C9—H9	119.9
C4—N8—N7	105.28 (10)	C10—C9—H9	119.9
N4—C1—N1	109.26 (11)	C5—C10—C9	120.13 (12)
N4—C1—H1	125.4	C5—C10—H10	119.9
N1—C1—H1	125.4	C9—C10—H10	119.9
N1—C2—C5	110.55 (9)	C12—C11—C16	119.73 (12)
N1—C2—C3	110.06 (9)	C12—C11—C3	118.49 (11)
C5—C2—C3	109.34 (9)	C16—C11—C3	121.77 (11)
N1—C2—H2	109.0	C11—C12—C13	120.15 (13)
C5—C2—H2	109.0	C11—C12—H12	119.9

C3—C2—H2	109.0	C13—C12—H12	119.9
N5—C3—C11	110.66 (9)	C14—C13—C12	120.31 (13)
N5—C3—C2	111.92 (9)	C14—C13—H13	119.8
C11—C3—C2	111.46 (10)	C12—C13—H13	119.8
N5—C3—H3	107.5	C13—C14—C15	119.75 (12)
C11—C3—H3	107.5	C13—C14—H14	120.1
C2—C3—H3	107.5	C15—C14—H14	120.1
N8—C4—N5	109.01 (11)	C14—C15—C16	120.40 (13)
N8—C4—H4	125.5	C14—C15—H15	119.8
N5—C4—H4	125.5	C16—C15—H15	119.8
C10—C5—C6	119.47 (12)	C15—C16—C11	119.65 (13)
C10—C5—C2	119.92 (10)	C15—C16—H16	120.2
C6—C5—C2	120.55 (11)	C11—C16—H16	120.2
C1—N1—N2—N3	-0.94 (14)	C3—N5—C4—N8	175.92 (12)
C2—N1—N2—N3	-174.78 (10)	N1—C2—C5—C10	137.82 (11)
N1—N2—N3—N4	0.90 (14)	C3—C2—C5—C10	-100.87 (13)
N2—N3—N4—C1	-0.50 (15)	N1—C2—C5—C6	-45.31 (15)
C4—N5—N6—N7	-0.14 (14)	C3—C2—C5—C6	76.01 (13)
C3—N5—N6—N7	-176.11 (11)	C10—C5—C6—C7	0.03 (19)
N5—N6—N7—N8	-0.09 (14)	C2—C5—C6—C7	-176.86 (12)
N6—N7—N8—C4	0.30 (14)	C5—C6—C7—C8	0.9 (2)
N3—N4—C1—N1	-0.11 (15)	C6—C7—C8—C9	-0.9 (2)
N2—N1—C1—N4	0.66 (15)	C7—C8—C9—C10	-0.1 (2)
C2—N1—C1—N4	173.81 (12)	C6—C5—C10—C9	-0.98 (19)
C1—N1—C2—C5	-81.28 (15)	C2—C5—C10—C9	175.93 (12)
N2—N1—C2—C5	91.05 (13)	C8—C9—C10—C5	1.0 (2)
C1—N1—C2—C3	157.83 (12)	N5—C3—C11—C12	108.50 (12)
N2—N1—C2—C3	-29.83 (14)	C2—C3—C11—C12	-126.26 (12)
C4—N5—C3—C11	-120.73 (14)	N5—C3—C11—C16	-72.66 (14)
N6—N5—C3—C11	54.32 (14)	C2—C3—C11—C16	52.57 (15)
C4—N5—C3—C2	114.30 (13)	C16—C11—C12—C13	1.08 (19)
N6—N5—C3—C2	-70.65 (13)	C3—C11—C12—C13	179.94 (12)
N1—C2—C3—N5	-53.30 (12)	C11—C12—C13—C14	-0.6 (2)
C5—C2—C3—N5	-174.92 (9)	C12—C13—C14—C15	-0.5 (2)
N1—C2—C3—C11	-177.83 (9)	C13—C14—C15—C16	1.0 (2)
C5—C2—C3—C11	60.56 (12)	C14—C15—C16—C11	-0.4 (2)
N7—N8—C4—N5	-0.38 (15)	C12—C11—C16—C15	-0.58 (19)
N6—N5—C4—N8	0.34 (15)	C3—C11—C16—C15	-179.40 (12)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C1—H1···N7 ⁱ	0.95	2.70	3.430 (2)	134
C2—H2···N7 ⁱ	1.00	2.55	3.392 (2)	142
C2—H2···N8 ^j	1.00	2.53	3.506 (2)	165
C3—H3···N3 ⁱⁱ	1.00	2.46	3.351 (2)	149
C4—H4···N3 ⁱⁱ	0.95	2.63	3.315 (2)	130

C4—H4···N4 ⁱⁱ	0.95	2.67	3.543 (2)	154
C6—H6···N2	0.95	2.62	3.256 (2)	124
C7—H7···N6 ⁱⁱⁱ	0.95	2.63	3.339 (2)	132
C12—H12···N2 ⁱⁱ	0.95	2.71	3.655 (2)	171
C13—H13···N7 ^{iv}	0.95	2.74	3.379 (2)	125

Symmetry codes: (i) $-x+2, y-1/2, -z+1/2$; (ii) $x-1/2, -y+3/2, -z$; (iii) $-x+3/2, -y+1, z-1/2$; (iv) $x-1, y, z$.