

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

Tetraaquabis(2-oxo-1,2-dihydropyridine-5-sulfonato- $\kappa O^2$ )zinc(II)Zhi-Biao Zhu,<sup>a</sup> Shan Gao<sup>a</sup> and Seik Weng Ng<sup>b\*</sup><sup>a</sup>College of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 28 September 2009; accepted 30 September 2009

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(C-C) = 0.003$  Å;  $R$  factor = 0.023;  $wR$  factor = 0.066; data-to-parameter ratio = 13.5.

The metal atom in the title compound,  $[Zn(C_5H_4NO_4S)_2(H_2O)_4]$ , lies on a center of inversion and is linked to the anionic ligand through the carbonyl O atom. In the crystal structure, the 2-oxo-1,2-dihydropyridine-5-sulfonate ligand interacts with other molecules through  $N-H \cdots O$  and  $O-H \cdots O$  hydrogen bonds, forming a three-dimensional network structure.

## Related literature

For the crystal structure of another zwitterionic tetraaquabis(amide)-metal<sup>II</sup> complex, see: Gao *et al.* (2004).



## Experimental

## Crystal data

 $[Zn(C_5H_4NO_4S)_2(H_2O)_4]$  $M_r = 485.74$ Monoclinic,  $P2_1/c$  $a = 6.7701$  (2) Å $b = 13.9725$  (5) Å $c = 10.0343$  (3) Å $\beta = 115.331$  (2)°  
 $V = 857.93$  (5) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation $\mu = 1.74$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.21 \times 0.16 \times 0.16$  mm

## Data collection

Rigaku R-Axis RAPID IP diffractometer  
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)  
 $T_{\min} = 0.711$ ,  $T_{\max} = 0.768$ 8224 measured reflections  
1951 independent reflections  
1866 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.023$   
 $wR(F^2) = 0.066$   
 $S = 1.06$   
1951 reflections  
144 parameters  
5 restraintsH atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.37$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.39$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$N1-H1 \cdots O2^i$	0.85 (1)	1.99 (1)	2.790 (2)	157 (2)
$O1w-H11 \cdots O2^{ii}$	0.84 (1)	1.98 (1)	2.809 (2)	171 (2)
$O1w-H12 \cdots O3^{iii}$	0.84 (1)	1.93 (1)	2.767 (2)	172 (3)
$O2w-H21 \cdots O3^{iv}$	0.83 (1)	2.13 (1)	2.926 (2)	160 (3)
$O2w-H22 \cdots O4^v$	0.84 (1)	1.93 (1)	2.765 (2)	174 (3)

Symmetry codes: (i)  $x - 1, y, z$ ; (ii)  $-x + 1, -y + 1, -z$ ; (iii)  $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (iv)  $x - 1, -y + \frac{3}{2}, z + \frac{1}{2}$ ; (v)  $x, y, z + 1$ .

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Natural Science Foundation of Heilongjiang Province (No. B200501), Heilongjiang University, China, and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2623).

## References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Gao, S., Zhang, Z.-Y., Huo, L.-H., Zhao, H. & Zhao, J.-G. (2004). *Acta Cryst. E* **60**, m1422–m1424.  
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.  
Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.  
Rigaku/MSC (2002). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
Westrip, S. P. (2009). *publCIF*. In preparation.

**supplementary materials**

*Acta Cryst.* (2009). E65, m1310 [ doi:10.1107/S1600536809039774 ]

## Tetraaquabis(2-oxo-1,2-dihydropyridine-5-sulfonato- $\kappa O^2$ )zinc(II)

Z.-B. Zhu, S. Gao and S. W. Ng

### Experimental

Zinc carbonate (0.25 g, 2 mmol) was added to a hot aqueous solution of 2-hydroxypyridine 5-sulfonic acid (0.35 g, 2 mmol); the pH value was adjusted to 6 with 0.1 M sodium hydroxide. The solution was allowed to evaporate slowly. Colorless prismatic crystals were isolated after five days. CH&N elemental analysis. Calc. for  $C_{10}H_{16}N_2O_{12}S_2Zn$ : C 24.73, H 3.32, N 5.77%; found: C 24.77, H 3.37, N 5.81%.

### Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 Å) and were included in the refinement in the riding model approximation, with  $U(H)$  set to  $1.2U(C)$ . The ammonium and water H-atoms were refined with a distance restraint of N—H = O—H  $0.85 \pm 0.01$  Å; their temperature factors were refined.

### Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of  $Zn(H_2O)_4(C_5H_4NO_4S)_2$  at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

## Tetraaquabis(2-oxo-1,2-dihydropyridine-5-sulfonato- $\kappa O^2$ )zinc(II)

### Crystal data

$[Zn(C_5H_4NO_4S)_2(H_2O)_4]$

$M_r = 485.74$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 6.7701$  (2) Å

$b = 13.9725$  (5) Å

$c = 10.0343$  (3) Å

$\beta = 115.331$  (2)°

$V = 857.93$  (5) Å<sup>3</sup>

$Z = 2$

$F_{000} = 496$

$D_x = 1.880$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 7685 reflections

$\theta = 3.3$ – $27.5$ °

$\mu = 1.74$  mm<sup>-1</sup>

$T = 293$  K

Prism, colorless

$0.21 \times 0.16 \times 0.16$  mm

## Data collection

Rigaku R-Axis RAPID IP diffractometer	1951 independent reflections
Radiation source: fine-focus sealed tube	1866 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.021$
$T = 293$ K	$\theta_{\text{max}} = 27.5^\circ$
$\omega$ scans	$\theta_{\text{min}} = 3.3^\circ$
Absorption correction: Multi-scan (ABSCOR; Higashi, 1995)	$h = -8 \rightarrow 8$
$T_{\text{min}} = 0.711$ , $T_{\text{max}} = 0.768$	$k = -17 \rightarrow 18$
8224 measured reflections	$l = -13 \rightarrow 13$

## Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.023$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.066$	$w = 1/[\sigma^2(F_o^2) + (0.037P)^2 + 0.5079P]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
1951 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
144 parameters	$\Delta\rho_{\text{max}} = 0.37 \text{ e } \text{\AA}^{-3}$
5 restraints	$\Delta\rho_{\text{min}} = -0.39 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	0.5000	0.5000	0.02207 (10)
S1	0.81951 (6)	0.69216 (3)	-0.07941 (4)	0.01923 (11)
O1	0.4186 (2)	0.54472 (10)	0.28771 (14)	0.0295 (3)
O3	0.9550 (2)	0.77155 (9)	0.00655 (14)	0.0290 (3)
O2	0.95353 (19)	0.61832 (9)	-0.10431 (13)	0.0264 (3)
O4	0.63570 (19)	0.72213 (9)	-0.21315 (13)	0.0271 (3)
O1W	0.20020 (19)	0.43039 (9)	0.40362 (13)	0.0252 (2)
O2W	0.3496 (2)	0.62596 (11)	0.53828 (17)	0.0406 (3)
C2	0.7484 (3)	0.55821 (12)	0.25390 (18)	0.0234 (3)
H2	0.8360	0.5271	0.3413	0.028*
C1	0.5190 (3)	0.56850 (11)	0.21274 (17)	0.0214 (3)
N1	0.4007 (2)	0.60683 (10)	0.07675 (15)	0.0229 (3)
C5	0.4880 (2)	0.64352 (13)	-0.01119 (17)	0.0221 (3)
H5	0.3979	0.6715	-0.1009	0.027*
C4	0.7072 (2)	0.63937 (11)	0.03179 (17)	0.0201 (3)

C3	0.8395 (3)	0.59340 (11)	0.16668 (18)	0.0227 (3)
H3	0.9891	0.5873	0.1956	0.027*
H1	0.2625 (16)	0.6084 (16)	0.045 (2)	0.035 (6)*
H11	0.158 (4)	0.4218 (18)	0.3125 (12)	0.048 (7)*
H12	0.166 (4)	0.3815 (12)	0.438 (3)	0.045 (7)*
H21	0.223 (2)	0.6479 (18)	0.509 (3)	0.051 (7)*
H22	0.431 (4)	0.6538 (17)	0.6168 (18)	0.049 (7)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.02021 (15)	0.02605 (16)	0.02093 (15)	-0.00103 (9)	0.00973 (11)	0.00145 (9)
S1	0.01744 (18)	0.02194 (19)	0.01890 (19)	-0.00059 (13)	0.00834 (14)	0.00027 (13)
O1	0.0244 (6)	0.0417 (7)	0.0246 (6)	-0.0023 (5)	0.0125 (5)	0.0061 (5)
O3	0.0276 (6)	0.0281 (6)	0.0320 (7)	-0.0085 (5)	0.0133 (5)	-0.0060 (5)
O2	0.0222 (5)	0.0329 (6)	0.0252 (6)	0.0046 (5)	0.0111 (5)	-0.0021 (5)
O4	0.0241 (6)	0.0313 (6)	0.0233 (6)	0.0017 (5)	0.0076 (5)	0.0071 (5)
O1W	0.0249 (6)	0.0286 (6)	0.0223 (6)	-0.0043 (5)	0.0104 (5)	0.0014 (5)
O2W	0.0268 (7)	0.0401 (8)	0.0447 (8)	0.0077 (6)	0.0056 (6)	-0.0145 (6)
C2	0.0214 (7)	0.0268 (8)	0.0198 (7)	0.0036 (6)	0.0069 (6)	0.0030 (6)
C1	0.0223 (7)	0.0218 (7)	0.0204 (7)	-0.0023 (6)	0.0094 (6)	-0.0005 (6)
N1	0.0151 (6)	0.0316 (7)	0.0216 (7)	-0.0009 (5)	0.0074 (5)	0.0019 (5)
C5	0.0203 (7)	0.0264 (8)	0.0190 (7)	0.0007 (6)	0.0079 (6)	0.0024 (5)
C4	0.0198 (7)	0.0215 (7)	0.0204 (7)	-0.0012 (6)	0.0101 (6)	-0.0006 (5)
C3	0.0178 (7)	0.0266 (8)	0.0229 (8)	0.0021 (6)	0.0078 (6)	0.0001 (6)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Zn1—O1 <sup>i</sup>	2.0560 (12)	O2W—H21	0.833 (10)
Zn1—O1	2.0560 (12)	O2W—H22	0.838 (10)
Zn1—O1W <sup>i</sup>	2.0788 (12)	C2—C3	1.360 (2)
Zn1—O1W	2.0788 (12)	C2—C1	1.434 (2)
Zn1—O2W	2.1487 (14)	C2—H2	0.9300
Zn1—O2W <sup>i</sup>	2.1487 (14)	C1—N1	1.362 (2)
S1—O4	1.4477 (12)	N1—C5	1.356 (2)
S1—O3	1.4626 (12)	N1—H1	0.850 (10)
S1—O2	1.4643 (12)	C5—C4	1.358 (2)
S1—C4	1.7588 (15)	C5—H5	0.9300
O1—C1	1.2553 (19)	C4—C3	1.418 (2)
O1W—H11	0.841 (10)	C3—H3	0.9300
O1W—H12	0.841 (10)		
O1 <sup>i</sup> —Zn1—O1	180.0	Zn1—O1W—H12	125.1 (18)
O1 <sup>i</sup> —Zn1—O1W <sup>i</sup>	83.37 (5)	H11—O1W—H12	108 (2)
O1—Zn1—O1W <sup>i</sup>	96.63 (5)	Zn1—O2W—H21	137.0 (19)
O1 <sup>i</sup> —Zn1—O1W	96.63 (5)	Zn1—O2W—H22	112.2 (18)
O1—Zn1—O1W	83.37 (5)	H21—O2W—H22	109 (3)
O1W <sup>i</sup> —Zn1—O1W	180.00 (6)	C3—C2—C1	120.81 (15)

## supplementary materials

O1 <sup>i</sup> —Zn1—O2W	90.07 (6)	C3—C2—H2	119.6
O1—Zn1—O2W	89.93 (6)	C1—C2—H2	119.6
O1W <sup>i</sup> —Zn1—O2W	88.66 (5)	O1—C1—N1	117.88 (14)
O1W—Zn1—O2W	91.34 (5)	O1—C1—C2	126.76 (15)
O1 <sup>i</sup> —Zn1—O2W <sup>i</sup>	89.93 (6)	N1—C1—C2	115.35 (14)
O1—Zn1—O2W <sup>i</sup>	90.07 (6)	C5—N1—C1	124.58 (13)
O1W <sup>i</sup> —Zn1—O2W <sup>i</sup>	91.34 (5)	C5—N1—H1	117.5 (16)
O1W—Zn1—O2W <sup>i</sup>	88.66 (5)	C1—N1—H1	118.0 (16)
O2W—Zn1—O2W <sup>i</sup>	180.0	N1—C5—C4	119.93 (14)
O4—S1—O3	113.63 (8)	N1—C5—H5	120.0
O4—S1—O2	113.28 (7)	C4—C5—H5	120.0
O3—S1—O2	110.91 (7)	C5—C4—C3	118.76 (14)
O4—S1—C4	106.01 (7)	C5—C4—S1	119.40 (12)
O3—S1—C4	106.05 (7)	C3—C4—S1	121.84 (12)
O2—S1—C4	106.28 (7)	C2—C3—C4	120.17 (14)
C1—O1—Zn1	136.67 (11)	C2—C3—H3	119.9
Zn1—O1W—H11	112.6 (17)	C4—C3—H3	119.9
O1W <sup>i</sup> —Zn1—O1—C1	28.28 (18)	N1—C5—C4—C3	2.1 (2)
O1W—Zn1—O1—C1	-151.72 (18)	N1—C5—C4—S1	-177.08 (12)
O2W—Zn1—O1—C1	116.92 (17)	O4—S1—C4—C5	-6.85 (15)
O2W <sup>i</sup> —Zn1—O1—C1	-63.08 (17)	O3—S1—C4—C5	114.25 (14)
Zn1—O1—C1—N1	-171.26 (12)	O2—S1—C4—C5	-127.66 (13)
Zn1—O1—C1—C2	9.9 (3)	O4—S1—C4—C3	173.98 (13)
C3—C2—C1—O1	-175.15 (17)	O3—S1—C4—C3	-64.92 (15)
C3—C2—C1—N1	6.0 (2)	O2—S1—C4—C3	53.17 (15)
O1—C1—N1—C5	173.90 (16)	C1—C2—C3—C4	-1.2 (2)
C2—C1—N1—C5	-7.1 (2)	C5—C4—C3—C2	-3.0 (2)
C1—N1—C5—C4	3.2 (3)	S1—C4—C3—C2	176.21 (13)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ .

### Hydrogen-bond geometry ( $\text{\AA}, ^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 <sup>ii</sup> ...O2 <sup>ii</sup>	0.85 (1)	1.99 (1)	2.790 (2)	157 (2)
O1w—H11 <sup>iii</sup> ...O2 <sup>iii</sup>	0.84 (1)	1.98 (1)	2.809 (2)	171 (2)
O1w—H12 <sup>iv</sup> ...O3 <sup>iv</sup>	0.84 (1)	1.93 (1)	2.767 (2)	172 (3)
O2w—H21 <sup>v</sup> ...O3 <sup>v</sup>	0.83 (1)	2.13 (1)	2.926 (2)	160 (3)
O2w—H22 <sup>vi</sup> ...O4 <sup>vi</sup>	0.84 (1)	1.93 (1)	2.765 (2)	174 (3)

Symmetry codes: (ii)  $x-1, y, z$ ; (iii)  $-x+1, -y+1, -z$ ; (iv)  $-x+1, y-1/2, -z+1/2$ ; (v)  $x-1, -y+3/2, z+1/2$ ; (vi)  $x, y, z+1$ .

Fig. 1

