

Aquabis(4-formylbenzoato- $\kappa^2 O^1, O^{1\prime}$)-bis(isonicotinamide- κN^1)cadmium(II) monohydrate

Tuncer Hökelek,^{a*} Filiz Yılmaz,^b Barış Tercan,^c Ferdi Gürgen^d and Hacali Necefoğlu^d

^aDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey,

^bDepartment of Chemistry, Faculty of Science, Anadolu University, 26470

Yenibağlar, Eskisehir, Turkey, ^cDepartment of Physics, Karabük University, 78050

Karabük, Turkey, and ^dDepartment of Chemistry, Kafkas University, 63100 Kars, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 9 October 2009; accepted 16 October 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.003$ Å; disorder in main residue; R factor = 0.028; wR factor = 0.075; data-to-parameter ratio = 16.6.

The asymmetric unit of the title Cd^{II} complex, $[Cd(C_8H_5O_3)_2(C_6H_6N_2O)_2 \cdot (H_2O)] \cdot H_2O$, contains two 4-formylbenzoate (FB), two isonicotinamide (INA) ligands, one coordinated and one uncoordinated water molecule; the FB ions act as bidentate ligands. The coordination number of the Cd(II) atom is seven within a CdO_5N_2 donor set. Intramolecular O—H···O hydrogen bonds link the uncoordinated water molecules to the carboxyl groups. The dihedral angle between the carboxylate groups and the adjacent benzene rings are 17.53 (13) and 16.55 (14)°. In the crystal structure, intermolecular O—H···O, N—H···O, N—H···N and C—H···O hydrogen bonds link the molecules into a supramolecular structure. The amide group of one of the INA ligands is disordered over two orientations, with an occupancy ratio of 0.759 (3):0.241 (3).

Related literature

For niacin, see: Krishnamachari (1974) and for the nicotinic acid derivative *N,N*-diethylnicotinamide, see: Bigoli *et al.* (1972). For related structures, see: Greenaway *et al.* (1984); Hökelek & Necefoğlu (1996); Hökelek *et al.* (2009).



Experimental

Crystal data

$[Cd(C_8H_5O_3)_2(C_6H_6N_2O)_2 \cdot (H_2O)] \cdot H_2O$

$M_r = 690.93$

Monoclinic, $P2_{1}/c$

$a = 9.3357$ (3) Å

$b = 19.0501$ (6) Å

$c = 16.3743$ (5) Å

$\beta = 93.203$ (1)°

$V = 2907.55$ (16) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.82$ mm⁻¹

$T = 100$ K

$0.27 \times 0.11 \times 0.10$ mm

Data collection

Bruker Kappa APEXII CCD area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2005)

$T_{\min} = 0.896$, $T_{\max} = 0.920$

26549 measured reflections

7151 independent reflections

5904 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$

$wR(F^2) = 0.075$

$S = 1.03$

7151 reflections

431 parameters

8 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 1.16$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.63$ e Å⁻³

Table 1
Selected bond lengths (Å).

Cd1—O1	2.6055 (13)	Cd1—O9	2.3271 (15)
Cd1—O2	2.3066 (13)	Cd1—N1	2.3200 (16)
Cd1—O3	2.3368 (13)	Cd1—N3	2.3362 (16)
Cd1—O4	2.5117 (13)		

Table 2
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N2—H2A···O7 ⁱ	0.86	2.04	2.891 (2)	172
N2—H2B···O10 ⁱⁱ	0.86	2.05	2.888 (3)	165
N4—H4A···O8 ⁱⁱⁱ	0.86	2.01	2.863 (3)	174
N4—H4B···O5 ^{iv}	0.86	2.01	2.852 (3)	166
N4—H4A···N4B ⁱⁱⁱ	0.86	2.35	3.096 (7)	146
N4B—H4B1···N4 ⁱⁱⁱ	0.86	2.35	3.096 (7)	145
N4B—H4B1···O8B ⁱⁱⁱ	0.86	2.02	2.880 (9)	178
N4B—H4B2···O7 ^v	0.86	2.12	2.941 (7)	158
O9—H91···O4 ^{vi}	0.884 (15)	1.86 (2)	2.740 (2)	175 (2)

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O9—H92···O1 ^v	0.875 (16)	1.87 (2)	2.740 (2)	174 (2)
O10—H101···O3	0.880 (16)	2.32 (3)	2.876 (2)	122 (2)
O10—H102···O2	0.864 (18)	1.90 (3)	2.766 (2)	175 (3)
C11—H11···O8 ⁱⁱ	0.93	2.55	3.315 (3)	140
C17—H17···O1 ^v	0.93	2.44	3.212 (2)	140
C20—H20···O10 ⁱⁱ	0.93	2.35	3.258 (3)	166
C23—H23···O4 ^{vi}	0.93	2.48	3.253 (2)	140
C26—H26···O6 ^{vii}	0.93	2.55	3.137 (3)	122
C27—H27···O6 ^{vii}	0.93	2.51	3.133 (3)	124

Symmetry codes: (i) $-x, -y + 1, -z - 1$; (ii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (v) $-x, -y + 1, -z$; (vi) $-x + 1, -y + 1, -z$; (vii) $x - 1, -y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors are indebted to Anadolu University and the Medicinal Plants and Medicine Research Centre of Anadolu

University, Eskişehir, Turkey, for the use of the X-ray diffractometer. This work was supported financially by Kafkas University Research Fund (grant No. 2009-FEF-03).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2635).

References

- Bigoli, F., Braibanti, A., Pellinghelli, M. A. & Tiripicchio, A. (1972). *Acta Cryst.* **B28**, 962–966.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Greenaway, F. T., Pazeshk, A., Cordes, A. W., Noble, M. C. & Sorenson, J. R. J. (1984). *Inorg. Chim. Acta*, **93**, 67–71.
- Hökelek, T., Dal, H., Tercan, B., Aybirdi, Ö. & Necefoglu, H. (2009). *Acta Cryst.* **E65**, m651–m652.
- Hökelek, T. & Necefoglu, H. (1996). *Acta Cryst.* **C52**, 1128–1131.
- Krishnamachari, K. A. V. R. (1974). *Am. J. Clin. Nutr.* **27**, 108–111.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2009). E65, m1416–m1417 [https://doi.org/10.1107/S1600536809042640]

Aquabis(4-formylbenzoato- $\kappa^2 O^1, O^{1\prime}$)bis(isonicotinamide- κN^1)cadmium(II) monohydrate

Tuncer Hökelek, Filiz Yılmaz, Barış Tercan, Ferdi Gürgen and Hacali Necefoğlu

S1. Comment

As a part of our ongoing investigation on transition metal complexes of nicotinamide (NA), one form of niacin (Krishnamachari, 1974), and/or the nicotinic acid derivative *N,N*-diethylnicotinamide (DENA), an important respiratory stimulant (Bigoli *et al.*, 1972), the title compound was synthesized and its crystal structure is reported herein.

In the monomeric title complex, (I), the Cd^{II} ion is surrounded by two formylbenzoate (FB) and two isonicotinamide (INA) ligands and one water molecule. The FB ions act as bidentate ligands, while the INA ions are monodentate ligands. The structures of similar complexes of Zn^{II} ion, [Zn₂(C₁₀H₁₄N₂O)₂(C₇H₅O₃)₄]₂H₂O, (II) (Hökelek & Necefoğlu, 1996) and [Zn(C₉H₁₀NO₂)₂(C₆H₆N₂O)(H₂O)₂], (III) (Hökelek *et al.*, 2009) have also been determined.

In the title compound (Fig. 1), the average Cd—O bond length (Table 1) is 2.4175 (13) Å and the Cd atom is displaced out of the least-squares planes of the carboxylate groups (O1/C1/O2) and (O3/C9/O4) by 0.1556 (2) Å and -0.0577 (2) Å, respectively. The dihedral angle between the planar carboxylate groups and the adjacent benzene rings A (C₂—C₇) and B (C₁₀—C₁₅) are 17.53 (13)° and 16.55 (14)°, respectively, while those between rings A, B, C (N₁/C₁₇—C₂₁) and D (N₃/C₂₃—C₂₇) are A/B = 7.05 (6), A/C = 69.34 (6), A/D = 66.49 (6), B/C = 68.78 (6), B/D = 73.54 (6) and C/D = 87.43 (5) °. The two four-membered rings, (Cd₁/O₁/O₂/C₁) and (Cd₁/O₃/O₄/C₉), are oriented at a dihedral angle of 20.06 (6)°. The intramolecular O—H···O hydrogen bonds (Table 2) link the uncoordinated water molecule to the carboxylate groups. In (I), the O₁—Cd₁—O₂ and O₃—Cd₁—O₄ angles are 52.91 (4) and 53.96 (4) °, respectively. The corresponding O—M—O (where M is a metal) angles are 58.3 (3)° in (II), 60.03 (6)° in (III) and 55.2 (1)° in [Cu(Asp)₂(py)₂] (where Asp is acetylsalicylate and py is pyridine) [(IV); Greenaway *et al.*, 1984].

In the crystal structure, intramolecular O—H···O and intermolecular O—H···O, N—H···O, N—H···N and C—H···O hydrogen bonds (Table 2) link the molecules into a supramolecular structure, in which they may be effective in the stabilization of the structure.

S2. Experimental

The title compound was prepared by the reaction of 3CdSO₄·8H₂O (3.85 g, 5 mmol) in H₂O (25 ml) and INA (1.22 g, 10 mmol) in H₂O (40 ml) with sodium 4-formylbenzoate (1.72 g, 10 mmol) in H₂O (50 ml). The mixture was filtered and set aside to crystallize at ambient temperature for several days, giving colorless single crystals.

S3. Refinement

Atoms H₈ and H₁₆ (for methine) and H₉₁, H₉₂, H₁₀₁ and H₁₀₂ (for H₂O) were located in difference Fourier map and refined isotropically, with restraints of O₉—H₉₁ = 0.894 (15), O₉—H₉₂ = 0.875 (16), O₁₀—H₁₀₁ = 0.880 (16) and O₁₀—H₁₀₂ = 0.864 (18) Å and H₉₁—O₉—H₉₂ = 106 (2) and H₁₀₁—O₁₀—H₁₀₂ = 106 (3) °. The remaining H atoms were positioned geometrically with N—H = 0.86 Å (for NH₂) and C—H = 0.93 Å for aromatic H atoms and constrained to ride

on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$. The O8, N4, H4A and H4B atoms are disordered over two orientations. During the refinement process, the disordered O8, N4, H4A, H4B and O8B, N4B, H4B1, H4B2 were refined with occupancies of 0.759 (3) and 0.241 (3), respectively.




Figure 1

The molecular structure of the title molecule with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Dashed lines indicate the hydrogen-bondings. Hydrogen atoms except of water molecules have been omitted for clarity.

Aquabis(4-formylbenzoato- $\kappa^2\text{O}^1,\text{O}^1'$)bis(isonicotinamide- κN^1)cadmium(II) monohydrate

Crystal data



$M_r = 690.93$

Monoclinic, $P2_1/c$

Hall symbol: -P 2yb_c

$a = 9.3357 (3)$ Å

$b = 19.0501 (6)$ Å

$c = 16.3743 (5)$ Å

$\beta = 93.203 (1)$ °

$V = 2907.55 (16)$ Å³

$Z = 4$

$F(000) = 1400$

$D_x = 1.578$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9959 reflections

$\theta = 2.4\text{--}28.4$ °

$\mu = 0.82$ mm⁻¹

$T = 100$ K

Block, colorless

$0.27 \times 0.11 \times 0.10$ mm

Data collection

Bruker Kappa APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2005)

$T_{\min} = 0.896$, $T_{\max} = 0.920$

26549 measured reflections

7151 independent reflections

5904 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.043$
 $\theta_{\text{max}} = 28.4^\circ$, $\theta_{\text{min}} = 1.6^\circ$
 $h = -12 \rightarrow 12$

$k = -20 \rightarrow 25$
 $l = -21 \rightarrow 17$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.075$
 $S = 1.03$
7151 reflections
431 parameters
8 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[c^2(F_o^2) + (0.0355P)^2 + 1.7245P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.002$
 $\Delta\rho_{\text{max}} = 1.16 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.63 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cd1	0.254193 (11)	0.604747 (6)	-0.000741 (7)	0.01128 (5)	
N1	0.16048 (16)	0.59088 (8)	-0.13399 (10)	0.0149 (3)	
N2	0.1001 (2)	0.56817 (10)	-0.44328 (10)	0.0292 (4)	
H2A	0.0791	0.5559	-0.4930	0.035*	
H2B	0.1589	0.6022	-0.4330	0.035*	
N3	0.34349 (16)	0.58841 (8)	0.13406 (9)	0.0147 (3)	
N4	0.4990 (3)	0.46566 (12)	0.39673 (14)	0.0269 (6)	0.759 (3)
H4A	0.5191	0.4518	0.4460	0.032*	0.759 (3)
H4B	0.5033	0.4366	0.3567	0.032*	0.759 (3)
N4B	0.3548 (7)	0.5230 (4)	0.4293 (4)	0.0264 (19)	0.241 (3)
H4B1	0.3709	0.5103	0.4794	0.032*	0.241 (3)
H4B2	0.2682	0.5300	0.4103	0.032*	0.241 (3)
O1	-0.00985 (14)	0.59254 (7)	0.04111 (9)	0.0179 (3)	
O2	0.10880 (13)	0.69266 (7)	0.04495 (8)	0.0184 (3)	
O3	0.39326 (13)	0.69982 (7)	-0.03952 (8)	0.0182 (3)	
O4	0.51129 (14)	0.59950 (7)	-0.03809 (9)	0.0171 (3)	
O5	-0.48043 (16)	0.85486 (9)	0.22050 (10)	0.0368 (4)	
O6	1.08430 (17)	0.78213 (9)	-0.20603 (11)	0.0406 (4)	
O7	-0.04317 (19)	0.48524 (9)	-0.39285 (9)	0.0374 (4)	
O8	0.4524 (2)	0.57695 (10)	0.43632 (11)	0.0255 (5)	0.759 (3)
O8B	0.5920 (5)	0.5229 (3)	0.4041 (3)	0.0222 (15)	0.241 (3)

O9	0.25468 (13)	0.48260 (8)	-0.00288 (8)	0.0204 (3)
H91	0.327 (2)	0.4540 (11)	0.0103 (16)	0.037 (7)*
H92	0.179 (2)	0.4560 (12)	-0.0130 (17)	0.042 (8)*
O10	0.28361 (18)	0.80970 (9)	0.06005 (11)	0.0374 (4)
H101	0.336 (3)	0.8070 (17)	0.0170 (15)	0.075 (11)*
H102	0.226 (3)	0.7741 (15)	0.058 (2)	0.100 (14)*
C1	-0.00109 (18)	0.65688 (10)	0.05789 (11)	0.0152 (4)
C2	-0.12241 (18)	0.69296 (10)	0.09768 (11)	0.0167 (4)
C3	-0.22577 (19)	0.65346 (11)	0.13505 (12)	0.0215 (4)
H3	-0.2209	0.6047	0.1349	0.026*
C4	-0.3363 (2)	0.68729 (12)	0.17261 (13)	0.0262 (5)
H4	-0.4053	0.6611	0.1979	0.031*
C5	-0.3441 (2)	0.75969 (12)	0.17262 (13)	0.0255 (5)
C6	-0.2407 (2)	0.79938 (12)	0.13549 (13)	0.0258 (5)
H6	-0.2464	0.8481	0.1354	0.031*
C7	-0.1293 (2)	0.76589 (11)	0.09865 (13)	0.0222 (4)
H7	-0.0591	0.7922	0.0746	0.027*
C8	-0.4633 (2)	0.79273 (13)	0.21423 (15)	0.0323 (5)
H8	-0.539 (3)	0.7581 (15)	0.2423 (17)	0.053 (8)*
C9	0.50454 (18)	0.66490 (10)	-0.05055 (11)	0.0142 (3)
C10	0.63231 (18)	0.70288 (10)	-0.08114 (11)	0.0158 (4)
C11	0.6381 (2)	0.77562 (10)	-0.07766 (13)	0.0216 (4)
H11	0.5634	0.8007	-0.0563	0.026*
C12	0.7556 (2)	0.81089 (11)	-0.10619 (13)	0.0257 (4)
H12	0.7610	0.8595	-0.1025	0.031*
C13	0.8650 (2)	0.77348 (11)	-0.14008 (13)	0.0228 (4)
C14	0.85761 (19)	0.70082 (11)	-0.14571 (13)	0.0215 (4)
H14	0.9300	0.6759	-0.1697	0.026*
C15	0.74230 (19)	0.66558 (10)	-0.11555 (12)	0.0186 (4)
H15	0.7382	0.6169	-0.1182	0.022*
C16	0.9887 (2)	0.81047 (13)	-0.17276 (16)	0.0330 (5)
H16	0.999 (3)	0.8554 (15)	-0.1612 (17)	0.049 (8)*
C17	0.05844 (19)	0.54301 (10)	-0.15291 (12)	0.0179 (4)
H17	0.0135	0.5210	-0.1106	0.021*
C18	0.0173 (2)	0.52511 (10)	-0.23229 (12)	0.0200 (4)
H18	-0.0532	0.4914	-0.2431	0.024*
C19	0.0830 (2)	0.55820 (10)	-0.29624 (11)	0.0180 (4)
C20	0.1849 (2)	0.60930 (10)	-0.27711 (12)	0.0190 (4)
H20	0.2289	0.6334	-0.3183	0.023*
C21	0.22006 (19)	0.62389 (10)	-0.19570 (11)	0.0168 (4)
H21	0.2885	0.6583	-0.1833	0.020*
C22	0.0421 (2)	0.53467 (11)	-0.38244 (12)	0.0250 (4)
C23	0.44196 (19)	0.53843 (10)	0.15065 (11)	0.0168 (4)
H23	0.4839	0.5165	0.1072	0.020*
C24	0.4839 (2)	0.51813 (10)	0.22937 (12)	0.0188 (4)
H24	0.5516	0.4828	0.2385	0.023*
C25	0.42334 (19)	0.55129 (10)	0.29467 (11)	0.0162 (4)
C26	0.3257 (2)	0.60491 (10)	0.27811 (12)	0.0179 (4)

H26	0.2861	0.6293	0.3205	0.021*
C27	0.28801 (19)	0.62164 (10)	0.19738 (11)	0.0165 (4)
H27	0.2216	0.6573	0.1866	0.020*
C28	0.4613 (2)	0.53171 (10)	0.38237 (11)	0.0196 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.01149 (8)	0.01184 (8)	0.01049 (8)	0.00038 (4)	0.00042 (5)	-0.00060 (5)
N1	0.0142 (7)	0.0165 (8)	0.0139 (8)	0.0007 (6)	0.0005 (6)	0.0003 (6)
N2	0.0474 (11)	0.0282 (10)	0.0115 (8)	-0.0121 (8)	-0.0024 (7)	-0.0003 (7)
N3	0.0147 (7)	0.0163 (8)	0.0130 (7)	0.0004 (6)	0.0006 (6)	-0.0012 (6)
N4	0.0490 (15)	0.0203 (12)	0.0109 (11)	0.0046 (10)	-0.0034 (10)	0.0018 (9)
N4B	0.020 (3)	0.048 (5)	0.011 (3)	-0.001 (3)	0.001 (3)	0.007 (3)
O1	0.0174 (6)	0.0163 (7)	0.0203 (7)	0.0021 (5)	0.0029 (5)	-0.0009 (5)
O2	0.0142 (6)	0.0177 (7)	0.0235 (7)	0.0011 (5)	0.0044 (5)	-0.0001 (6)
O3	0.0124 (6)	0.0183 (7)	0.0241 (7)	0.0004 (5)	0.0038 (5)	-0.0024 (6)
O4	0.0141 (6)	0.0162 (7)	0.0212 (7)	-0.0011 (5)	0.0035 (5)	-0.0003 (5)
O5	0.0259 (8)	0.0447 (11)	0.0394 (10)	0.0106 (7)	-0.0015 (7)	-0.0184 (8)
O6	0.0297 (8)	0.0375 (10)	0.0567 (11)	-0.0091 (7)	0.0201 (8)	0.0063 (8)
O7	0.0580 (11)	0.0372 (10)	0.0165 (8)	-0.0249 (8)	-0.0030 (7)	-0.0033 (7)
O8	0.0446 (12)	0.0181 (10)	0.0132 (9)	-0.0015 (8)	-0.0033 (8)	-0.0001 (7)
O8B	0.015 (3)	0.034 (4)	0.018 (3)	-0.002 (2)	0.000 (2)	0.006 (3)
O9	0.0179 (7)	0.0133 (7)	0.0293 (8)	0.0001 (5)	-0.0048 (6)	-0.0007 (6)
O10	0.0360 (9)	0.0361 (10)	0.0407 (10)	-0.0002 (8)	0.0079 (8)	-0.0205 (8)
C1	0.0140 (8)	0.0186 (9)	0.0128 (9)	0.0027 (7)	0.0003 (6)	0.0009 (7)
C2	0.0121 (8)	0.0210 (10)	0.0168 (9)	0.0015 (7)	-0.0004 (7)	-0.0032 (8)
C3	0.0188 (9)	0.0216 (10)	0.0245 (10)	-0.0022 (7)	0.0038 (8)	-0.0072 (8)
C4	0.0173 (9)	0.0334 (12)	0.0284 (11)	-0.0064 (8)	0.0063 (8)	-0.0119 (9)
C5	0.0141 (8)	0.0359 (12)	0.0264 (11)	0.0018 (8)	0.0000 (8)	-0.0150 (9)
C6	0.0208 (9)	0.0243 (11)	0.0322 (12)	0.0058 (8)	0.0002 (8)	-0.0085 (9)
C7	0.0173 (9)	0.0220 (10)	0.0275 (11)	0.0023 (7)	0.0022 (8)	-0.0020 (9)
C8	0.0186 (10)	0.0434 (14)	0.0350 (13)	0.0030 (9)	0.0016 (9)	-0.0222 (11)
C9	0.0131 (8)	0.0168 (9)	0.0127 (8)	-0.0013 (7)	0.0008 (6)	-0.0012 (7)
C10	0.0133 (8)	0.0175 (9)	0.0166 (9)	-0.0016 (7)	0.0004 (7)	0.0000 (7)
C11	0.0195 (9)	0.0186 (10)	0.0272 (11)	0.0000 (7)	0.0056 (8)	-0.0009 (8)
C12	0.0269 (10)	0.0175 (10)	0.0330 (12)	-0.0065 (8)	0.0047 (9)	-0.0012 (9)
C13	0.0189 (9)	0.0234 (10)	0.0263 (11)	-0.0038 (8)	0.0034 (8)	0.0052 (9)
C14	0.0159 (8)	0.0223 (10)	0.0269 (11)	0.0015 (7)	0.0055 (8)	0.0054 (8)
C15	0.0159 (8)	0.0178 (9)	0.0223 (10)	0.0005 (7)	0.0028 (7)	0.0032 (8)
C16	0.0291 (11)	0.0270 (12)	0.0439 (14)	-0.0085 (9)	0.0103 (10)	0.0046 (11)
C17	0.0200 (9)	0.0180 (9)	0.0156 (9)	-0.0029 (7)	0.0008 (7)	0.0023 (7)
C18	0.0252 (9)	0.0176 (10)	0.0168 (10)	-0.0064 (7)	-0.0022 (7)	0.0011 (8)
C19	0.0232 (9)	0.0173 (9)	0.0131 (9)	-0.0002 (7)	-0.0011 (7)	-0.0004 (7)
C20	0.0199 (9)	0.0226 (10)	0.0146 (9)	-0.0023 (7)	0.0015 (7)	0.0018 (8)
C21	0.0162 (8)	0.0197 (9)	0.0146 (9)	-0.0040 (7)	0.0006 (7)	-0.0004 (7)
C22	0.0354 (11)	0.0243 (11)	0.0148 (10)	-0.0056 (9)	-0.0033 (8)	-0.0001 (8)
C23	0.0188 (8)	0.0169 (9)	0.0147 (9)	0.0023 (7)	0.0008 (7)	-0.0023 (7)

C24	0.0210 (9)	0.0175 (9)	0.0176 (9)	0.0042 (7)	-0.0010 (7)	0.0013 (8)
C25	0.0179 (8)	0.0172 (9)	0.0133 (9)	-0.0029 (7)	-0.0009 (7)	0.0023 (7)
C26	0.0180 (9)	0.0217 (10)	0.0141 (9)	0.0015 (7)	0.0017 (7)	-0.0013 (7)
C27	0.0161 (8)	0.0191 (9)	0.0143 (9)	0.0022 (7)	0.0011 (7)	-0.0008 (7)
C28	0.0248 (9)	0.0204 (10)	0.0135 (9)	-0.0014 (8)	-0.0014 (7)	0.0031 (8)

Geometric parameters (\AA , $^{\circ}$)

Cd1—O1	2.6055 (13)	C5—C6	1.392 (3)
Cd1—O2	2.3066 (13)	C5—C8	1.478 (3)
Cd1—O3	2.3368 (13)	C6—C7	1.387 (3)
Cd1—O4	2.5117 (13)	C6—H6	0.9300
Cd1—O9	2.3271 (15)	C7—H7	0.9300
Cd1—N1	2.3200 (16)	C8—H8	1.08 (3)
Cd1—N3	2.3362 (16)	C9—C10	1.505 (2)
N1—C17	1.342 (2)	C10—C11	1.388 (3)
N1—C21	1.337 (2)	C10—C15	1.393 (2)
N2—C22	1.324 (3)	C11—C12	1.390 (3)
N2—H2A	0.8600	C11—H11	0.9300
N2—H2B	0.8600	C12—C13	1.386 (3)
N3—C23	1.341 (2)	C12—H12	0.9300
N3—C27	1.343 (2)	C13—C14	1.389 (3)
N4—H4A	0.8600	C13—C16	1.479 (3)
N4—H4B	0.8600	C14—C15	1.383 (2)
N4B—C28	1.301 (7)	C14—H14	0.9300
N4B—H4B1	0.8600	C15—H15	0.9300
N4B—H4B2	0.8600	C16—H16	0.88 (3)
O1—C1	1.258 (2)	C17—C18	1.378 (3)
O2—C1	1.259 (2)	C17—H17	0.9300
O3—C9	1.255 (2)	C18—C19	1.393 (3)
O4—C9	1.263 (2)	C18—H18	0.9300
O5—C8	1.200 (3)	C19—C20	1.385 (3)
O6—C16	1.199 (3)	C19—C22	1.509 (3)
O7—C22	1.239 (3)	C20—C21	1.383 (3)
O8B—C28	1.263 (5)	C20—H20	0.9300
O9—H91	0.884 (15)	C21—H21	0.9300
O9—H92	0.875 (16)	C23—C24	1.381 (3)
O10—H101	0.880 (16)	C23—H23	0.9300
O10—H102	0.864 (18)	C24—C25	1.389 (3)
C1—C2	1.504 (2)	C24—H24	0.9300
C2—C7	1.391 (3)	C25—C26	1.386 (3)
C2—C3	1.392 (3)	C25—C28	1.507 (3)
C3—C4	1.389 (3)	C26—C27	1.386 (3)
C3—H3	0.9300	C26—H26	0.9300
C4—C5	1.381 (3)	C27—H27	0.9300
C4—H4	0.9300		
O2—Cd1—N1	100.95 (5)	O5—C8—C5	124.6 (2)

O2—Cd1—O9	137.10 (4)	O5—C8—H8	118.1 (15)
N1—Cd1—O9	82.70 (5)	C5—C8—H8	117.3 (15)
O2—Cd1—N3	88.61 (5)	O3—C9—O4	122.24 (16)
N1—Cd1—N3	165.76 (6)	O3—C9—C10	118.10 (16)
O9—Cd1—N3	83.11 (5)	O4—C9—C10	119.65 (15)
O2—Cd1—O3	82.61 (5)	C11—C10—C15	119.88 (17)
N1—Cd1—O3	91.08 (5)	C11—C10—C9	119.74 (16)
O9—Cd1—O3	140.29 (4)	C15—C10—C9	120.34 (16)
N3—Cd1—O3	100.70 (5)	C10—C11—C12	119.89 (18)
O2—Cd1—O4	134.07 (4)	C10—C11—H11	120.1
N1—Cd1—O4	94.69 (5)	C12—C11—H11	120.1
O9—Cd1—O4	87.35 (4)	C13—C12—C11	119.94 (19)
N3—Cd1—O4	85.95 (5)	C13—C12—H12	120.0
O3—Cd1—O4	53.96 (4)	C11—C12—H12	120.0
O2—Cd1—O1	52.91 (4)	C12—C13—C14	120.29 (18)
N1—Cd1—O1	85.68 (5)	C12—C13—C16	120.5 (2)
O9—Cd1—O1	85.26 (4)	C14—C13—C16	119.19 (19)
N3—Cd1—O1	91.86 (5)	C15—C14—C13	119.78 (18)
O3—Cd1—O1	133.54 (4)	C15—C14—H14	120.1
O4—Cd1—O1	172.50 (4)	C13—C14—H14	120.1
C21—N1—C17	117.70 (16)	C14—C15—C10	120.18 (18)
C21—N1—Cd1	120.40 (12)	C14—C15—H15	119.9
C17—N1—Cd1	121.32 (12)	C10—C15—H15	119.9
C22—N2—H2A	120.0	O6—C16—C13	124.5 (2)
C22—N2—H2B	120.0	O6—C16—H16	117.5 (18)
H2A—N2—H2B	120.0	C13—C16—H16	117.5 (18)
C23—N3—C27	117.90 (16)	N1—C17—C18	122.86 (17)
C23—N3—Cd1	119.37 (12)	N1—C17—H17	118.6
C27—N3—Cd1	122.27 (12)	C18—C17—H17	118.6
H4A—N4—H4B	120.0	C17—C18—C19	119.09 (17)
C28—N4B—H4B1	120.0	C17—C18—H18	120.5
C28—N4B—H4B2	120.0	C19—C18—H18	120.5
H4B1—N4B—H4B2	120.0	C20—C19—C18	118.26 (17)
C1—O1—Cd1	85.39 (10)	C20—C19—C22	123.69 (17)
C1—O2—Cd1	99.28 (11)	C18—C19—C22	118.01 (17)
C9—O3—Cd1	96.04 (11)	C21—C20—C19	118.83 (17)
C9—O4—Cd1	87.73 (10)	C21—C20—H20	120.6
Cd1—O9—H91	128.0 (16)	C19—C20—H20	120.6
Cd1—O9—H92	125.4 (16)	N1—C21—C20	123.19 (17)
H91—O9—H92	106 (2)	N1—C21—H21	118.4
H101—O10—H102	106 (3)	C20—C21—H21	118.4
O1—C1—O2	122.29 (16)	O7—C22—N2	123.34 (19)
O1—C1—C2	119.91 (16)	O7—C22—C19	118.77 (18)
O2—C1—C2	117.76 (16)	N2—C22—C19	117.89 (18)
C7—C2—C3	120.09 (17)	N3—C23—C24	122.90 (17)
C7—C2—C1	119.85 (17)	N3—C23—H23	118.5
C3—C2—C1	120.04 (17)	C24—C23—H23	118.5
C4—C3—C2	119.61 (19)	C23—C24—C25	119.00 (17)

C4—C3—H3	120.2	C23—C24—H24	120.5
C2—C3—H3	120.2	C25—C24—H24	120.5
C5—C4—C3	120.26 (19)	C26—C25—C24	118.43 (17)
C5—C4—H4	119.9	C26—C25—C28	119.09 (17)
C3—C4—H4	119.9	C24—C25—C28	122.48 (17)
C4—C5—C6	120.29 (18)	C25—C26—C27	119.04 (17)
C4—C5—C8	117.8 (2)	C25—C26—H26	120.5
C6—C5—C8	121.9 (2)	C27—C26—H26	120.5
C7—C6—C5	119.7 (2)	N3—C27—C26	122.65 (17)
C7—C6—H6	120.2	N3—C27—H27	118.7
C5—C6—H6	120.2	C26—C27—H27	118.7
C6—C7—C2	120.04 (19)	O8B—C28—N4B	125.1 (4)
C6—C7—H7	120.0	O8B—C28—C25	118.3 (3)
C2—C7—H7	120.0	N4B—C28—C25	116.6 (3)
O2—Cd1—N1—C21	95.74 (14)	C3—C4—C5—C6	-0.5 (3)
O9—Cd1—N1—C21	-127.58 (14)	C3—C4—C5—C8	-179.42 (19)
N3—Cd1—N1—C21	-132.88 (19)	C4—C5—C6—C7	-0.2 (3)
O3—Cd1—N1—C21	13.06 (14)	C8—C5—C6—C7	178.7 (2)
O4—Cd1—N1—C21	-40.86 (14)	C5—C6—C7—C2	1.1 (3)
O1—Cd1—N1—C21	146.65 (14)	C3—C2—C7—C6	-1.3 (3)
O2—Cd1—N1—C17	-93.18 (14)	C1—C2—C7—C6	-179.81 (18)
O9—Cd1—N1—C17	43.49 (14)	C4—C5—C8—O5	178.0 (2)
N3—Cd1—N1—C17	38.2 (3)	C6—C5—C8—O5	-0.9 (4)
O3—Cd1—N1—C17	-175.86 (14)	Cd1—O3—C9—O4	-1.43 (19)
O4—Cd1—N1—C17	130.21 (14)	Cd1—O3—C9—C10	177.44 (14)
O1—Cd1—N1—C17	-42.27 (14)	Cd1—O4—C9—O3	1.33 (18)
O2—Cd1—N3—C23	179.10 (14)	Cd1—O4—C9—C10	-177.53 (15)
N1—Cd1—N3—C23	46.6 (2)	O3—C9—C10—C11	15.4 (3)
O9—Cd1—N3—C23	41.28 (13)	O4—C9—C10—C11	-165.66 (18)
O3—Cd1—N3—C23	-98.69 (13)	O3—C9—C10—C15	-162.27 (18)
O4—Cd1—N3—C23	-46.53 (13)	O4—C9—C10—C15	16.6 (3)
O1—Cd1—N3—C23	126.28 (13)	C15—C10—C11—C12	-2.0 (3)
O2—Cd1—N3—C27	7.06 (14)	C9—C10—C11—C12	-179.72 (18)
N1—Cd1—N3—C27	-125.5 (2)	C10—C11—C12—C13	1.8 (3)
O9—Cd1—N3—C27	-130.76 (14)	C11—C12—C13—C14	0.0 (3)
O3—Cd1—N3—C27	89.27 (14)	C11—C12—C13—C16	178.3 (2)
O4—Cd1—N3—C27	141.42 (14)	C12—C13—C14—C15	-1.6 (3)
O1—Cd1—N3—C27	-45.76 (14)	C16—C13—C14—C15	-179.9 (2)
O2—Cd1—O1—C1	-1.99 (10)	C13—C14—C15—C10	1.3 (3)
N1—Cd1—O1—C1	-109.20 (11)	C11—C10—C15—C14	0.5 (3)
O9—Cd1—O1—C1	167.79 (11)	C9—C10—C15—C14	178.16 (18)
N3—Cd1—O1—C1	84.86 (11)	C12—C13—C16—O6	-176.7 (2)
O3—Cd1—O1—C1	-21.82 (13)	C14—C13—C16—O6	1.6 (4)
N1—Cd1—O2—C1	77.97 (12)	C21—N1—C17—C18	2.4 (3)
O9—Cd1—O2—C1	-13.06 (14)	Cd1—N1—C17—C18	-168.89 (15)
N3—Cd1—O2—C1	-91.39 (11)	N1—C17—C18—C19	-0.5 (3)
O3—Cd1—O2—C1	167.64 (12)	C17—C18—C19—C20	-1.6 (3)

O4—Cd1—O2—C1	−174.38 (10)	C17—C18—C19—C22	176.54 (18)
O1—Cd1—O2—C1	2.00 (10)	C18—C19—C20—C21	1.8 (3)
O2—Cd1—O3—C9	164.84 (12)	C22—C19—C20—C21	−176.23 (19)
N1—Cd1—O3—C9	−94.27 (11)	C17—N1—C21—C20	−2.2 (3)
O9—Cd1—O3—C9	−14.42 (14)	Cd1—N1—C21—C20	169.18 (15)
N3—Cd1—O3—C9	77.67 (11)	C19—C20—C21—N1	0.1 (3)
O4—Cd1—O3—C9	0.75 (10)	C20—C19—C22—O7	175.1 (2)
O1—Cd1—O3—C9	−179.32 (10)	C18—C19—C22—O7	−2.9 (3)
O2—Cd1—O4—C9	−22.99 (13)	C20—C19—C22—N2	−4.7 (3)
N1—Cd1—O4—C9	87.17 (11)	C18—C19—C22—N2	177.2 (2)
O9—Cd1—O4—C9	169.62 (11)	C27—N3—C23—C24	2.6 (3)
N3—Cd1—O4—C9	−107.10 (11)	Cd1—N3—C23—C24	−169.79 (14)
O3—Cd1—O4—C9	−0.74 (10)	N3—C23—C24—C25	−0.9 (3)
Cd1—O1—C1—O2	3.43 (17)	C23—C24—C25—C26	−1.7 (3)
Cd1—O1—C1—C2	−174.23 (16)	C23—C24—C25—C28	179.10 (17)
Cd1—O2—C1—O1	−3.9 (2)	C24—C25—C26—C27	2.4 (3)
Cd1—O2—C1—C2	173.79 (13)	C28—C25—C26—C27	−178.32 (17)
O1—C1—C2—C7	−164.72 (18)	C23—N3—C27—C26	−1.8 (3)
O2—C1—C2—C7	17.5 (3)	Cd1—N3—C27—C26	170.37 (14)
O1—C1—C2—C3	16.7 (3)	C25—C26—C27—N3	−0.7 (3)
O2—C1—C2—C3	−161.05 (18)	C26—C25—C28—O8B	−134.3 (4)
C7—C2—C3—C4	0.6 (3)	C24—C25—C28—O8B	44.9 (4)
C1—C2—C3—C4	179.12 (18)	C26—C25—C28—N4B	47.9 (5)
C2—C3—C4—C5	0.3 (3)	C24—C25—C28—N4B	−132.9 (4)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N2—H2A···O7 ⁱ	0.86	2.04	2.891 (2)	172
N2—H2B···O10 ⁱⁱ	0.86	2.05	2.888 (3)	165
N4—H4A···O8 ⁱⁱⁱ	0.86	2.01	2.863 (3)	174
N4—H4B···O5 ^{iv}	0.86	2.01	2.852 (3)	166
N4—H4A···N4B ⁱⁱⁱ	0.86	2.35	3.096 (7)	146
N4B—H4B1···N4 ⁱⁱⁱ	0.86	2.35	3.096 (7)	145
N4B—H4B1···O8B ⁱⁱⁱ	0.86	2.02	2.880 (9)	178
N4B—H4B2···O7 ^v	0.86	2.12	2.941 (7)	158
O9—H91···O4 ^{vi}	0.88 (2)	1.86 (2)	2.740 (2)	175 (2)
O9—H92···O1 ^v	0.88 (2)	1.87 (2)	2.740 (2)	174 (2)
O10—H101···O3	0.88 (2)	2.32 (3)	2.876 (2)	122 (2)
O10—H102···O2	0.86 (2)	1.90 (3)	2.766 (2)	175 (3)
C11—H11···O8 ⁱⁱ	0.93	2.55	3.315 (3)	140
C17—H17···O1 ^v	0.93	2.44	3.212 (2)	140
C20—H20···O10 ⁱⁱ	0.93	2.35	3.258 (3)	166
C23—H23···O4 ^{vi}	0.93	2.48	3.253 (2)	140

C26—H26···O6 ^{vii}	0.93	2.55	3.137 (3)	122
C27—H27···O6 ^{vii}	0.93	2.51	3.133 (3)	124

Symmetry codes: (i) $-x, -y+1, -z-1$; (ii) $x, -y+3/2, z-1/2$; (iii) $-x+1, -y+1, -z+1$; (iv) $-x, y-1/2, -z+1/2$; (v) $-x, -y+1, -z$; (vi) $-x+1, -y+1, -z$; (vii) $x-1, -y+3/2, z+1/2$.