

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Hydroxy-*N'*-(5-hydroxy-2-nitrobenzylidene)-2-naphthohydrazide

De-Suo Yang

Department of Chemistry and Chemical Engineering, Baoji University of Arts and Sciences, Baoji 721007, People's Republic of China
Correspondence e-mail: desuoyang@yahoo.com.cn


Received 21 October 2009; accepted 29 October 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.044; wR factor = 0.119; data-to-parameter ratio = 14.2.

The molecule of the title compound, $\text{C}_{18}\text{H}_{13}\text{N}_3\text{O}_5$, displays an *E* configuration with respect to the $\text{C}=\text{N}$ double bond. The dihedral angle between the benzene ring and the naphthyl system is $1.1(2)^\circ$. In the crystal structure, molecules are linked through intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a three-dimensional network.

Related literature

For the biological and structural chemistry of hydrazone compounds, see: Avaji *et al.* (2009); Charkoudian *et al.* (2007); Cukurovali *et al.* (2006). For related structures, see: Yang (2008*a,b,c,d,e*, 2007*a,b,c*); Yang & Guo (2006).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{13}\text{N}_3\text{O}_5$
 $M_r = 351.31$
Monoclinic, $P2_1/n$
 $a = 10.1588(3)$ Å
 $b = 8.2562(2)$ Å

$c = 19.5268(5)$ Å
 $\beta = 104.867(1)^\circ$
 $V = 1582.95(7)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.11$ mm⁻¹
 $T = 298$ K

0.23 × 0.20 × 0.20 mm

Data collection

Bruker SMART CCD diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.975$, $T_{\max} = 0.978$

9168 measured reflections
3425 independent reflections
2436 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.119$
 $S = 1.04$
3425 reflections
241 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.19$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> ⋯ <i>A</i>	<i>D</i> —H⋯ <i>A</i>
O2—H2 \cdots O1	0.82	1.88	2.5987 (18)	146
N1—H1 \cdots O4 ⁱ	0.898 (9)	2.144 (10)	3.0361 (19)	172.8 (18)
O5—H5 \cdots O1 ⁱⁱ	0.82	1.88	2.6889 (17)	168

Symmetry codes: (i) $-x + 2, -y + 1, -z$; (ii) $-x + \frac{5}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The author acknowledges Baoji University of Arts and Sciences for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2256).

References

- Avaji, P. G., Vinod Kumar, C. H., Patil, S. A., Shivananda, K. N. & Nagaraju, C. (2009). *Eur. J. Med. Chem.* **44**, 3552–3559.
Bruker (2002). SAINT and SMART. Bruker AXS Inc., Madison, Wisconsin, USA.
Charkoudian, L. K., Pham, D. M., Kwon, A. M., Vangeloff, A. D. & Franz, K. J. (2007). *J. Chem. Soc. Dalton Trans.* pp. 5031–5042.
Cukurovali, A., Yilmaz, I., Gur, S. & Kazaz, C. (2006). *Eur. J. Med. Chem.* **41**, 201–207.
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Yang, D.-S. (2007*a*). *J. Chem. Crystallogr.* **37**, 343–348.
Yang, D.-S. (2007*b*). *Acta Cryst.* **E63**, o3738.
Yang, D.-S. (2007*c*). *Acta Cryst.* **E63**, o3739.
Yang, D.-S. (2008*a*). *Acta Cryst.* **E64**, o1758.
Yang, D.-S. (2008*b*). *Acta Cryst.* **E64**, o1759.
Yang, D.-S. (2008*c*). *Acta Cryst.* **E64**, o1849.
Yang, D.-S. (2008*d*). *Acta Cryst.* **E64**, o1850.
Yang, D.-S. & Guo, J.-B. (2006). *Acta Cryst.* **E62**, o4414–o4415.

supplementary materials

Acta Cryst. (2009). E65, o2978 [doi:10.1107/S1600536809045279]

3-Hydroxy-*N'*-(5-hydroxy-2-nitrobenzylidene)-2-naphthohydrazide

D.-S. Yang

Comment

Hydrazone compounds have been of great interest for their versatile biological and structural chemistry (Avaji *et al.*, 2009; Charkoudian *et al.*, 2007; Cukurovali *et al.*, 2006). Recently, we have reported a few hydrazone compounds (Yang, 2008*a,b,c,d,e*, 2007*a,b,c*; Yang & Guo, 2006). As a further investigation of this work, the crystal structure of the title new hydrazone compound is reported.

In the title compound (Fig. 1), the molecule displays an *E* configuration with respect to the C=N double bond. The C13...C18 benzene ring forms dihedral angles of 28.6 (2) and 1.1 (2)°, respectively, with the O3—N3—O4 nitro group and the C1...C10 naphthyl ring. All bond lengths are within normal ranges. The C12=N2 bond length of 1.266 (2) Å, conforms to the value for a formal double bond. The bond length of 1.341 (2) Å between atoms C11 and N1 is intermediate between a C—N single bond and a C=N double bond, because of conjugation effects in the molecule.

In the crystal structure, molecules are linked through intermolecular N—H...O and O—H...O hydrogen bonds (Table 1), forming a three-dimensional network (Fig. 2).

Experimental

5-Hydroxy-2-nitrobenzaldehyde (0.1 mmol, 16.7 mg) and 3-hydroxy-2-naphthohydrazide (0.1 mmol, 20.2 mg) were dissolved in CHCl₃ (10 ml). The mixture was stirred at room temperature to give a clear colorless solution. Crystals of the title compound were formed by gradual evaporation of the solvent over a period of 3 days, at room temperature.

Refinement

Atom H1 was located in a difference map and refined isotropically, with N—H distance restrained to 0.90 (1) Å. Other H atoms were placed in idealized positions and constrained to ride on their parent atoms, with O—H distances of 0.82 Å, C—H distances of 0.93 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The structure of the title compound showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. Hydrogen bond is shown as a dashed line.




Fig. 2. Molecular packing as viewed along the *b* axis. Hydrogen bonds are shown as dashed lines.

3-Hydroxy-*N'*-(5-hydroxy-2-nitrobenzylidene)-2-naphthohydrazide

Crystal data

$C_{18}H_{13}N_3O_5$	$F_{000} = 728$
$M_r = 351.31$	$D_x = 1.474 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 2463 reflections
$a = 10.1588 (3) \text{ \AA}$	$\theta = 2.5\text{--}26.6^\circ$
$b = 8.2562 (2) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$c = 19.5268 (5) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 104.8670 (10)^\circ$	Block, colorless
$V = 1582.95 (7) \text{ \AA}^3$	$0.23 \times 0.20 \times 0.20 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART CCD diffractometer	3425 independent reflections
Radiation source: fine-focus sealed tube	2436 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.028$
$T = 298 \text{ K}$	$\theta_{\text{max}} = 27.0^\circ$
ω scans	$\theta_{\text{min}} = 2.1^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -12 \rightarrow 12$
$T_{\text{min}} = 0.975$, $T_{\text{max}} = 0.978$	$k = -10 \rightarrow 10$
9168 measured reflections	$l = -24 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.119$	$w = 1/[\sigma^2(F_o^2) + (0.0503P)^2 + 0.3232P]$
	where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.04$ $(\Delta/\sigma)_{\max} < 0.001$
 3425 reflections $\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$
 241 parameters $\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$
 1 restraint Extinction correction: none
 Primary atom site location: structure-invariant direct methods

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.88109 (14)	0.92607 (17)	0.05530 (7)	0.0462 (4)
N2	1.00455 (13)	0.88709 (16)	0.10140 (7)	0.0446 (3)
N3	1.20940 (14)	0.48961 (18)	0.04801 (7)	0.0461 (3)
O1	0.86564 (13)	1.16125 (15)	0.11112 (6)	0.0575 (4)
O2	0.63397 (15)	1.31345 (18)	0.08033 (7)	0.0689 (4)
H2	0.7084	1.2854	0.1052	0.103*
O3	1.13711 (14)	0.56430 (16)	-0.00173 (6)	0.0612 (4)
O4	1.24674 (14)	0.34954 (16)	0.04217 (7)	0.0695 (4)
O5	1.38574 (15)	0.75885 (19)	0.31532 (6)	0.0721 (4)
H5	1.4565	0.7160	0.3377	0.108*
C1	0.69456 (16)	1.10569 (19)	0.00579 (8)	0.0407 (4)
C2	0.60609 (17)	1.2299 (2)	0.01840 (9)	0.0485 (4)
C3	0.49048 (18)	1.2674 (3)	-0.03235 (10)	0.0601 (5)
H3	0.4332	1.3482	-0.0234	0.072*
C4	0.45611 (17)	1.1865 (3)	-0.09781 (10)	0.0582 (5)
C5	0.3359 (2)	1.2237 (3)	-0.15133 (13)	0.0833 (8)
H5A	0.2756	1.3013	-0.1428	0.100*
C6	0.3090 (3)	1.1465 (4)	-0.21477 (15)	0.1025 (11)
H6	0.2299	1.1717	-0.2493	0.123*
C7	0.3980 (3)	1.0294 (3)	-0.22928 (13)	0.1001 (10)
H7	0.3782	0.9787	-0.2733	0.120*
C8	0.5134 (2)	0.9896 (3)	-0.17908 (11)	0.0769 (7)
H8	0.5721	0.9116	-0.1889	0.092*
C9	0.54456 (19)	1.0664 (2)	-0.11211 (9)	0.0541 (5)
C10	0.66309 (18)	1.0299 (2)	-0.05915 (9)	0.0477 (4)
H10	0.7224	0.9519	-0.0683	0.057*
C11	0.81986 (16)	1.06728 (19)	0.06131 (8)	0.0417 (4)
C12	1.04643 (17)	0.7455 (2)	0.09371 (9)	0.0468 (4)
H12	0.9935	0.6774	0.0596	0.056*
C13	1.17752 (16)	0.68736 (18)	0.13789 (8)	0.0398 (4)
C14	1.25482 (16)	0.56598 (18)	0.11681 (8)	0.0381 (3)
C15	1.37767 (16)	0.51385 (19)	0.16009 (9)	0.0429 (4)
H15	1.4276	0.4342	0.1443	0.051*
C16	1.42572 (16)	0.5796 (2)	0.22626 (8)	0.0440 (4)
H16	1.5094	0.5471	0.2550	0.053*
C17	1.34847 (18)	0.6951 (2)	0.24990 (8)	0.0466 (4)
C18	1.22688 (17)	0.7487 (2)	0.20549 (8)	0.0481 (4)
H18	1.1772	0.8280	0.2216	0.058*

supplementary materials

H1 0.8395 (18) 0.8513 (19) 0.0237 (8) 0.066 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0433 (8)	0.0423 (8)	0.0426 (8)	0.0066 (6)	-0.0077 (6)	-0.0034 (6)
N2	0.0416 (7)	0.0449 (8)	0.0399 (7)	0.0064 (6)	-0.0033 (6)	0.0013 (6)
N3	0.0441 (8)	0.0456 (8)	0.0486 (8)	-0.0067 (6)	0.0117 (7)	-0.0072 (7)
O1	0.0632 (8)	0.0469 (7)	0.0482 (7)	0.0076 (6)	-0.0119 (6)	-0.0087 (6)
O2	0.0754 (10)	0.0743 (10)	0.0563 (8)	0.0261 (8)	0.0159 (7)	-0.0026 (7)
O3	0.0687 (9)	0.0642 (9)	0.0433 (7)	-0.0101 (7)	0.0007 (6)	-0.0002 (6)
O4	0.0683 (9)	0.0559 (8)	0.0786 (9)	0.0076 (7)	0.0085 (7)	-0.0274 (7)
O5	0.0725 (10)	0.0899 (11)	0.0412 (7)	0.0308 (8)	-0.0088 (6)	-0.0124 (7)
C1	0.0390 (8)	0.0370 (8)	0.0416 (8)	-0.0019 (7)	0.0022 (7)	0.0075 (7)
C2	0.0475 (10)	0.0517 (10)	0.0469 (10)	0.0047 (8)	0.0133 (8)	0.0101 (8)
C3	0.0431 (10)	0.0710 (13)	0.0687 (13)	0.0150 (9)	0.0189 (9)	0.0243 (11)
C4	0.0378 (9)	0.0711 (13)	0.0591 (12)	-0.0109 (9)	0.0004 (8)	0.0282 (10)
C5	0.0427 (11)	0.1106 (19)	0.0836 (16)	-0.0086 (12)	-0.0074 (11)	0.0461 (15)
C6	0.0652 (15)	0.120 (2)	0.0910 (19)	-0.0332 (16)	-0.0366 (14)	0.0452 (18)
C7	0.113 (2)	0.0834 (18)	0.0686 (15)	-0.0412 (17)	-0.0400 (15)	0.0161 (13)
C8	0.0945 (16)	0.0613 (13)	0.0542 (12)	-0.0199 (12)	-0.0187 (11)	0.0049 (10)
C9	0.0536 (11)	0.0486 (10)	0.0485 (10)	-0.0154 (9)	-0.0080 (8)	0.0144 (8)
C10	0.0518 (10)	0.0375 (8)	0.0459 (9)	-0.0020 (8)	-0.0022 (8)	0.0045 (7)
C11	0.0437 (9)	0.0382 (8)	0.0384 (8)	0.0000 (7)	0.0016 (7)	0.0026 (7)
C12	0.0455 (9)	0.0440 (9)	0.0434 (9)	0.0023 (8)	-0.0022 (7)	-0.0052 (7)
C13	0.0403 (8)	0.0366 (8)	0.0387 (8)	0.0020 (7)	0.0031 (7)	0.0030 (7)
C14	0.0422 (8)	0.0333 (8)	0.0376 (8)	-0.0044 (7)	0.0082 (7)	0.0001 (6)
C15	0.0438 (9)	0.0363 (8)	0.0497 (9)	0.0043 (7)	0.0141 (8)	0.0035 (7)
C16	0.0409 (9)	0.0443 (9)	0.0430 (9)	0.0060 (7)	0.0038 (7)	0.0090 (7)
C17	0.0506 (10)	0.0494 (10)	0.0350 (8)	0.0053 (8)	0.0022 (7)	0.0009 (7)
C18	0.0499 (10)	0.0462 (9)	0.0436 (9)	0.0140 (8)	0.0037 (8)	-0.0029 (7)

Geometric parameters (\AA , $^\circ$)

N1—C11	1.341 (2)	C5—H5A	0.9300
N1—N2	1.3812 (18)	C6—C7	1.402 (4)
N1—H1	0.898 (9)	C6—H6	0.9300
N2—C12	1.266 (2)	C7—C8	1.362 (3)
N3—O3	1.2235 (18)	C7—H7	0.9300
N3—O4	1.2315 (18)	C8—C9	1.414 (3)
N3—C14	1.448 (2)	C8—H8	0.9300
O1—C11	1.2378 (18)	C9—C10	1.404 (2)
O2—C2	1.357 (2)	C10—H10	0.9300
O2—H2	0.8200	C12—C13	1.469 (2)
O5—C17	1.343 (2)	C12—H12	0.9300
O5—H5	0.8200	C13—C18	1.382 (2)
C1—C10	1.376 (2)	C13—C14	1.399 (2)
C1—C2	1.426 (2)	C14—C15	1.384 (2)
C1—C11	1.479 (2)	C15—C16	1.371 (2)

C2—C3	1.364 (2)	C15—H15	0.9300
C3—C4	1.405 (3)	C16—C17	1.387 (2)
C3—H3	0.9300	C16—H16	0.9300
C4—C9	1.414 (3)	C17—C18	1.387 (2)
C4—C5	1.422 (3)	C18—H18	0.9300
C5—C6	1.357 (4)		
C11—N1—N2	120.64 (13)	C9—C8—H8	119.8
C11—N1—H1	120.7 (13)	C10—C9—C4	118.26 (17)
N2—N1—H1	118.4 (13)	C10—C9—C8	122.1 (2)
C12—N2—N1	114.48 (14)	C4—C9—C8	119.63 (18)
O3—N3—O4	122.43 (14)	C1—C10—C9	122.28 (17)
O3—N3—C14	120.03 (14)	C1—C10—H10	118.9
O4—N3—C14	117.54 (14)	C9—C10—H10	118.9
C2—O2—H2	109.5	O1—C11—N1	121.64 (14)
C17—O5—H5	109.5	O1—C11—C1	121.38 (15)
C10—C1—C2	118.58 (15)	N1—C11—C1	116.98 (14)
C10—C1—C11	122.05 (15)	N2—C12—C13	120.90 (15)
C2—C1—C11	119.31 (14)	N2—C12—H12	119.5
O2—C2—C3	118.43 (17)	C13—C12—H12	119.5
O2—C2—C1	121.50 (15)	C18—C13—C14	116.75 (14)
C3—C2—C1	120.07 (17)	C18—C13—C12	119.55 (15)
C2—C3—C4	121.34 (18)	C14—C13—C12	123.64 (14)
C2—C3—H3	119.3	C15—C14—C13	121.96 (14)
C4—C3—H3	119.3	C15—C14—N3	116.99 (14)
C3—C4—C9	119.42 (16)	C13—C14—N3	121.04 (14)
C3—C4—C5	122.1 (2)	C16—C15—C14	119.98 (15)
C9—C4—C5	118.4 (2)	C16—C15—H15	120.0
C6—C5—C4	120.2 (3)	C14—C15—H15	120.0
C6—C5—H5A	119.9	C15—C16—C17	119.39 (15)
C4—C5—H5A	119.9	C15—C16—H16	120.3
C5—C6—C7	121.3 (2)	C17—C16—H16	120.3
C5—C6—H6	119.4	O5—C17—C16	122.72 (15)
C7—C6—H6	119.4	O5—C17—C18	117.25 (16)
C8—C7—C6	120.1 (3)	C16—C17—C18	120.03 (15)
C8—C7—H7	119.9	C13—C18—C17	121.79 (16)
C6—C7—H7	119.9	C13—C18—H18	119.1
C7—C8—C9	120.3 (3)	C17—C18—H18	119.1
C7—C8—H8	119.8		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 \cdots O1	0.82	1.88	2.5987 (18)	146
N1—H1 \cdots O4 ⁱ	0.898 (9)	2.144 (10)	3.0361 (19)	172.8 (18)
O5—H5 \cdots O1 ⁱⁱ	0.82	1.88	2.6889 (17)	168

Symmetry codes: (i) $-x+2, -y+1, -z$; (ii) $-x+5/2, y-1/2, -z+1/2$.

Fig. 1




Fig. 2

