

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# *cis*-1,2-Bis[[4-(4-pyridyl)pyrimidin-2-yl]-sulfanylmethyl]benzene

 Hua-Ze Dong,<sup>a\*</sup> Hai-Bin Zhu<sup>b</sup> and Shao-Hua Gou<sup>b</sup>

<sup>a</sup>Department of Chemistry and Chemical Engineering, Hefei Normal University, Hefei 230061, People's Republic of China, and <sup>b</sup>School of Chemistry and Chemical Engineering, Southeast University, Nanjing 211189, People's Republic of China  
Correspondence e-mail: dapdong@163.com


Received 25 October 2009; accepted 29 October 2009

Key indicators: single-crystal X-ray study;  $T = 291$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.052;  $wR$  factor = 0.108; data-to-parameter ratio = 16.0.

The molecular skeleton of the title molecule,  $\text{C}_{26}\text{H}_{20}\text{N}_6\text{S}_2$ , adopts a *cis* conformation with the two arms positioned on one side of the benzene ring plane. Intramolecular  $\pi$ - $\pi$  interactions between the pyrimidine rings [centroid-centroid distance = 3.654 (2) Å] and between the pyridine rings [centroid-centroid distance = 3.775 (2) Å] help to set the molecular conformation; the pyrimidine rings, as well as the pyridine rings, are nearly parallel, forming dihedral angles of 4.12 (14) and 2.46 (14)°, respectively.

## Related literature

For related compounds, see: Dong *et al.* (2008, 2009); Huang *et al.* (2007).



## Experimental

## Crystal data

$\text{C}_{26}\text{H}_{20}\text{N}_6\text{S}_2$   
 $M_r = 480.62$   
 Monoclinic,  $P2_1/n$   
 $a = 8.6078$  (11) Å  
 $b = 27.102$  (3) Å  
 $c = 10.0282$  (12) Å  
 $\beta = 103.661$  (2)°  
 $V = 2273.3$  (5) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.26$  mm<sup>-1</sup>  
 $T = 291$  K  
 $0.32 \times 0.18 \times 0.16$  mm

## Data collection

Bruker SMART CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2000)  
 $T_{\min} = 0.917$ ,  $T_{\max} = 0.966$   
 13064 measured reflections  
 4903 independent reflections  
 3067 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.052$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$   
 $wR(F^2) = 0.108$   
 $S = 0.90$   
 4903 reflections  
 307 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.32$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.22$  e Å<sup>-3</sup>

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

The authors are indebted to the National Natural Science Foundation of China (grant No. 20801011) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2646).

## References

- Bruker (2000). SADABS, SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Dong, H. Z., Zhao, J., Zhu, H. B. & Gou, S. H. (2009). *Polyhedron*, **28**, 1040–1048.  
 Dong, H. Z., Zhu, H. B., Tong, T. F. & Gou, S. H. (2008). *J. Mol. Struct.* **891**, 266–271.  
 Huang, C. H., Gou, S. H., Zhu, H. B. & Huang, W. (2007). *Inorg. Chem.* **46**, 5537–5543.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supporting information

*Acta Cryst.* (2009). E65, o2977 [doi:10.1107/S1600536809045450]

***cis*-1,2-Bis{[4-(4-pyridyl)pyrimidin-2-yl]sulfanylmethyl}benzene**

Hua-Ze Dong, Hai-Bin Zhu and Shao-Hua Gou

**S1. Comment**

Remarkable attention has been paid to the rational design and assembly of new coordination polymers with heterocyclic thiolates or thioethers in recent years. In our previous work, we reported a series of dithioether ligands (Dong *et al.*, 2008; 2009; Huang *et al.*, 2007). As our continuing study, herein we report the molecular structure of the title compound - the newly synthesized ligand derived from 4-(4-pyridinyl)pyrimidine-2-thiol.


The molecular structure of the title compound is shown in Fig. 1. The molecule adopts a *cis* conformation with two arms positioned on one side of the benzene ring plane. It is noted that intramolecular  $\pi$ - $\pi$  interactions between the pyrimidinyl rings [centroid-centroid distance of 3.654 (2) Å] and between the pyridinyl rings [centroid-centroid distance of 3.775 (2) Å] set the molecular conformation - the pyrimidinyl rings, as well as the pyridinyl ones, are nearly parallel forming dihedral angles of 4.12 (14)° and 2.46 (14)°, respectively.

**S2. Experimental**

All solvents and chemicals were of analytical grade and were used without further purification. The title compound was prepared by similar procedure reported in the literature (Dong *et al.*, 2008; 2009). To a solution of 4-(4-pyridinyl)pyrimidine-2-thiol (3.78 g, 20 mmol) and sodium hydroxide (0.80 g, 20 mmol) in dry ethanol (300 ml), 1,2-bis-(bromomethyl)benzene (2.64 g, 10 mmol) in CCl<sub>4</sub> (30 ml) was added. The mixture was stirred and refluxed for 8 h. After cooling, precipitates were filtered, washed in water and ethanol, and dried in vacuum. Anal. Calcd for C<sub>26</sub>H<sub>20</sub>N<sub>6</sub>S<sub>2</sub>: C, 64.98; H, 4.19; N, 17.49%. Found: C, 65.07; H, 4.52; N, 17.34%. Single crystals of ligand suitable for X-ray diffraction were grown from methanol solution by slow evaporation in air at room temperature.

**S3. Refinement**

All hydrogen atoms were geometrically positioned (C—H 0.93–0.97 Å) and refined as riding, with  $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$ .

**Figure 1**

The molecular structure of the title compound showing 30% probability displacement ellipsoids and the atom-numbering scheme.

### ***cis*-1,2-Bis[[4-(4-pyridyl)pyrimidin-2-yl]sulfanylmethyl]benzene**

#### *Crystal data*

$C_{26}H_{20}N_6S_2$

$M_r = 480.62$

Monoclinic,  $P2_1/n$

Hall symbol:  $-P\ 2_1n$

$a = 8.6078$  (11) Å

$b = 27.102$  (3) Å

$c = 10.0282$  (12) Å

$\beta = 103.661$  (2)°

$V = 2273.3$  (5) Å<sup>3</sup>

$Z = 4$

$F(000) = 1000.0$

$D_x = 1.404$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4903 reflections

$\theta = 2.2$ – $27.0$ °

$\mu = 0.26$  mm<sup>-1</sup>

$T = 291$  K

Block, pale yellow

$0.32 \times 0.18 \times 0.16$  mm

#### *Data collection*

Bruker SMART CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2000)

$T_{\min} = 0.917$ ,  $T_{\max} = 0.966$

13064 measured reflections

4903 independent reflections

3067 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.052$

$\theta_{\max} = 27.0$ °,  $\theta_{\min} = 2.2$ °

$h = -10 \rightarrow 8$

$k = -34 \rightarrow 31$

$l = -12 \rightarrow 12$

Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.052$   
 $wR(F^2) = 0.108$   
 $S = 0.90$   
 4903 reflections  
 307 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0421P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7345 (3)	0.22381 (8)	0.8229 (2)	0.0384 (5)
C2	0.9748 (3)	0.25202 (9)	0.9324 (2)	0.0494 (6)
H2	1.0799	0.2469	0.9807	0.059*
C3	0.9230 (3)	0.29935 (8)	0.9059 (2)	0.0474 (6)
H3	0.9909	0.3259	0.9351	0.057*
C4	0.7668 (3)	0.30680 (8)	0.8343 (2)	0.0411 (6)
C5	0.6967 (3)	0.35654 (8)	0.8048 (2)	0.0436 (6)
C6	0.7521 (3)	0.39620 (9)	0.8867 (3)	0.0623 (8)
H6	0.8355	0.3922	0.9640	0.075*
C7	0.6845 (4)	0.44177 (10)	0.8545 (3)	0.0774 (9)
H7	0.7240	0.4680	0.9125	0.093*
C8	0.5126 (3)	0.41215 (9)	0.6685 (3)	0.0690 (8)
H8	0.4291	0.4172	0.5919	0.083*
C9	0.5716 (3)	0.36519 (8)	0.6935 (3)	0.0529 (7)
H9	0.5275	0.3394	0.6357	0.063*
C10	0.6875 (3)	0.12421 (7)	0.8693 (2)	0.0472 (6)
H10A	0.7817	0.1124	0.8418	0.057*
H10B	0.7188	0.1342	0.9648	0.057*
C11	0.5630 (3)	0.08400 (8)	0.8510 (2)	0.0443 (6)
C12	0.5488 (3)	0.05264 (9)	0.7392 (3)	0.0592 (7)
H12	0.6183	0.0560	0.6815	0.071*
C13	0.4334 (4)	0.01686 (9)	0.7133 (3)	0.0687 (8)
H13	0.4248	-0.0038	0.6379	0.082*
C14	0.3317 (3)	0.01133 (9)	0.7966 (3)	0.0660 (8)
H14	0.2519	-0.0126	0.7777	0.079*

C15	0.3473 (3)	0.04123 (8)	0.9091 (3)	0.0566 (7)
H15	0.2794	0.0367	0.9677	0.068*
C16	0.4617 (3)	0.07813 (8)	0.9376 (2)	0.0446 (6)
C17	0.4698 (3)	0.11019 (7)	1.0609 (2)	0.0502 (6)
H17A	0.5805	0.1184	1.1017	0.060*
H17B	0.4283	0.0921	1.1285	0.060*
C18	0.4950 (3)	0.21357 (8)	1.0767 (2)	0.0380 (5)
C19	0.7376 (3)	0.24131 (9)	1.1805 (2)	0.0484 (6)
H19	0.8442	0.2358	1.2238	0.058*
C20	0.6847 (3)	0.28872 (8)	1.1637 (2)	0.0446 (6)
H20	0.7523	0.3151	1.1955	0.053*
C21	0.5267 (3)	0.29608 (8)	1.0977 (2)	0.0393 (5)
C22	0.4562 (3)	0.34572 (8)	1.0709 (2)	0.0399 (5)
C23	0.5154 (3)	0.38590 (9)	1.1509 (3)	0.0589 (7)
H23	0.6024	0.3823	1.2253	0.071*
C24	0.4452 (4)	0.43133 (9)	1.1201 (3)	0.0701 (9)
H24	0.4868	0.4579	1.1761	0.084*
C25	0.3284 (3)	0.35403 (8)	0.9626 (2)	0.0497 (6)
H25	0.2835	0.3282	0.9052	0.060*
C26	0.2679 (3)	0.40072 (9)	0.9402 (3)	0.0597 (7)
H26	0.1816	0.4054	0.8656	0.072*
N1	0.8822 (2)	0.21295 (7)	0.89261 (18)	0.0458 (5)
N2	0.6713 (2)	0.26820 (6)	0.79073 (17)	0.0405 (5)
N3	0.5667 (3)	0.45081 (8)	0.7465 (3)	0.0771 (8)
N4	0.6443 (2)	0.20240 (7)	1.13813 (19)	0.0463 (5)
N5	0.4294 (2)	0.25802 (6)	1.05275 (17)	0.0394 (5)
N6	0.3221 (3)	0.43947 (7)	1.0157 (2)	0.0640 (6)
S1	0.60063 (8)	0.17575 (2)	0.76342 (6)	0.04810 (19)
S2	0.35623 (8)	0.16649 (2)	1.01669 (6)	0.04634 (19)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0353 (14)	0.0411 (13)	0.0382 (13)	-0.0043 (11)	0.0074 (11)	0.0000 (10)
C2	0.0363 (16)	0.0586 (16)	0.0487 (15)	-0.0038 (12)	0.0005 (12)	0.0005 (12)
C3	0.0409 (16)	0.0496 (15)	0.0480 (15)	-0.0112 (12)	0.0031 (12)	-0.0033 (11)
C4	0.0424 (15)	0.0443 (13)	0.0377 (13)	-0.0058 (12)	0.0116 (11)	-0.0023 (10)
C5	0.0423 (16)	0.0436 (14)	0.0462 (14)	-0.0055 (11)	0.0131 (12)	-0.0066 (11)
C6	0.068 (2)	0.0518 (16)	0.0583 (17)	-0.0030 (15)	-0.0019 (14)	-0.0126 (13)
C7	0.090 (3)	0.0455 (17)	0.086 (2)	-0.0035 (16)	-0.001 (2)	-0.0243 (15)
C8	0.063 (2)	0.0504 (16)	0.081 (2)	0.0088 (14)	-0.0070 (16)	-0.0122 (15)
C9	0.0461 (17)	0.0422 (14)	0.0656 (17)	-0.0020 (12)	0.0036 (14)	-0.0122 (12)
C10	0.0394 (15)	0.0401 (13)	0.0559 (15)	0.0036 (11)	-0.0010 (12)	0.0066 (11)
C11	0.0447 (16)	0.0305 (12)	0.0509 (15)	0.0019 (11)	-0.0025 (12)	0.0012 (11)
C12	0.064 (2)	0.0480 (15)	0.0640 (18)	-0.0002 (14)	0.0131 (15)	-0.0030 (13)
C13	0.079 (2)	0.0455 (16)	0.072 (2)	-0.0027 (15)	-0.0014 (18)	-0.0151 (14)
C14	0.058 (2)	0.0367 (15)	0.094 (2)	-0.0075 (13)	-0.0005 (17)	-0.0006 (15)
C15	0.0512 (18)	0.0380 (14)	0.079 (2)	0.0012 (12)	0.0129 (15)	0.0064 (13)

C16	0.0457 (16)	0.0304 (12)	0.0526 (15)	0.0027 (11)	0.0016 (12)	0.0056 (11)
C17	0.0563 (17)	0.0387 (13)	0.0527 (16)	0.0057 (12)	0.0071 (13)	0.0101 (11)
C18	0.0354 (15)	0.0429 (13)	0.0355 (13)	0.0000 (11)	0.0084 (11)	-0.0030 (10)
C19	0.0335 (15)	0.0601 (16)	0.0471 (15)	0.0018 (13)	0.0005 (12)	0.0007 (12)
C20	0.0357 (15)	0.0464 (14)	0.0473 (14)	-0.0037 (11)	0.0014 (12)	-0.0047 (11)
C21	0.0379 (14)	0.0453 (13)	0.0340 (13)	-0.0012 (11)	0.0070 (11)	-0.0043 (10)
C22	0.0364 (14)	0.0410 (13)	0.0417 (13)	-0.0033 (11)	0.0080 (11)	-0.0054 (10)
C23	0.0475 (17)	0.0524 (16)	0.0654 (18)	0.0002 (13)	-0.0096 (14)	-0.0141 (13)
C24	0.065 (2)	0.0455 (16)	0.088 (2)	-0.0040 (14)	-0.0062 (17)	-0.0218 (14)
C25	0.0514 (17)	0.0407 (14)	0.0511 (15)	0.0003 (12)	0.0003 (13)	-0.0089 (11)
C26	0.0584 (19)	0.0521 (16)	0.0590 (17)	0.0073 (14)	-0.0057 (14)	-0.0017 (13)
N1	0.0354 (13)	0.0505 (12)	0.0475 (12)	-0.0011 (10)	0.0016 (10)	0.0037 (9)
N2	0.0357 (12)	0.0397 (11)	0.0440 (11)	-0.0041 (9)	0.0054 (9)	-0.0016 (9)
N3	0.079 (2)	0.0508 (14)	0.0889 (19)	0.0089 (13)	-0.0057 (15)	-0.0148 (13)
N4	0.0392 (13)	0.0472 (11)	0.0494 (12)	0.0050 (10)	0.0041 (10)	0.0038 (9)
N5	0.0350 (12)	0.0394 (11)	0.0420 (11)	0.0016 (9)	0.0053 (9)	-0.0027 (8)
N6	0.0645 (17)	0.0434 (13)	0.0754 (16)	0.0052 (11)	-0.0007 (13)	-0.0055 (11)
S1	0.0409 (4)	0.0417 (3)	0.0538 (4)	-0.0057 (3)	-0.0044 (3)	0.0051 (3)
S2	0.0425 (4)	0.0402 (3)	0.0539 (4)	-0.0003 (3)	0.0067 (3)	-0.0024 (3)

*Geometric parameters (Å, °)*

C1—N2	1.328 (2)	C13—H13	0.9300
C1—N1	1.331 (3)	C14—C15	1.369 (3)
C1—S1	1.748 (2)	C14—H14	0.9300
C2—N1	1.329 (3)	C15—C16	1.385 (3)
C2—C3	1.364 (3)	C15—H15	0.9300
C2—H2	0.9300	C16—C17	1.500 (3)
C3—C4	1.381 (3)	C17—S2	1.810 (2)
C3—H3	0.9300	C17—H17A	0.9700
C4—N2	1.338 (2)	C17—H17B	0.9700
C4—C5	1.478 (3)	C18—N4	1.323 (3)
C5—C6	1.369 (3)	C18—N5	1.328 (2)
C5—C9	1.375 (3)	C18—S2	1.754 (2)
C6—C7	1.371 (3)	C19—N4	1.333 (3)
C6—H6	0.9300	C19—C20	1.360 (3)
C7—N3	1.320 (3)	C19—H19	0.9300
C7—H7	0.9300	C20—C21	1.379 (3)
C8—N3	1.324 (3)	C20—H20	0.9300
C8—C9	1.372 (3)	C21—N5	1.339 (2)
C8—H8	0.9300	C21—C22	1.474 (3)
C9—H9	0.9300	C22—C25	1.369 (3)
C10—C11	1.509 (3)	C22—C23	1.377 (3)
C10—S1	1.807 (2)	C23—C24	1.374 (3)
C10—H10A	0.9700	C23—H23	0.9300
C10—H10B	0.9700	C24—N6	1.321 (3)
C11—C16	1.377 (3)	C24—H24	0.9300
C11—C12	1.389 (3)	C25—C26	1.367 (3)

C12—C13	1.368 (3)	C25—H25	0.9300
C12—H12	0.9300	C26—N6	1.314 (3)
C13—C14	1.354 (4)	C26—H26	0.9300
N2—C1—N1	127.8 (2)	C14—C15—H15	119.2
N2—C1—S1	113.12 (17)	C16—C15—H15	119.2
N1—C1—S1	119.04 (17)	C11—C16—C15	118.6 (2)
N1—C2—C3	123.0 (2)	C11—C16—C17	122.7 (2)
N1—C2—H2	118.5	C15—C16—C17	118.7 (2)
C3—C2—H2	118.5	C16—C17—S2	111.68 (15)
C2—C3—C4	118.2 (2)	C16—C17—H17A	109.3
C2—C3—H3	120.9	S2—C17—H17A	109.3
C4—C3—H3	120.9	C16—C17—H17B	109.3
N2—C4—C3	120.2 (2)	S2—C17—H17B	109.3
N2—C4—C5	117.2 (2)	H17A—C17—H17B	107.9
C3—C4—C5	122.6 (2)	N4—C18—N5	128.1 (2)
C6—C5—C9	116.6 (2)	N4—C18—S2	120.10 (17)
C6—C5—C4	121.9 (2)	N5—C18—S2	111.83 (17)
C9—C5—C4	121.5 (2)	N4—C19—C20	123.3 (2)
C5—C6—C7	119.8 (2)	N4—C19—H19	118.4
C5—C6—H6	120.1	C20—C19—H19	118.4
C7—C6—H6	120.1	C19—C20—C21	117.4 (2)
N3—C7—C6	124.2 (2)	C19—C20—H20	121.3
N3—C7—H7	117.9	C21—C20—H20	121.3
C6—C7—H7	117.9	N5—C21—C20	121.2 (2)
N3—C8—C9	124.2 (2)	N5—C21—C22	116.3 (2)
N3—C8—H8	117.9	C20—C21—C22	122.4 (2)
C9—C8—H8	117.9	C25—C22—C23	116.7 (2)
C8—C9—C5	119.5 (2)	C25—C22—C21	120.9 (2)
C8—C9—H9	120.3	C23—C22—C21	122.4 (2)
C5—C9—H9	120.3	C24—C23—C22	119.7 (2)
C11—C10—S1	107.66 (15)	C24—C23—H23	120.1
C11—C10—H10A	110.2	C22—C23—H23	120.1
S1—C10—H10A	110.2	N6—C24—C23	123.7 (2)
C11—C10—H10B	110.2	N6—C24—H24	118.1
S1—C10—H10B	110.2	C23—C24—H24	118.1
H10A—C10—H10B	108.5	C26—C25—C22	119.1 (2)
C16—C11—C12	119.3 (2)	C26—C25—H25	120.4
C16—C11—C10	122.9 (2)	C22—C25—H25	120.4
C12—C11—C10	117.7 (2)	N6—C26—C25	125.1 (2)
C13—C12—C11	120.6 (3)	N6—C26—H26	117.4
C13—C12—H12	119.7	C25—C26—H26	117.4
C11—C12—H12	119.7	C2—N1—C1	114.35 (19)
C14—C13—C12	120.5 (3)	C1—N2—C4	116.4 (2)
C14—C13—H13	119.8	C7—N3—C8	115.6 (2)
C12—C13—H13	119.8	C18—N4—C19	114.42 (19)
C13—C14—C15	119.4 (3)	C18—N5—C21	115.62 (19)
C13—C14—H14	120.3	C26—N6—C24	115.6 (2)

## supporting information

---

C15—C14—H14	120.3	C1—S1—C10	103.16 (11)
C14—C15—C16	121.6 (3)	C18—S2—C17	104.16 (11)

---