

Tetra- μ -benzoato-bis[[*trans*-1-(2-pyridyl)-2-(4-pyridyl)ethylene]zinc(II)]

Young Joo Song,^a Soo-Won Lee,^b Kyung Hwan Jang,^c
Cheal Kim^{a*} and Youngmee Kim^{d*}

^aDepartment of Fine Chemistry, and Eco-Product and Materials Education Center, Seoul National University of Technology, Seoul 139-743, Republic of Korea, ^bForest Practice Research Center, Korea Forest Research Institute, Pocheon 487-821, Republic of Korea, ^cKorea Forest Research Institute 44-3, Suwon 441-350, Republic of Korea, and ^dDepartment of Chemistry and Nano Science, Ewha Womans University, Seoul 120-750, Republic of Korea

Correspondence e-mail: chealkim@sunt.ac.kr, ymeekim@ewha.ac.kr


Received 21 October 2009; accepted 28 October 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.039; wR factor = 0.090; data-to-parameter ratio = 14.8.

The paddle-wheel-type centrosymmetric dinuclear title complex, $[\text{Zn}_2(\text{C}_7\text{H}_5\text{O}_2)_4(\text{C}_{12}\text{H}_{10}\text{N}_2)_2]$, contains four bridging benzoate groups and two terminal *trans*-1-(2-pyridyl)-2-(4-pyridyl)ethylene (*L*) ligands. The inversion center is located between the two Zn^{II} atoms. The octahedral coordination around the Zn^{II} atom, with four O atoms in the equatorial plane, is completed by an N atom of the *L* molecule [$\text{Zn}-\text{N} = 2.0198$ (15) Å] and by the second Zn^{II} atom [$\text{Zn}\cdots\text{Zn} = 2.971$ (8) Å]. The Zn^{II} atom is 0.372 Å out of the plane of the four coordinating O atoms.

Related literature

For structures containing $[\text{Zn}_2(\text{O}_2\text{CPh})_4]$, see: Necefoglu *et al.* (2002); Zeleňák *et al.* (2004); Karmakar *et al.* (2006); Ohmura *et al.* (2005). For the structures of copper(II) and zinc(II) benzoates with quinoxaline, 6-methylquinoline, 3-methylquinoline, and di-2-pyridyl ketone, see: Lee *et al.* (2008); Yu *et al.* (2008, 2009); Park *et al.* (2008); Shin *et al.* (2009). For transition metal ions as the major cation contributors to the inorganic composition of natural water and biological fluids, see: Daniele *et al.* (2008); Parkin (2004); Tshuva & Lippard (2004).



Experimental

Crystal data

$[\text{Zn}_2(\text{C}_7\text{H}_5\text{O}_2)_4(\text{C}_{12}\text{H}_{10}\text{N}_2)_2]$

$M_r = 979.66$

Monoclinic, $C2/c$

$a = 24.919$ (6) Å

$b = 12.186$ (3) Å

$c = 15.742$ (4) Å

$\beta = 109.857$ (4)°

$V = 4496.0$ (19) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 1.13$ mm⁻¹

$T = 293$ K

0.20 × 0.15 × 0.15 mm

Data collection

Bruker SMART CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 1997)

$T_{\text{min}} = 0.816$, $T_{\text{max}} = 0.884$

12326 measured reflections

4416 independent reflections

2947 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.090$

$S = 1.03$

4416 reflections

298 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.26$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.27$ e Å⁻³

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINTE* (Bruker, 1997); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Financial support from the Korean Ministry of the Environment "ET-Human resource development Project" and the Cooperative Research Program for Agricultural Science & Technology Development (20070301-036-019-02) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2505).

References

- Bruker (1997). *SMART*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Daniele, P. G., Foti, C., Gianguzza, A., Prenesti, E. & Sammartano, S. (2008). *Coord. Chem. Rev.* **252**, 1093–1107.
- Karmakar, A., Sarma, R. J. & Baruah, J. B. (2006). *Inorg. Chem. Commun.* **9**, 1169–1172.
- Lee, E. Y., Park, B. K., Kim, C., Kim, S.-J. & Kim, Y. (2008). *Acta Cryst.* **E64**, m286.
- Necefoglu, H., Clegg, W. & Scott, A. J. (2002). *Acta Cryst.* **E58**, m121–m122.

- Ohmura, T., Mori, W., Takei, T., Ikeda, T. & Maeda, A. (2005). *Mater. Sci. Pol.* **23**, 729–736.
- Park, B. K., Jang, K.-H., Kim, P.-G., Kim, C. & Kim, Y. (2008). *Acta Cryst.* **E64**, m1141.
- Parkin, G. (2004). *Chem. Rev.* **104**, 699–767.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shin, D. H., Han, S.-H., Kim, P.-G., Kim, C. & Kim, Y. (2009). *Acta Cryst.* **E65**, m658–m659.
- Tshuva, E. Y. & Lippard, S. J. (2004). *Chem. Rev.* **104**, 987–1012.
- Yu, S. M., Park, C.-H., Kim, P.-G., Kim, C. & Kim, Y. (2008). *Acta Cryst.* **E64**, m881–m882.
- Yu, S. M., Shin, D. H., Kim, P.-G., Kim, C. & Kim, Y. (2009). *Acta Cryst.* **E65**, m1045–m1046.
- Zeleňák, V., Sabo, M., Massa, W. & Černák, J. (2004). *Acta Cryst.* **C60**, m85–m87.

supplementary materials

Acta Cryst. (2009). E65, m1495-m1496 [doi:10.1107/S1600536809045048]

Tetra- μ -benzoato-bis{[*trans*-1-(2-pyridyl)-2-(4-pyridyl)ethylene]zinc(II)}

Y. J. Song, S.-W. Lee, K. H. Jang, C. Kim and Y. Kim

Comment

A great attention has been paid to transition metal ions as the major cation contributors to the inorganic composition of natural water and biological fluids (Daniele, *et al.*, 2008; Parkin, 2004; Tshuva & Lippard, 2004). While the main attention was focused on the interaction of transition metal ions with biologically active molecules such as amino acids, proteins, sugars, nucleotides *etc.*, the study on the interaction of the transition metal ions with fulvic acids and humic acids, mainly found in soil, is about to start. As models to examine the interaction, therefore, we have previously used copper(II) and zinc(II) benzoates as building blocks and reported the structures of copper(II) and zinc(II) benzoates with quinoxaline, 6-methylquinoline, 3-methylquinoline, and di-2-pyridyl ketone (Lee, *et al.*, 2008; Yu, *et al.*, 2008; Park, *et al.*, 2008; Shin, *et al.*, 2009; Yu, *et al.*, 2009). The related paddle-wheel type structures for Zn complexes have been previously reported (Necefoglu *et al.*, 2002; Zeleňák, *et al.*, 2004; Karmakar, *et al.*, 2006; Ohmura, *et al.*, 2005). In this work, we have employed zinc(II) benzoate as a building block and *trans*-1-(2-pyridyl)-2-(4-pyridyl)ethylene as a ligand. We report hereon the structure of new zinc(II) benzoate with *trans*-1-(2-pyridyl)-2-(4-pyridyl)ethylene.

Asymmetric unit contains half of whole molecule, and there is an inversion center in the middle of Zn \cdots Zn bond. Symmetric operation (1-*x*, 1-*y*, 1-*z*) produces a paddle-wheel type dinuclear zinc-benzoate complex (Fig. 1). The paddle-wheel type dinuclear complex is constructed by four bridging benzoate groups and two terminal *L* ligands (*L* = *trans*-1-(2-pyridyl)-2-(4-pyridyl)ethylene). The octahedral coordination around the zinc atom, with four O atoms in the equatorial plane, is completed by nitrogen atom of *L* molecule (Zn—N 2.0198 (15) Å) and by the second zinc atom (Zn \cdots Zn 2.971 (8) Å). The zinc atom is 0.372 Å out of the plane of the four oxygen atoms.

Experimental

30.4 mg (0.1 mmol) of Zn(NO₃)₂·6H₂O and 28.0 mg (0.2 mmol) of C₆H₅COONH₄ were dissolved in 4 ml H₂O and carefully layered by 4 ml methanol solution of *trans*-1-(2-pyridyl)-2-(4-pyridyl)ethylene (37.6 mg, 0.2 mmol). Suitable crystals of the title compound for X-ray analysis were obtained in a few weeks.

Refinement

H atoms were placed in calculated positions with C—H distances of 0.93 Å. They were included in the refinement in a riding-motion approximation with U_{iso}(H) = 1.2U_{eq}(C).

Figures




Fig. 1. The structure of the title compound showing the atom-labeling scheme. Displacement ellipsoids are shown at the 30% probability level. H atoms have been omitted for clarity. [Symmetry code: (i) $-x+1, -y+1, -z+1$].

Tetra- μ -benzoato-bis{[*trans*-1-(2-pyridyl)-2-(4-pyridyl)ethylene]zinc(II)}

Crystal data

$[\text{Zn}_2(\text{C}_7\text{H}_5\text{O}_2)_4(\text{C}_{12}\text{H}_{10}\text{N}_2)_2]$

$M_r = 979.66$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 24.919\ (6)\ \text{\AA}$

$b = 12.186\ (3)\ \text{\AA}$

$c = 15.742\ (4)\ \text{\AA}$

$\beta = 109.857\ (4)^\circ$

$V = 4496.0\ (19)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 2016$

$D_x = 1.447\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1818 reflections

$\theta = 2.5\text{--}19.6^\circ$

$\mu = 1.13\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Block, colorless

$0.20 \times 0.15 \times 0.15\ \text{mm}$

Data collection

Bruker SMART CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293\ \text{K}$

ϕ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 1997)

$T_{\min} = 0.816$, $T_{\max} = 0.884$

12326 measured reflections

4416 independent reflections

2947 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.039$

$\theta_{\max} = 26.0^\circ$

$\theta_{\min} = 1.9^\circ$

$h = -20 \rightarrow 30$

$k = -15 \rightarrow 15$

$l = -19 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.090$

$S = 1.03$

4416 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0205P)^2 + 1.48P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.26\ \text{e \AA}^{-3}$

298 parameters

$$\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.545232 (12)	0.50480 (2)	0.590326 (19)	0.03852 (11)
O11	0.48081 (8)	0.42174 (16)	0.61458 (13)	0.0536 (5)
O12	0.58816 (8)	0.58560 (17)	0.51974 (13)	0.0589 (5)
O21	0.56818 (8)	0.35925 (15)	0.54865 (13)	0.0549 (5)
O22	0.50102 (8)	0.64874 (15)	0.58505 (13)	0.0581 (6)
N31	0.60407 (9)	0.52228 (16)	0.71558 (14)	0.0391 (5)
N32	0.75201 (11)	0.7049 (2)	1.18938 (17)	0.0710 (8)
C11	0.43281 (12)	0.3906 (2)	0.56234 (19)	0.0420 (7)
C12	0.39861 (11)	0.3173 (2)	0.60095 (18)	0.0399 (6)
C13	0.41930 (13)	0.2875 (3)	0.6908 (2)	0.0584 (8)
H13	0.4539	0.3159	0.7282	0.070*
C14	0.38945 (18)	0.2165 (3)	0.7258 (3)	0.0805 (11)
H14	0.4041	0.1973	0.7865	0.097*
C15	0.33831 (18)	0.1736 (3)	0.6724 (3)	0.0803 (11)
H15	0.3184	0.1251	0.6963	0.096*
C16	0.31681 (14)	0.2032 (3)	0.5828 (3)	0.0747 (10)
H16	0.2821	0.1745	0.5459	0.090*
C17	0.34645 (12)	0.2755 (2)	0.5472 (2)	0.0563 (8)
H17	0.3312	0.2962	0.4869	0.068*
C21	0.53915 (12)	0.3100 (2)	0.47807 (19)	0.0423 (6)
C22	0.55306 (11)	0.1917 (2)	0.46933 (19)	0.0450 (7)
C23	0.51906 (15)	0.1303 (3)	0.3980 (3)	0.0773 (11)
H23	0.4894	0.1636	0.3525	0.093*
C24	0.5289 (2)	0.0196 (3)	0.3940 (4)	0.1091 (17)
H24	0.5051	-0.0218	0.3464	0.131*
C25	0.5726 (2)	-0.0295 (3)	0.4583 (4)	0.1087 (17)
H25	0.5786	-0.1044	0.4551	0.130*
C26	0.6079 (2)	0.0306 (3)	0.5279 (3)	0.0902 (13)
H26	0.6384	-0.0034	0.5714	0.108*
C27	0.59863 (14)	0.1418 (3)	0.5344 (2)	0.0628 (9)

supplementary materials

H27	0.6228	0.1826	0.5820	0.075*
C31	0.65674 (12)	0.4837 (2)	0.73577 (19)	0.0531 (8)
H31	0.6658	0.4437	0.6922	0.064*
C32	0.69861 (12)	0.5004 (2)	0.81848 (19)	0.0568 (8)
H32	0.7351	0.4728	0.8292	0.068*
C33	0.68646 (11)	0.5579 (2)	0.88547 (17)	0.0412 (7)
C34	0.63111 (11)	0.5948 (2)	0.86490 (17)	0.0473 (7)
H34	0.6203	0.6323	0.9080	0.057*
C35	0.59211 (11)	0.5760 (2)	0.78060 (17)	0.0456 (7)
H35	0.5552	0.6024	0.7681	0.055*
C36	0.73130 (12)	0.5774 (2)	0.97322 (18)	0.0505 (7)
H36	0.7680	0.5540	0.9794	0.061*
C37	0.72419 (12)	0.6248 (2)	1.04352 (18)	0.0509 (8)
H37	0.6872	0.6455	1.0378	0.061*
C38	0.76886 (13)	0.6485 (2)	1.13035 (18)	0.0473 (7)
C39	0.82430 (14)	0.6161 (3)	1.1499 (2)	0.0647 (9)
H39	0.8352	0.5770	1.1077	0.078*
C310	0.86366 (15)	0.6418 (3)	1.2323 (2)	0.0819 (12)
H310	0.9014	0.6194	1.2467	0.098*
C311	0.84732 (15)	0.7006 (3)	1.2933 (2)	0.0639 (9)
H311	0.8735	0.7201	1.3491	0.077*
C312	0.79166 (16)	0.7295 (3)	1.2699 (2)	0.0722 (10)
H312	0.7802	0.7684	1.3116	0.087*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.03822 (19)	0.04063 (17)	0.03006 (17)	-0.00220 (15)	0.00295 (12)	-0.00148 (14)
O11	0.0470 (12)	0.0593 (12)	0.0528 (12)	-0.0127 (10)	0.0146 (10)	-0.0025 (10)
O12	0.0622 (13)	0.0699 (13)	0.0446 (12)	-0.0098 (11)	0.0180 (10)	0.0091 (11)
O21	0.0599 (13)	0.0486 (11)	0.0522 (13)	0.0073 (10)	0.0140 (10)	-0.0082 (10)
O22	0.0580 (13)	0.0474 (11)	0.0584 (14)	0.0102 (10)	0.0063 (11)	0.0015 (10)
N31	0.0405 (13)	0.0400 (12)	0.0331 (12)	-0.0012 (10)	0.0078 (10)	-0.0030 (9)
N32	0.0614 (18)	0.105 (2)	0.0411 (15)	-0.0057 (16)	0.0108 (13)	-0.0143 (15)
C11	0.0491 (18)	0.0344 (14)	0.0460 (17)	0.0026 (13)	0.0207 (14)	-0.0019 (13)
C12	0.0426 (16)	0.0373 (14)	0.0431 (16)	0.0024 (12)	0.0190 (13)	-0.0015 (12)
C13	0.062 (2)	0.0643 (19)	0.052 (2)	-0.0050 (17)	0.0227 (16)	0.0023 (16)
C14	0.100 (3)	0.086 (3)	0.066 (2)	0.002 (2)	0.042 (2)	0.022 (2)
C15	0.094 (3)	0.057 (2)	0.111 (3)	-0.002 (2)	0.062 (3)	0.015 (2)
C16	0.056 (2)	0.069 (2)	0.102 (3)	-0.0149 (18)	0.031 (2)	-0.007 (2)
C17	0.0473 (19)	0.0579 (18)	0.062 (2)	-0.0040 (15)	0.0168 (16)	-0.0014 (16)
C21	0.0444 (17)	0.0416 (14)	0.0452 (17)	0.0015 (13)	0.0206 (14)	0.0010 (13)
C22	0.0479 (17)	0.0401 (14)	0.0545 (18)	0.0015 (13)	0.0270 (14)	-0.0016 (13)
C23	0.068 (2)	0.059 (2)	0.097 (3)	-0.0027 (18)	0.016 (2)	-0.0217 (19)
C24	0.103 (4)	0.063 (3)	0.163 (5)	-0.014 (2)	0.046 (3)	-0.050 (3)
C25	0.123 (4)	0.042 (2)	0.192 (6)	0.007 (2)	0.094 (4)	-0.003 (3)
C26	0.103 (3)	0.064 (2)	0.121 (4)	0.035 (2)	0.061 (3)	0.034 (2)
C27	0.071 (2)	0.061 (2)	0.062 (2)	0.0168 (17)	0.0303 (18)	0.0140 (16)

C31	0.0490 (18)	0.0643 (19)	0.0413 (16)	0.0072 (15)	0.0093 (13)	-0.0153 (14)
C32	0.0391 (16)	0.074 (2)	0.0488 (18)	0.0100 (16)	0.0043 (13)	-0.0124 (17)
C33	0.0440 (17)	0.0416 (15)	0.0331 (15)	-0.0033 (13)	0.0067 (12)	-0.0023 (12)
C34	0.0447 (17)	0.0595 (17)	0.0356 (15)	0.0031 (14)	0.0110 (13)	-0.0087 (13)
C35	0.0360 (16)	0.0582 (17)	0.0374 (16)	0.0044 (14)	0.0056 (12)	-0.0004 (14)
C36	0.0397 (17)	0.0608 (18)	0.0410 (17)	0.0010 (14)	0.0009 (13)	-0.0077 (14)
C37	0.0445 (18)	0.0638 (19)	0.0371 (16)	-0.0022 (14)	0.0045 (13)	-0.0036 (14)
C38	0.0547 (19)	0.0500 (16)	0.0324 (16)	-0.0109 (14)	0.0083 (14)	-0.0012 (13)
C39	0.059 (2)	0.077 (2)	0.0444 (18)	0.0088 (17)	0.0002 (16)	-0.0159 (16)
C310	0.062 (2)	0.099 (3)	0.062 (2)	0.005 (2)	-0.0074 (19)	-0.013 (2)
C311	0.071 (2)	0.069 (2)	0.0367 (18)	-0.0147 (19)	-0.0020 (16)	-0.0016 (16)
C312	0.079 (3)	0.096 (3)	0.0382 (18)	-0.009 (2)	0.0162 (17)	-0.0134 (18)

Geometric parameters (Å, °)

Zn1—N31	2.029 (2)	C23—C24	1.376 (5)
Zn1—O12	2.039 (2)	C23—H23	0.9300
Zn1—O21	2.0392 (19)	C24—C25	1.349 (6)
Zn1—O11	2.0407 (19)	C24—H24	0.9300
Zn1—O22	2.0580 (19)	C25—C26	1.362 (6)
Zn1—Zn1 ⁱ	2.9711 (8)	C25—H25	0.9300
O11—C11	1.258 (3)	C26—C27	1.385 (4)
O12—C11 ⁱ	1.252 (3)	C26—H26	0.9300
O21—C21	1.254 (3)	C27—H27	0.9300
O22—C21 ⁱ	1.251 (3)	C31—C32	1.379 (4)
N31—C31	1.327 (3)	C31—H31	0.9300
N31—C35	1.331 (3)	C32—C33	1.383 (4)
N32—C38	1.333 (4)	C32—H32	0.9300
N32—C312	1.350 (4)	C33—C34	1.381 (3)
C11—O12 ⁱ	1.252 (3)	C33—C36	1.471 (3)
C11—C12	1.498 (4)	C34—C35	1.372 (3)
C12—C13	1.380 (4)	C34—H34	0.9300
C12—C17	1.385 (4)	C35—H35	0.9300
C13—C14	1.372 (4)	C36—C37	1.313 (4)
C13—H13	0.9300	C36—H36	0.9300
C14—C15	1.370 (5)	C37—C38	1.468 (3)
C14—H14	0.9300	C37—H37	0.9300
C15—C16	1.375 (5)	C38—C39	1.368 (4)
C15—H15	0.9300	C39—C310	1.371 (4)
C16—C17	1.385 (4)	C39—H39	0.9300
C16—H16	0.9300	C310—C311	1.366 (5)
C17—H17	0.9300	C310—H310	0.9300
C21—O22 ⁱ	1.251 (3)	C311—C312	1.355 (4)
C21—C22	1.500 (4)	C311—H311	0.9300
C22—C23	1.375 (4)	C312—H312	0.9300
C22—C27	1.385 (4)		
N31—Zn1—O12	98.00 (8)	C24—C23—H23	120.0
N31—Zn1—O21	102.41 (8)	C22—C23—H23	120.0

supplementary materials

O12—Zn1—O21	89.34 (8)	C25—C24—C23	120.7 (4)
N31—Zn1—O11	103.00 (8)	C25—C24—H24	119.7
O12—Zn1—O11	158.97 (8)	C23—C24—H24	119.7
O21—Zn1—O11	87.31 (8)	C24—C25—C26	120.1 (4)
N31—Zn1—O22	98.62 (8)	C24—C25—H25	119.9
O12—Zn1—O22	86.52 (9)	C26—C25—H25	119.9
O21—Zn1—O22	158.93 (8)	C25—C26—C27	120.4 (4)
O11—Zn1—O22	89.19 (8)	C25—C26—H26	119.8
N31—Zn1—Zn1 ⁱ	175.50 (6)	C27—C26—H26	119.8
O12—Zn1—Zn1 ⁱ	82.26 (6)	C26—C27—C22	119.4 (3)
O21—Zn1—Zn1 ⁱ	82.08 (6)	C26—C27—H27	120.3
O11—Zn1—Zn1 ⁱ	76.71 (6)	C22—C27—H27	120.3
O22—Zn1—Zn1 ⁱ	76.89 (5)	N31—C31—C32	122.9 (3)
C11—O11—Zn1	131.38 (19)	N31—C31—H31	118.5
C11 ⁱ —O12—Zn1	124.19 (18)	C32—C31—H31	118.5
C21—O21—Zn1	124.11 (17)	C31—C32—C33	120.2 (3)
C21 ⁱ —O22—Zn1	130.32 (18)	C31—C32—H32	119.9
C31—N31—C35	116.8 (2)	C33—C32—H32	119.9
C31—N31—Zn1	121.66 (18)	C34—C33—C32	116.5 (2)
C35—N31—Zn1	121.47 (18)	C34—C33—C36	123.2 (2)
C38—N32—C312	117.7 (3)	C32—C33—C36	120.3 (3)
O12 ⁱ —C11—O11	125.1 (3)	C35—C34—C33	119.7 (3)
O12 ⁱ —C11—C12	117.5 (2)	C35—C34—H34	120.1
O11—C11—C12	117.4 (3)	C33—C34—H34	120.1
C13—C12—C17	118.4 (3)	N31—C35—C34	123.8 (3)
C13—C12—C11	120.5 (2)	N31—C35—H35	118.1
C17—C12—C11	121.1 (3)	C34—C35—H35	118.1
C14—C13—C12	120.8 (3)	C37—C36—C33	125.9 (3)
C14—C13—H13	119.6	C37—C36—H36	117.1
C12—C13—H13	119.6	C33—C36—H36	117.1
C15—C14—C13	120.8 (3)	C36—C37—C38	126.5 (3)
C15—C14—H14	119.6	C36—C37—H37	116.8
C13—C14—H14	119.6	C38—C37—H37	116.8
C14—C15—C16	119.1 (3)	N32—C38—C39	121.7 (3)
C14—C15—H15	120.4	N32—C38—C37	115.6 (3)
C16—C15—H15	120.4	C39—C38—C37	122.8 (3)
C15—C16—C17	120.4 (3)	C38—C39—C310	119.3 (3)
C15—C16—H16	119.8	C38—C39—H39	120.3
C17—C16—H16	119.8	C310—C39—H39	120.3
C16—C17—C12	120.4 (3)	C311—C310—C39	119.8 (3)
C16—C17—H17	119.8	C311—C310—H310	120.1
C12—C17—H17	119.8	C39—C310—H310	120.1
O22 ⁱ —C21—O21	125.2 (2)	C312—C311—C310	117.8 (3)
O22 ⁱ —C21—C22	117.4 (2)	C312—C311—H311	121.1
O21—C21—C22	117.3 (2)	C310—C311—H311	121.1
C23—C22—C27	119.2 (3)	N32—C312—C311	123.6 (3)
C23—C22—C21	120.0 (3)	N32—C312—H312	118.2

C27—C22—C21	120.8 (3)	C311—C312—H312	118.2
C24—C23—C22	120.1 (4)		
O12 ⁱ —C11—C12—C13	179.8 (3)	C21—C22—C27—C26	175.3 (3)
O11—C11—C12—C13	1.0 (4)	C35—N31—C31—C32	-2.1 (4)
O12 ⁱ —C11—C12—C17	1.5 (4)	N31—C31—C32—C33	1.1 (5)
O11—C11—C12—C17	-177.2 (3)	C31—C32—C33—C34	0.9 (4)
C17—C12—C13—C14	1.3 (5)	C31—C32—C33—C36	-178.9 (3)
C11—C12—C13—C14	-177.0 (3)	C32—C33—C34—C35	-1.7 (4)
C12—C13—C14—C15	-0.1 (5)	C36—C33—C34—C35	178.1 (3)
C13—C14—C15—C16	-0.5 (6)	C31—N31—C35—C34	1.3 (4)
C14—C15—C16—C17	-0.1 (6)	C33—C34—C35—N31	0.7 (4)
C15—C16—C17—C12	1.2 (5)	C34—C33—C36—C37	4.5 (5)
C13—C12—C17—C16	-1.8 (4)	C32—C33—C36—C37	-175.6 (3)
C11—C12—C17—C16	176.5 (3)	C33—C36—C37—C38	-177.6 (3)
O22 ⁱ —C21—C22—C23	-5.7 (4)	C312—N32—C38—C39	0.0 (5)
O21—C21—C22—C23	173.3 (3)	C312—N32—C38—C37	-179.4 (3)
O22 ⁱ —C21—C22—C27	176.9 (3)	C36—C37—C38—N32	175.1 (3)
O21—C21—C22—C27	-4.1 (4)	C36—C37—C38—C39	-4.4 (5)
C27—C22—C23—C24	2.9 (6)	N32—C38—C39—C310	0.2 (5)
C21—C22—C23—C24	-174.5 (4)	C37—C38—C39—C310	179.6 (3)
C22—C23—C24—C25	-1.6 (7)	C38—C39—C310—C311	-0.9 (5)
C23—C24—C25—C26	-0.5 (8)	C39—C310—C311—C312	1.3 (5)
C24—C25—C26—C27	1.4 (7)	C38—N32—C312—C311	0.4 (5)
C25—C26—C27—C22	0.0 (6)	C310—C311—C312—N32	-1.1 (5)
C23—C22—C27—C26	-2.1 (5)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

