

Bis(2,2'-bi-1*H*-imidazole- κ^2 N³,N^{3'})bis(4-methylbenzoato- κ O)copper(II)

Zhou Hui

College of Chemistry and Chemical Engineering, Henan University, Kaifeng 475001, People's Republic of China
Correspondence e-mail: zhouchui@163.com


Received 5 October 2009; accepted 24 October 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.045; wR factor = 0.104; data-to-parameter ratio = 13.3.

In title compound, $[\text{Cu}(\text{C}_8\text{H}_7\text{O}_2)_2(\text{C}_6\text{H}_6\text{N}_4)_2]$, the Cu^{II} atom (site symmetry $\bar{1}$) is coordinated by two *N,N'*-bidentate 2,2'-biimidazole ligands and two weakly bonded 4-methylbenzoate anions, resulting in a strongly elongated *trans*-CuO₂N₄ octahedral geometry. In the crystal, adjacent molecules are linked *via* pairs of N—H···O hydrogen bonds into chains propagating in [010].

Related literature

For a related structure, see: Yang *et al.* (2009).



Experimental

Crystal data

$[\text{Cu}(\text{C}_8\text{H}_7\text{O}_2)_2(\text{C}_6\text{H}_6\text{N}_4)_2]$

$M_r = 602.11$

Monoclinic, $P2_1/n$
 $a = 12.2839 (9)$ Å
 $b = 7.3150 (5)$ Å
 $c = 14.9755 (11)$ Å
 $\beta = 93.673 (1)$ °
 $V = 1342.89 (17)$ Å³

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.87$ mm⁻¹
 $T = 296$ K
 $0.25 \times 0.18 \times 0.15$ mm

Data collection

Bruker SMART APEX CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2001)
 $T_{\min} = 0.813$, $T_{\max} = 0.881$

6459 measured reflections
2611 independent reflections
1807 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.104$
 $S = 1.02$
2611 reflections
196 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.26$ e Å⁻³
 $\Delta\rho_{\min} = -0.29$ e Å⁻³

Table 1
Selected bond lengths (Å).

Cu1—N3	2.012 (2)	Cu1—O2	2.685 (2)
Cu1—N1	2.021 (2)		

Table 2
Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
N4—H4A···O1 ⁱ	0.890 (10)	1.755 (11)	2.644 (3)	178 (3)
N2—H2A···O2 ⁱ	0.894 (10)	1.755 (11)	2.647 (3)	176 (3)

Symmetry code: (i) $x, y - 1, z$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5128).

References

- Bruker (2001). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Yang, L. F., Cao, M. L., Mo, H. J., Hao, H. G., Wu, J. J., Zhang, J. P. & Ye, B. H. (2009). *CrystEngComm*, **11**, 1114–1121.

supporting information

Acta Cryst. (2009). E65, m1498 [doi:10.1107/S1600536809044262]

Bis(2,2'-bi-1*H*-imidazole- κ^2N^3,N^3')bis(4-methylbenzoato- κO)copper(II)

Zhou Hui

S1. Comment

2,2'-Biimidazole is an interesting ligand because it has two N atoms and two –NH donors. Both N-donors having the stronger coordination ability and flexible coordination modes. Additionally, two –NH donors can interact with other hydrogen bonding acceptors *via* hydrogen bonds (Yang *et al.*, 2009). Herein, we report the title compound of bis(4-methyl-benzenecarboxylate- κN)bis(2,2'-biimidazole-*N,N'*) copper, (I).


In the symmetric unit of (I), Cu²⁺ having an inversion centre was chelated by two 2,2'-biimidazole ligands through and weakly interacts with two 4-methyl benzenecarboxylate ligands acting as monodentate, which results in a elongated octahedron (Cu1—N1 2.021 (2) Å; Cu1—N3 2.012 (2) Å; Cu1—O2 2.685 (2) Å). Adjacent two Cu(C₈H₇O₂)(C₆H₃N₄) units are linked together by two pairs of N—H \cdots O hydrogen bonds forming two R₂²(9) motifs, which are further arranged into a one-dimensional chain along the [010] direction.

S2. Experimental


CuCl₂·2H₂O (0.17 g, 1.0 mmol) was added into an aqueous solution (15 ml) of 4-methyl-benzenecarboxylic acid (0.14 g, 1.0 mmol) and NaOH (0.04 g, 1.0 mmol) and refluxed for 30 min. Then an ethanol solution (10 ml) containing 2,2'-biimidazole (0.13 g, 1.0 mmol) was slowly added with continuous stirring. The resulting solution was refluxed for 3 h, filtered and kept for crystallization. After nine days, blue blocks of (I) were obtained.

S3. Refinement

H atoms bonded to N are located in a difference maps and refined isotropically 0.89 (1) for N—H using *DFIX* commands. All the remaining H atoms were positioned geometrically with C—H = 0.93 Å (aromatic) and 0.96 Å (methyl) and were refined as riding with U_{iso}(H) = 1.2U_{eq}(C) (aromatic) and 1.5U_{eq}(C) (methyl).

**Figure 1**

The molecular structure of (I), with displacement ellipsoids drawn at the 30% probability level. H-bonds are shown as dashed lines. Unlabeled atoms are related to labeled atoms by the symmetry transformation ($2 - x, 2 - y, -z$).

**Figure 2**

Part of the crystal structure of (I), showing the formation of the one-dimensional chain along the [010] direction. Weak contacts and hydrogen bonds are shown as thick and thin dashed lines, respectively. H atoms not involved in the hydrogen bonds have been omitted for the clarity.

Bis(2,2'-bi-1H-imidazole- κ^2N^3,N^3')bis(4-methylbenzoato- κO)copper(II)*Crystal data* $[\text{Cu}(\text{C}_8\text{H}_7\text{O}_2)_2(\text{C}_6\text{H}_6\text{N}_4)_2]$ $M_r = 602.11$ Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

 $a = 12.2839 (9) \text{ \AA}$ $b = 7.3150 (5) \text{ \AA}$ $c = 14.9755 (11) \text{ \AA}$ $\beta = 93.673 (1)^\circ$ $V = 1342.89 (17) \text{ \AA}^3$ $Z = 2$ $F(000) = 622$ $D_x = 1.489 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1035 reflections

 $\theta = 2.7-22.3^\circ$ $\mu = 0.87 \text{ mm}^{-1}$ $T = 296 \text{ K}$

Block, blue

 $0.25 \times 0.18 \times 0.15 \text{ mm}$ *Data collection*Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

0.3° wide ω exposures scansAbsorption correction: multi-scan
(SADABS; Bruker, 2001) $T_{\min} = 0.813$, $T_{\max} = 0.881$

6459 measured reflections

2611 independent reflections

1807 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.043$ $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 2.1^\circ$ $h = -13 \rightarrow 15$ $k = -8 \rightarrow 9$ $l = -18 \rightarrow 10$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.045$ $wR(F^2) = 0.104$ $S = 1.02$

2611 reflections

196 parameters

2 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0405P)^2 + 0.305P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.26 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.29 \text{ e \AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	1.0000	1.0000	0.0000	0.03766 (19)
N1	0.89630 (18)	0.7883 (3)	0.01223 (16)	0.0349 (6)
N2	0.8736 (2)	0.5161 (3)	0.07173 (18)	0.0397 (6)

H2A	0.895 (3)	0.410 (3)	0.096 (2)	0.067 (11)*
N3	1.09622 (18)	0.8469 (3)	0.08419 (16)	0.0347 (6)
N4	1.1064 (2)	0.5980 (4)	0.16805 (17)	0.0408 (6)
H4A	1.087 (2)	0.494 (3)	0.193 (2)	0.053 (10)*
C9	0.7954 (2)	0.7212 (4)	-0.0174 (2)	0.0417 (8)
H9	0.7454	0.7807	-0.0565	0.050*
C10	0.7809 (2)	0.5549 (4)	0.0195 (2)	0.0450 (8)
H10	0.7195	0.4809	0.0110	0.054*
C11	0.9395 (2)	0.6593 (4)	0.06569 (19)	0.0333 (7)
C12	1.0464 (2)	0.6939 (4)	0.10720 (19)	0.0342 (7)
C13	1.2008 (2)	0.6939 (4)	0.1851 (2)	0.0478 (8)
H13	1.2587	0.6610	0.2249	0.057*
C14	1.1944 (2)	0.8452 (4)	0.1337 (2)	0.0420 (8)
H14	1.2481	0.9346	0.1320	0.050*
O1	1.05155 (19)	1.2894 (3)	0.24540 (17)	0.0673 (8)
O2	0.93115 (18)	1.1925 (3)	0.13825 (16)	0.0527 (6)
C1	0.9823 (3)	1.1777 (4)	0.2138 (2)	0.0444 (8)
C2	0.9573 (2)	1.0140 (4)	0.2697 (2)	0.0379 (7)
C3	0.8761 (2)	0.8939 (4)	0.2408 (2)	0.0415 (8)
H3	0.8384	0.9127	0.1857	0.050*
C4	0.8499 (3)	0.7468 (5)	0.2923 (2)	0.0500 (9)
H4	0.7947	0.6676	0.2715	0.060*
C5	0.9039 (3)	0.7145 (5)	0.3741 (2)	0.0506 (9)
C6	0.9860 (3)	0.8328 (5)	0.4026 (2)	0.0566 (10)
H6	1.0242	0.8122	0.4573	0.068*
C7	1.0128 (3)	0.9816 (4)	0.3515 (2)	0.0507 (9)
H7	1.0683	1.0602	0.3722	0.061*
C8	0.8718 (3)	0.5549 (6)	0.4308 (3)	0.0838 (14)
H8A	0.8992	0.4436	0.4067	0.126*
H8B	0.7937	0.5485	0.4307	0.126*
H8C	0.9021	0.5710	0.4910	0.126*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0373 (3)	0.0273 (3)	0.0474 (4)	-0.0028 (2)	-0.0042 (2)	0.0112 (2)
N1	0.0373 (13)	0.0266 (13)	0.0404 (15)	0.0007 (11)	-0.0007 (11)	0.0035 (11)
N2	0.0399 (14)	0.0276 (15)	0.0512 (16)	-0.0012 (12)	-0.0003 (13)	0.0073 (12)
N3	0.0383 (13)	0.0242 (13)	0.0411 (15)	-0.0033 (11)	-0.0010 (12)	0.0036 (11)
N4	0.0411 (15)	0.0313 (15)	0.0488 (17)	-0.0016 (12)	-0.0071 (13)	0.0134 (13)
C9	0.0405 (17)	0.0315 (17)	0.052 (2)	0.0030 (14)	-0.0065 (15)	0.0054 (14)
C10	0.0370 (17)	0.0360 (18)	0.061 (2)	-0.0050 (14)	-0.0052 (16)	0.0028 (16)
C11	0.0381 (16)	0.0254 (16)	0.0364 (17)	-0.0013 (13)	0.0039 (14)	0.0040 (13)
C12	0.0375 (16)	0.0270 (16)	0.0378 (18)	0.0013 (13)	0.0007 (14)	0.0005 (13)
C13	0.0412 (18)	0.046 (2)	0.054 (2)	-0.0018 (15)	-0.0124 (16)	0.0117 (17)
C14	0.0429 (18)	0.0359 (18)	0.046 (2)	-0.0071 (14)	-0.0050 (16)	0.0042 (15)
O1	0.0665 (16)	0.0519 (16)	0.0802 (19)	-0.0196 (13)	-0.0211 (14)	0.0323 (14)
O2	0.0590 (14)	0.0433 (14)	0.0549 (15)	-0.0006 (11)	-0.0050 (12)	0.0187 (11)

C1	0.0415 (18)	0.0378 (19)	0.054 (2)	0.0051 (15)	0.0002 (17)	0.0120 (16)
C2	0.0374 (16)	0.0327 (17)	0.0437 (19)	0.0052 (14)	0.0023 (14)	0.0073 (14)
C3	0.0453 (18)	0.0395 (19)	0.0392 (19)	0.0028 (15)	-0.0010 (15)	0.0092 (15)
C4	0.0507 (19)	0.0427 (19)	0.056 (2)	-0.0127 (16)	-0.0030 (17)	0.0077 (17)
C5	0.0470 (19)	0.044 (2)	0.060 (2)	-0.0069 (16)	0.0045 (17)	0.0199 (17)
C6	0.057 (2)	0.058 (2)	0.053 (2)	-0.0043 (18)	-0.0108 (18)	0.0261 (18)
C7	0.0451 (18)	0.050 (2)	0.055 (2)	-0.0119 (16)	-0.0118 (16)	0.0160 (17)
C8	0.085 (3)	0.078 (3)	0.086 (3)	-0.029 (2)	-0.010 (2)	0.045 (3)

Geometric parameters (\AA , $^\circ$)

Cu1—N3 ⁱ	2.012 (2)	C13—C14	1.347 (4)
Cu1—N3	2.012 (2)	C13—H13	0.9300
Cu1—N1 ⁱ	2.021 (2)	C14—H14	0.9300
Cu1—N1	2.021 (2)	O1—C1	1.251 (4)
Cu1—O2	2.685 (2)	O2—C1	1.263 (4)
Cu1—O2 ⁱ	2.685 (2)	C1—C2	1.504 (4)
N1—C11	1.326 (3)	C2—C3	1.378 (4)
N1—C9	1.380 (3)	C2—C7	1.384 (4)
N2—C11	1.331 (3)	C3—C4	1.374 (4)
N2—C10	1.370 (4)	C3—H3	0.9300
N2—H2A	0.894 (10)	C4—C5	1.375 (4)
N3—C12	1.332 (3)	C4—H4	0.9300
N3—C14	1.375 (4)	C5—C6	1.376 (5)
N4—C12	1.334 (3)	C5—C8	1.510 (4)
N4—C13	1.365 (4)	C6—C7	1.382 (4)
N4—H4A	0.890 (10)	C6—H6	0.9300
C9—C10	1.353 (4)	C7—H7	0.9300
C9—H9	0.9300	C8—H8A	0.9600
C10—H10	0.9300	C8—H8B	0.9600
C11—C12	1.439 (4)	C8—H8C	0.9600
N3 ⁱ —Cu1—N3	180.0	N4—C13—H13	126.3
N3 ⁱ —Cu1—N1 ⁱ	82.25 (9)	C13—C14—N3	109.4 (3)
N3—Cu1—N1 ⁱ	97.75 (9)	C13—C14—H14	125.3
N3 ⁱ —Cu1—N1	97.75 (9)	N3—C14—H14	125.3
N3—Cu1—N1	82.25 (9)	O1—C1—O2	124.8 (3)
N1 ⁱ —Cu1—N1	180.0	O1—C1—C2	117.9 (3)
C11—N1—C9	104.9 (2)	O2—C1—C2	117.3 (3)
C11—N1—Cu1	111.68 (18)	C3—C2—C7	118.3 (3)
C9—N1—Cu1	143.3 (2)	C3—C2—C1	120.1 (3)
C11—N2—C10	106.6 (3)	C7—C2—C1	121.5 (3)
C11—N2—H2A	123 (2)	C4—C3—C2	120.9 (3)
C10—N2—H2A	129 (2)	C4—C3—H3	119.5
C12—N3—C14	104.7 (2)	C2—C3—H3	119.5
C12—N3—Cu1	111.65 (18)	C3—C4—C5	121.2 (3)
C14—N3—Cu1	143.40 (19)	C3—C4—H4	119.4
C12—N4—C13	106.5 (2)	C5—C4—H4	119.4

C12—N4—H4A	126 (2)	C4—C5—C6	118.0 (3)
C13—N4—H4A	128 (2)	C4—C5—C8	120.3 (3)
C10—C9—N1	109.2 (3)	C6—C5—C8	121.6 (3)
C10—C9—H9	125.4	C5—C6—C7	121.3 (3)
N1—C9—H9	125.4	C5—C6—H6	119.4
C9—C10—N2	107.0 (3)	C7—C6—H6	119.4
C9—C10—H10	126.5	C6—C7—C2	120.2 (3)
N2—C10—H10	126.5	C6—C7—H7	119.9
N1—C11—N2	112.3 (3)	C2—C7—H7	119.9
N1—C11—C12	117.1 (2)	C5—C8—H8A	109.5
N2—C11—C12	130.6 (3)	C5—C8—H8B	109.5
N3—C12—N4	112.1 (2)	H8A—C8—H8B	109.5
N3—C12—C11	117.2 (3)	C5—C8—H8C	109.5
N4—C12—C11	130.7 (3)	H8A—C8—H8C	109.5
C14—C13—N4	107.3 (3)	H8B—C8—H8C	109.5
C14—C13—H13	126.3		

Symmetry code: (i) $-x+2, -y+2, -z$.

Hydrogen-bond geometry (\AA , $^{\circ}$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N4—H4A ⁱⁱ —O1 ⁱⁱ	0.89 (1)	1.76 (1)	2.644 (3)	178 (3)
N2—H2A ⁱⁱ —O2 ⁱⁱ	0.89 (1)	1.76 (1)	2.647 (3)	176 (3)

Symmetry code: (ii) $x, y-1, z$.