

Di- μ -ethanolato-bis[diethanolato(2-methylquinolin-8-olato)titanium(IV)]

Yousef Fazaeli, Ezzatollah Najafi, Mostafa M. Amini and Hamid Reza Khavasi*

Department of Chemistry, Shahid Beheshti University, G. C., Evin, Tehran 1983963113, Iran

Correspondence e-mail: m-pouramini@cc.sbu.ac.ir


Received 21 October 2009; accepted 31 October 2009

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.097; wR factor = 0.197; data-to-parameter ratio = 22.2.

In the centrosymmetric dinuclear title compound, $[\text{Ti}_2(\text{C}_{10}\text{H}_8\text{NO})_2(\text{C}_2\text{H}_5\text{O})_6]$, the Ti atom is bonded to an *N,O*-bidentate quinolin-8-olate ligand, two terminal ethanolate anions and two bridging ethanolate anions in a distorted TiNO_5 octahedral geometry. An intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond occurs; in the crystal, intermolecular $\text{C}-\text{H}\cdots\text{O}$ interactions help to establish the packing.

Related literature

For Ti^{IV} -8-hydroxyquinolinates, see: Amini *et al.* (2004); Birdet *et al.* (1973); Studd & Swallow (1968). For a related structure, see: Fazaeli *et al.* (2008).



Experimental

Crystal data

$[\text{Ti}_2(\text{C}_{10}\text{H}_8\text{NO})_2(\text{C}_2\text{H}_5\text{O})_6]$
 $M_r = 682.51$
 Monoclinic, $P2_1/n$
 $a = 9.0497$ (18) Å

$b = 13.086$ (3) Å
 $c = 14.189$ (3) Å
 $\beta = 95.21$ (3)°
 $V = 1673.4$ (6) Å³

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.53$ mm⁻¹

$T = 120$ K
 $0.45 \times 0.28 \times 0.23$ mm

Data collection

Stoe IPDS II diffractometer
 Absorption correction: numerical
 (*X-SHAPE*; Stoe & Cie, 2005)
 $T_{\text{min}} = 0.686$, $T_{\text{max}} = 0.905$

12962 measured reflections
 4503 independent reflections
 3540 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.099$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.097$
 $wR(F^2) = 0.197$
 $S = 1.14$
 4503 reflections

203 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.14$ e Å⁻³

Table 1

Selected bond lengths (Å).

Ti1—N1	2.387 (3)	Ti1—O3	1.817 (3)
Ti1—O1	1.950 (3)	Ti1—O4 ⁱ	2.008 (3)
Ti1—O2	1.808 (3)	Ti1—O4	2.061 (2)

Symmetry code: (i) $-x + 2, -y + 2, -z$.

Table 2

Hydrogen-bond geometry (Å, °).

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C15—H15A \cdots O1	0.97	2.46	3.061 (5)	120
C1—H1C \cdots O3 ⁱ	0.96	2.38	3.292 (5)	159
C3—H3 \cdots O1 ⁱⁱ	0.93	2.41	3.310 (5)	163

Symmetry codes: (i) $-x + 2, -y + 2, -z$; (ii) $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: *X-AREA* (Stoe & Cie, 2005); cell refinement: *X-AREA*; data reduction: *X-RED* (Stoe & Cie, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We are grateful to Shahid Beheshti University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5165).

References

- Amini, M. M., Mirzaee, M. & Ng, S. W. (2004). *Acta Cryst.* E60, m145–m146.
 Birdet, P. H., Fraser, A. R. & Lau, C. F. (1973). *Inorg. Chem.* 12, 1322–1328.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* 30, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* 32, 837–838.
 Fazaeli, Y., Amini, M. M., Gao, S. & Ng, S. W. (2008). *Acta Cryst.* E64, o97.
 Sheldrick, G. M. (2008). *Acta Cryst.* A64, 112–122.
 Stoe & Cie (2005). *X-AREA*, *X-SHAPE* and *X-RED*. Stoe & Cie, Darmstadt, Germany.
 Studd, B. F. & Swallow, A. G. (1968). *J. Chem. Soc. A*, pp. 1961–1967.

supplementary materials

Acta Cryst. (2009). E65, m1531 [doi:10.1107/S1600536809045796]

Di- μ -ethanolato-bis[diethanolato(2-methylquinolin-8-olato)titanium(IV)]

Y. Fazaeli, E. Najafi, M. M. Amini and H. R. Khavasi

Comment

8-Hydroxyquinoline and its derivatives are reagents for the gravimetric analysis of metal ions and the crystal structures of a large number of metal 8-hydroxyquinolinates have been documented. However, for Ti^{IV} , only three 8-hydroxyquinolinates are known (Amini *et al.*, 2004), (Birdet *et al.*, 1973; Studd & Swallow, 1968). Recently, we have reported the structure of 2-methyl-8-hydroxyquinoline (Fazaeli *et al.*, 2008). In continuation our work in preparation of 8-hydroxyquinoline derivatives of transition metal elements from corresponding alkoxides, here we report synthesis and characterization of the title compound, (I).

This molecule lie across crystallographic inversion centre and the assymmetric unit therefore contains one-half of a molecule. The 8-hydroxy-2-methylquinolinate anion chelates to Ti in the triethanolate derivative; however, the coordination number is raised to six as two of the four ethanolate groups are bridging (Table 1) (Fig. 1).

Experimental

Manipulations were carried out under nitrogen, using standard Schlenk techniques. Titanium^{IV} tetraethoxide was prepared from titanium^{IV} tetrapropoxide (Fluka) and dry ethanol by the alkoxide exchange method and it was puried by vacuum distillation. 8-Hydroxyquinoline (1.6 g, 10 mmol) was added to the reagent (2.28 g, 10 mmol) in toluene (10 ml). The mixture was stirred for a day and the solvent then removed under reduced pressure to furnish a yellow solid. The solid was crystallized from a dichloromethane n-hexane mixture to give yellow prisms of (I).

Refinement

All H atoms were positioned geometrically, with C—H = 0.93 Å, 0.96 Å and 0.97 Å for aromatic, methyl and CH₂ hydrogen atoms respectively and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of (I) with displacement ellipsoids drawn at the 30% probability level. Atoms with the suffix a are generated by $(2-x, 2-y, -z)$.

Di- μ -ethanolato-bis[diethanolato(2-methylquinolin-8-olato)titanium(IV)]

Crystal data

$[\text{Ti}_2(\text{C}_{10}\text{H}_8\text{NO})_2(\text{C}_2\text{H}_5\text{O})_6]$	$F_{000} = 720$
$M_r = 682.51$	$D_x = 1.354 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 1633 reflections
$a = 9.0497 (18) \text{ \AA}$	$\theta = 2.1\text{--}29.2^\circ$
$b = 13.086 (3) \text{ \AA}$	$\mu = 0.53 \text{ mm}^{-1}$
$c = 14.189 (3) \text{ \AA}$	$T = 120 \text{ K}$
$\beta = 95.21 (3)^\circ$	Prism, yellow
$V = 1673.4 (6) \text{ \AA}^3$	$0.45 \times 0.28 \times 0.23 \text{ mm}$
$Z = 2$	

Data collection

Stoe IPDS II diffractometer	4503 independent reflections
Radiation source: fine-focus sealed tube	3540 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.099$
Detector resolution: $0.15 \text{ mm pixels mm}^{-1}$	$\theta_{\text{max}} = 29.2^\circ$
$T = 120 \text{ K}$	$\theta_{\text{min}} = 2.1^\circ$
rotation method scans	$h = -12 \rightarrow 12$
Absorption correction: numerical (X-SHAPE; Stoe & Cie, 2005)	$k = -17 \rightarrow 17$
$T_{\text{min}} = 0.686$, $T_{\text{max}} = 0.905$	$l = -19 \rightarrow 15$
12962 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.097$	H-atom parameters constrained
$wR(F^2) = 0.197$	$w = 1/[\sigma^2(F_o^2) + (0.1579P)^2 + 0.3709P]$
$S = 1.14$	where $P = (F_o^2 + 2F_c^2)/3$
4503 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
203 parameters	$\Delta\rho_{\text{max}} = 1.26 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -1.14 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7281 (4)	0.8994 (3)	-0.1775 (3)	0.0353 (7)
H1A	0.6738	0.9058	-0.1227	0.053*
H1B	0.6598	0.8914	-0.2329	0.053*
H1C	0.7869	0.9596	-0.1839	0.053*
C2	0.8274 (3)	0.8079 (3)	-0.1667 (2)	0.0306 (7)
C3	0.8266 (4)	0.7345 (3)	-0.2396 (3)	0.0354 (7)
H3	0.7650	0.7436	-0.2950	0.042*
C4	0.9155 (4)	0.6503 (3)	-0.2294 (3)	0.0349 (7)
H4	0.9165	0.6032	-0.2784	0.042*
C5	1.0064 (4)	0.6351 (3)	-0.1439 (2)	0.0310 (6)
C6	1.0989 (4)	0.5492 (3)	-0.1246 (3)	0.0356 (7)
H6	1.1049	0.4985	-0.1700	0.043*
C7	1.1805 (4)	0.5410 (3)	-0.0376 (3)	0.0380 (8)
H7	1.2414	0.4844	-0.0252	0.046*
C8	1.1734 (4)	0.6166 (3)	0.0329 (3)	0.0349 (7)
H8	1.2298	0.6094	0.0907	0.042*
C9	1.0831 (4)	0.7011 (3)	0.0164 (2)	0.0305 (7)
C10	0.9996 (3)	0.7110 (3)	-0.0735 (2)	0.0284 (6)
C11	0.6511 (5)	0.7812 (4)	0.0572 (4)	0.0516 (11)
H11A	0.6560	0.7490	-0.0040	0.062*
H11B	0.5557	0.8149	0.0573	0.062*
C12	0.6658 (9)	0.7025 (4)	0.1327 (6)	0.081 (2)
H12A	0.7618	0.6710	0.1340	0.122*
H12B	0.5903	0.6515	0.1202	0.122*
H12C	0.6549	0.7340	0.1928	0.122*
C13	0.9606 (5)	0.8822 (4)	0.2620 (3)	0.0431 (9)
H13A	0.9889	0.8109	0.2585	0.052*
H13B	0.8540	0.8853	0.2650	0.052*
C14	1.0373 (5)	0.9294 (4)	0.3496 (3)	0.0512 (11)
H14A	1.1427	0.9274	0.3462	0.077*
H14B	1.0125	0.8918	0.4042	0.077*
H14C	1.0057	0.9990	0.3547	0.077*
C15	1.2662 (4)	0.9251 (3)	-0.0154 (3)	0.0347 (7)

supplementary materials

H15A	1.2825	0.8655	0.0249	0.042*
H15B	1.3421	0.9750	0.0042	0.042*
C16	1.2818 (4)	0.8944 (3)	-0.1167 (3)	0.0444 (9)
H16A	1.2054	0.8464	-0.1371	0.067*
H16B	1.3772	0.8636	-0.1210	0.067*
H16C	1.2728	0.9539	-0.1565	0.067*
N1	0.9147 (3)	0.7980 (2)	-0.0847 (2)	0.0284 (6)
O1	1.0681 (3)	0.7744 (2)	0.07988 (17)	0.0338 (5)
O2	0.7640 (3)	0.8529 (2)	0.07192 (19)	0.0356 (6)
O3	0.9989 (3)	0.9345 (2)	0.17872 (19)	0.0364 (6)
O4	1.1230 (2)	0.96763 (19)	-0.00251 (17)	0.0302 (5)
Ti1	0.94929 (6)	0.89765 (5)	0.05650 (4)	0.0280 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0283 (15)	0.0410 (18)	0.0341 (17)	0.0002 (13)	-0.0110 (13)	0.0000 (13)
C2	0.0241 (13)	0.0379 (16)	0.0286 (15)	-0.0053 (12)	-0.0048 (11)	0.0032 (12)
C3	0.0296 (15)	0.0430 (18)	0.0320 (17)	-0.0047 (13)	-0.0059 (12)	0.0010 (14)
C4	0.0361 (16)	0.0386 (18)	0.0288 (16)	-0.0035 (14)	-0.0032 (13)	-0.0046 (13)
C5	0.0260 (13)	0.0359 (16)	0.0312 (16)	-0.0044 (12)	0.0024 (11)	0.0014 (13)
C6	0.0354 (16)	0.0327 (16)	0.0387 (18)	0.0010 (13)	0.0034 (14)	-0.0022 (14)
C7	0.0332 (16)	0.0360 (17)	0.044 (2)	0.0052 (14)	0.0013 (14)	0.0007 (15)
C8	0.0311 (15)	0.0390 (18)	0.0331 (17)	0.0036 (13)	-0.0055 (13)	0.0010 (13)
C9	0.0254 (13)	0.0358 (17)	0.0293 (16)	-0.0001 (12)	-0.0031 (11)	0.0016 (12)
C10	0.0237 (13)	0.0350 (16)	0.0258 (14)	-0.0027 (12)	-0.0007 (11)	-0.0008 (12)
C11	0.0382 (19)	0.061 (3)	0.056 (3)	-0.0186 (19)	0.0044 (18)	-0.009 (2)
C12	0.112 (5)	0.039 (3)	0.101 (5)	-0.006 (3)	0.055 (4)	0.002 (3)
C13	0.0396 (18)	0.060 (2)	0.0292 (17)	-0.0043 (17)	0.0005 (14)	0.0001 (16)
C14	0.045 (2)	0.076 (3)	0.0309 (19)	0.013 (2)	-0.0030 (16)	-0.0061 (19)
C15	0.0218 (13)	0.0404 (18)	0.0420 (19)	0.0031 (12)	0.0028 (12)	-0.0011 (14)
C16	0.0322 (17)	0.056 (2)	0.046 (2)	-0.0024 (16)	0.0101 (16)	-0.0103 (18)
N1	0.0224 (11)	0.0346 (14)	0.0269 (13)	-0.0025 (10)	-0.0045 (10)	0.0011 (10)
O1	0.0329 (11)	0.0380 (13)	0.0287 (12)	0.0023 (10)	-0.0073 (9)	-0.0014 (9)
O2	0.0270 (11)	0.0408 (14)	0.0385 (13)	-0.0043 (10)	0.0009 (9)	-0.0014 (10)
O3	0.0329 (12)	0.0463 (15)	0.0295 (12)	-0.0052 (10)	0.0000 (9)	-0.0013 (10)
O4	0.0217 (9)	0.0372 (12)	0.0313 (12)	0.0014 (9)	0.0008 (8)	-0.0026 (9)
Ti1	0.0224 (3)	0.0344 (4)	0.0264 (3)	-0.0008 (2)	-0.0020 (2)	-0.0018 (2)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.496 (5)	C12—H12A	0.9600
C1—H1A	0.9600	C12—H12B	0.9600
C1—H1B	0.9600	C12—H12C	0.9600
C1—H1C	0.9600	C13—O3	1.435 (5)
C2—N1	1.352 (4)	C13—C14	1.501 (6)
C2—C3	1.411 (5)	C13—H13A	0.9700
C3—C4	1.364 (5)	C13—H13B	0.9700
C3—H3	0.9300	C14—H14A	0.9600

C4—C5	1.417 (5)	C14—H14B	0.9600
C4—H4	0.9300	C14—H14C	0.9600
C5—C6	1.414 (5)	C15—O4	1.438 (4)
C5—C10	1.414 (5)	C15—C16	1.512 (6)
C6—C7	1.384 (5)	C15—H15A	0.9700
C6—H6	0.9300	C15—H15B	0.9700
C7—C8	1.412 (5)	C16—H16A	0.9600
C7—H7	0.9300	C16—H16B	0.9600
C8—C9	1.383 (5)	C16—H16C	0.9600
C8—H8	0.9300	Ti1—N1	2.387 (3)
C9—O1	1.331 (4)	Ti1—O1	1.950 (3)
C9—C10	1.428 (4)	Ti1—O2	1.808 (3)
C10—N1	1.375 (4)	Ti1—O3	1.817 (3)
C11—O2	1.389 (5)	Ti1—O4 ⁱ	2.008 (3)
C11—C12	1.483 (9)	Ti1—O4	2.061 (2)
C11—H11A	0.9700	Ti1—Ti1 ⁱ	3.2948 (13)
C11—H11B	0.9700		
C2—C1—H1A	109.5	C14—C13—H13A	109.4
C2—C1—H1B	109.5	O3—C13—H13B	109.4
H1A—C1—H1B	109.5	C14—C13—H13B	109.4
C2—C1—H1C	109.5	H13A—C13—H13B	108.0
H1A—C1—H1C	109.5	C13—C14—H14A	109.5
H1B—C1—H1C	109.5	C13—C14—H14B	109.5
N1—C2—C3	121.9 (3)	H14A—C14—H14B	109.5
N1—C2—C1	117.6 (3)	C13—C14—H14C	109.5
C3—C2—C1	120.4 (3)	H14A—C14—H14C	109.5
C4—C3—C2	120.6 (3)	H14B—C14—H14C	109.5
C4—C3—H3	119.7	O4—C15—C16	112.8 (3)
C2—C3—H3	119.7	O4—C15—H15A	109.0
C3—C4—C5	119.6 (3)	C16—C15—H15A	109.0
C3—C4—H4	120.2	O4—C15—H15B	109.0
C5—C4—H4	120.2	C16—C15—H15B	109.0
C6—C5—C10	119.0 (3)	H15A—C15—H15B	107.8
C6—C5—C4	124.4 (3)	C15—C16—H16A	109.5
C10—C5—C4	116.6 (3)	C15—C16—H16B	109.5
C7—C6—C5	119.5 (3)	H16A—C16—H16B	109.5
C7—C6—H6	120.2	C15—C16—H16C	109.5
C5—C6—H6	120.2	H16A—C16—H16C	109.5
C6—C7—C8	121.5 (3)	H16B—C16—H16C	109.5
C6—C7—H7	119.2	C2—N1—C10	117.1 (3)
C8—C7—H7	119.2	C2—N1—Ti1	133.8 (2)
C9—C8—C7	120.3 (3)	C10—N1—Ti1	109.0 (2)
C9—C8—H8	119.9	C9—O1—Ti1	124.7 (2)
C7—C8—H8	119.9	C11—O2—Ti1	151.4 (3)
O1—C9—C8	123.8 (3)	C13—O3—Ti1	127.0 (2)
O1—C9—C10	117.4 (3)	C15—O4—Ti1 ⁱ	123.9 (2)
C8—C9—C10	118.7 (3)	C15—O4—Ti1	127.5 (2)
N1—C10—C5	124.0 (3)	Ti1 ⁱ —O4—Ti1	108.13 (10)

supplementary materials

N1—C10—C9	115.0 (3)	O2—Ti1—O3	97.01 (12)
C5—C10—C9	120.9 (3)	O2—Ti1—O1	102.40 (12)
O2—C11—C12	110.1 (5)	O3—Ti1—O1	88.43 (12)
O2—C11—H11A	109.6	O2—Ti1—O4 ⁱ	93.27 (11)
C12—C11—H11A	109.6	O3—Ti1—O4 ⁱ	100.06 (12)
O2—C11—H11B	109.6	O1—Ti1—O4 ⁱ	161.18 (11)
C12—C11—H11B	109.6	O2—Ti1—O4	160.52 (12)
H11A—C11—H11B	108.2	O3—Ti1—O4	97.94 (11)
C11—C12—H12A	109.5	O1—Ti1—O4	90.44 (10)
C11—C12—H12B	109.5	O4 ⁱ —Ti1—O4	71.87 (10)
H12A—C12—H12B	109.5	O2—Ti1—N1	82.66 (11)
C11—C12—H12C	109.5	O3—Ti1—N1	161.53 (12)
H12A—C12—H12C	109.5	O1—Ti1—N1	73.70 (10)
H12B—C12—H12C	109.5	O4 ⁱ —Ti1—N1	98.39 (10)
O3—C13—C14	111.0 (4)	O4—Ti1—N1	87.05 (10)
O3—C13—H13A	109.4		
N1—C2—C3—C4	-0.2 (5)	C11—O2—Ti1—N1	27.3 (6)
C1—C2—C3—C4	-179.1 (3)	C11—O2—Ti1—Ti1 ⁱ	115.5 (6)
C2—C3—C4—C5	1.7 (5)	C13—O3—Ti1—O2	38.1 (3)
C3—C4—C5—C6	177.8 (3)	C13—O3—Ti1—O1	-64.2 (3)
C3—C4—C5—C10	-0.7 (5)	C13—O3—Ti1—O4 ⁱ	132.7 (3)
C10—C5—C6—C7	0.1 (5)	C13—O3—Ti1—O4	-154.4 (3)
C4—C5—C6—C7	-178.5 (4)	C13—O3—Ti1—N1	-49.7 (5)
C5—C6—C7—C8	0.2 (6)	C13—O3—Ti1—Ti1 ⁱ	169.8 (3)
C6—C7—C8—C9	0.3 (6)	C9—O1—Ti1—O2	81.3 (3)
C7—C8—C9—O1	178.5 (3)	C9—O1—Ti1—O3	178.2 (3)
C7—C8—C9—C10	-1.0 (5)	C9—O1—Ti1—O4 ⁱ	-64.3 (4)
C6—C5—C10—N1	179.4 (3)	C9—O1—Ti1—O4	-83.9 (3)
C4—C5—C10—N1	-1.9 (5)	C9—O1—Ti1—N1	2.9 (2)
C6—C5—C10—C9	-0.8 (5)	C9—O1—Ti1—Ti1 ⁱ	-79.3 (3)
C4—C5—C10—C9	177.9 (3)	C15—O4—Ti1—O2	-130.7 (4)
O1—C9—C10—N1	1.5 (4)	Ti1 ⁱ —O4—Ti1—O2	41.8 (4)
C8—C9—C10—N1	-178.9 (3)	C15—O4—Ti1—O3	89.5 (3)
O1—C9—C10—C5	-178.3 (3)	Ti1 ⁱ —O4—Ti1—O3	-98.04 (13)
C8—C9—C10—C5	1.3 (5)	C15—O4—Ti1—O1	1.0 (3)
C3—C2—N1—C10	-2.3 (5)	Ti1 ⁱ —O4—Ti1—O1	173.48 (12)
C1—C2—N1—C10	176.6 (3)	C15—O4—Ti1—O4 ⁱ	-172.5 (3)
C3—C2—N1—Ti1	-178.5 (2)	Ti1 ⁱ —O4—Ti1—O4 ⁱ	0.0
C1—C2—N1—Ti1	0.5 (5)	C15—O4—Ti1—N1	-72.7 (3)
C5—C10—N1—C2	3.4 (5)	Ti1 ⁱ —O4—Ti1—N1	99.83 (12)
C9—C10—N1—C2	-176.4 (3)	C15—O4—Ti1—Ti1 ⁱ	-172.5 (3)
C5—C10—N1—Ti1	-179.5 (3)	C2—N1—Ti1—O2	69.3 (3)
C9—C10—N1—Ti1	0.7 (3)	C10—N1—Ti1—O2	-107.0 (2)
C8—C9—O1—Ti1	176.8 (3)	C2—N1—Ti1—O3	159.5 (3)
C10—C9—O1—Ti1	-3.6 (4)	C10—N1—Ti1—O3	-16.8 (4)

C12—C11—O2—Ti1	88.3 (7)	C2—N1—Ti1—O1	174.6 (3)
C14—C13—O3—Ti1	173.2 (3)	C10—N1—Ti1—O1	-1.8 (2)
C16—C15—O4—Ti1 ⁱ	-67.7 (4)	C2—N1—Ti1—O4 ⁱ	-22.9 (3)
C16—C15—O4—Ti1	103.7 (3)	C10—N1—Ti1—O4 ⁱ	160.7 (2)
C11—O2—Ti1—O3	-134.1 (6)	C2—N1—Ti1—O4	-94.1 (3)
C11—O2—Ti1—O1	-44.2 (6)	C10—N1—Ti1—O4	89.6 (2)
C11—O2—Ti1—O4 ⁱ	125.3 (6)	C2—N1—Ti1—Ti1 ⁱ	-59.2 (3)
C11—O2—Ti1—O4	85.9 (7)	C10—N1—Ti1—Ti1 ⁱ	124.4 (2)

Symmetry codes: (i) $-x+2, -y+2, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C15—H15A \cdots O1	0.97	2.46	3.061 (5)	120
C1—H1C \cdots O3 ⁱ	0.96	2.38	3.292 (5)	159
C3—H3 \cdots O1 ⁱⁱ	0.93	2.41	3.310 (5)	163

Symmetry codes: (i) $-x+2, -y+2, -z$; (ii) $x-1/2, -y+3/2, z-1/2$.

