

(E)-Benzyl 3-(3-nitrobenzylidene)dithiocarbazate

Huan-Qiu Li, Yin Luo, Xuan Qin and Hai-Liang Zhu*

State Key Laboratory of Pharmaceutical Biotechnology, Nanjing University, Nanjing 210093, People's Republic of China

Correspondence e-mail: hailiang_zhu@163.com


Received 30 October 2009; accepted 30 October 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$; R factor = 0.045; wR factor = 0.106; data-to-parameter ratio = 13.2.

In the title compound, $\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_2\text{S}_2$, the dihedral angle between the aromatic rings is $87.8(2)^\circ$. In the crystal, inversion dimers occur linked by pairs of $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds.

Related literature

For background to carbodithioates, see: Tarafder *et al.* (2002).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_2\text{S}_2$
 $M_r = 331.40$

Monoclinic, $P2_1/c$
 $a = 5.2175(10)\text{ \AA}$

$b = 26.213(5)\text{ \AA}$
 $c = 11.887(2)\text{ \AA}$
 $\beta = 90.67(3)^\circ$
 $V = 1625.6(6)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.34\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.30 \times 0.30 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4
diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.906$, $T_{\max} = 0.967$
9486 measured reflections

2788 independent reflections
951 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.097$
200 standard reflections
every 3 reflections
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.106$
 $S = 0.73$
2788 reflections
211 parameters

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\max} = 0.15\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.16\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots\text{A}$	$D\cdots\text{A}$	$D-\text{H}\cdots\text{A}$
$\text{N}1-\text{H}1\text{A}\cdots\text{S}2^i$	0.93 (3)	2.45 (4)	3.365 (4)	170 (3)

Symmetry code: (i) $-x + 2, -y + 1, -z + 1$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was financed by a grant from the National Natural Science Foundation of China (project 30772627) and the China Postdoctoral Science Foundation (project 20080441043).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5203).

References

- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tarafder, M. T. H., Chew, K.-B., Crouse, K. C., Ali, A. M., Yamin, B. M. & Fun, H.-K. (2002). *Polyhedron*, **21**, 2683–2690.

supporting information

Acta Cryst. (2009). E65, o3080 [doi:10.1107/S1600536809045723]

(E)-Benzyl 3-(3-nitrobenzylidene)dithiocarbazate

Huan-Qiu Li, Yin Luo, Xuan Qin and Hai-Liang Zhu

S1. Experimental

The title compound was prepared by stirring a mixture of benzyl hydrazinecarbodithioate (396 mg, 2 mmol), 3-nitrobenzaldehyde (302 mg, 2 mmol) in methanol (10 ml) for 1 h. After keeping the filtrate in air for 7 d, yellow blocks of (I) were formed.

S2. Refinement

The N-bound N atom was located in a difference map and freely refined. The other H atoms were positioned geometrically (C—H = 0.93 Å for the aromatic H atoms and C—H = 0.96 Å for the aliphatic H atoms) and were refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$.




Figure 1

The structure of the title compound (I) showing 50% displacement ellipsoids.

(E)-Benzyl 3-(3-nitrobenzylidene)dithiocarbazate

Crystal data

$\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_2\text{S}_2$
 $M_r = 331.40$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 5.2175 (10)$ Å
 $b = 26.213 (5)$ Å
 $c = 11.887 (2)$ Å

$\beta = 90.67 (3)^\circ$
 $V = 1625.6 (6)$ Å³
 $Z = 4$
 $F(000) = 688$
 $D_x = 1.354 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 25 reflections

$\theta = 9\text{--}12^\circ$
 $\mu = 0.34 \text{ mm}^{-1}$
 $T = 293 \text{ K}$

Block, yellow
 $0.30 \times 0.30 \times 0.10 \text{ mm}$

Data collection

Enraf–Nonius CAD-4
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 $\omega/2\theta$ scans
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.906$, $T_{\max} = 0.967$
9486 measured reflections

2788 independent reflections
951 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.097$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.9^\circ$
 $h = -6 \rightarrow 5$
 $k = -31 \rightarrow 29$
 $l = -14 \rightarrow 14$
200 standard reflections every 3 reflections
intensity decay: 1%

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.106$
 $S = 0.73$
2788 reflections
211 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0448P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.5564 (10)	0.38939 (18)	1.0058 (4)	0.1065 (16)
H1	1.6613	0.3787	0.9477	0.128*
C2	1.5833 (10)	0.36729 (18)	1.1102 (5)	0.1215 (18)
H2	1.7070	0.3423	1.1222	0.146*
C3	1.4312 (11)	0.38176 (19)	1.1951 (4)	0.0986 (14)
H3	1.4491	0.3668	1.2658	0.118*
C4	1.2549 (11)	0.4177 (2)	1.1770 (4)	0.1203 (18)
H4	1.1474	0.4275	1.2349	0.144*
C5	1.2307 (9)	0.44050 (17)	1.0726 (5)	0.1170 (17)
H5	1.1095	0.4661	1.0616	0.140*
C6	1.3817 (9)	0.42606 (17)	0.9862 (4)	0.0740 (11)
C7	1.3604 (11)	0.4511 (2)	0.8717 (4)	0.0893 (15)

C8	1.0763 (8)	0.45488 (13)	0.6740 (3)	0.0662 (10)
C9	0.5747 (8)	0.38503 (16)	0.5578 (3)	0.0740 (12)
H9	0.5412	0.4063	0.4967	0.089*
C10	0.4214 (8)	0.33846 (15)	0.5727 (3)	0.0648 (10)
C11	0.2307 (8)	0.32751 (16)	0.4958 (3)	0.0716 (11)
H11	0.1940	0.3499	0.4371	0.086*
C12	0.0946 (8)	0.28268 (18)	0.5071 (4)	0.0763 (12)
C13	0.1352 (9)	0.24961 (16)	0.5946 (4)	0.0997 (15)
H13	0.0389	0.2199	0.6014	0.120*
C14	0.3244 (9)	0.26178 (18)	0.6727 (4)	0.1020 (15)
H14	0.3563	0.2399	0.7329	0.122*
C15	0.4658 (8)	0.30577 (16)	0.6623 (3)	0.0838 (12)
H15	0.5917	0.3136	0.7156	0.101*
H1A	0.851 (7)	0.4589 (13)	0.540 (3)	0.100 (15)*
N1	0.8928 (7)	0.43995 (14)	0.6035 (3)	0.0779 (10)
N2	0.7523 (7)	0.39663 (13)	0.6274 (3)	0.0739 (9)
N3	-0.1075 (8)	0.27046 (17)	0.4218 (4)	0.0953 (12)
O1	-0.1536 (6)	0.30218 (13)	0.3506 (3)	0.1166 (11)
O2	-0.2120 (8)	0.22932 (15)	0.4285 (3)	0.1572 (16)
S1	1.1060 (2)	0.41799 (4)	0.79464 (9)	0.0806 (4)
S2	1.2592 (2)	0.50568 (4)	0.64490 (9)	0.0898 (4)
H7A	1.508 (7)	0.4456 (14)	0.827 (3)	0.109 (17)*
H7B	1.308 (8)	0.4842 (16)	0.871 (3)	0.14 (2)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.121 (4)	0.107 (4)	0.092 (4)	0.028 (3)	0.039 (3)	0.019 (3)
C2	0.128 (4)	0.128 (4)	0.109 (4)	0.051 (3)	0.027 (4)	0.034 (4)
C3	0.108 (4)	0.102 (4)	0.085 (4)	0.000 (3)	-0.004 (3)	0.005 (3)
C4	0.135 (5)	0.142 (5)	0.084 (4)	0.029 (4)	0.022 (3)	-0.002 (3)
C5	0.123 (4)	0.132 (4)	0.097 (4)	0.060 (3)	0.015 (3)	0.008 (3)
C6	0.070 (3)	0.076 (3)	0.076 (3)	-0.005 (2)	-0.005 (3)	-0.001 (3)
C7	0.077 (4)	0.109 (5)	0.082 (4)	-0.009 (3)	-0.010 (3)	0.016 (3)
C8	0.063 (3)	0.068 (3)	0.068 (3)	-0.002 (2)	0.001 (2)	0.007 (2)
C9	0.080 (3)	0.078 (3)	0.064 (3)	-0.005 (3)	0.002 (3)	0.005 (2)
C10	0.063 (3)	0.063 (3)	0.068 (3)	-0.006 (2)	0.006 (2)	0.000 (2)
C11	0.068 (3)	0.079 (3)	0.067 (3)	-0.007 (3)	0.003 (3)	0.005 (2)
C12	0.070 (3)	0.089 (4)	0.069 (3)	-0.012 (3)	-0.002 (3)	0.000 (3)
C13	0.103 (4)	0.089 (3)	0.107 (4)	-0.033 (3)	-0.017 (3)	0.027 (3)
C14	0.111 (4)	0.087 (4)	0.108 (4)	-0.020 (3)	-0.015 (3)	0.035 (3)
C15	0.094 (3)	0.078 (3)	0.079 (3)	-0.006 (3)	-0.013 (2)	0.012 (3)
N1	0.080 (3)	0.079 (3)	0.075 (3)	-0.010 (2)	-0.008 (2)	0.015 (2)
N2	0.076 (3)	0.068 (2)	0.077 (2)	-0.005 (2)	0.002 (2)	0.0053 (19)
N3	0.096 (3)	0.097 (4)	0.094 (3)	-0.010 (3)	-0.005 (3)	0.000 (3)
O1	0.114 (3)	0.127 (3)	0.108 (2)	-0.019 (2)	-0.031 (2)	0.016 (2)
O2	0.180 (4)	0.141 (3)	0.150 (3)	-0.082 (3)	-0.045 (3)	0.017 (3)
S1	0.0798 (8)	0.0879 (8)	0.0741 (7)	-0.0113 (6)	0.0010 (6)	0.0101 (6)

S2	0.0950 (9)	0.0826 (8)	0.0917 (8)	-0.0215 (7)	-0.0063 (6)	0.0142 (6)
----	------------	------------	------------	-------------	-------------	------------

Geometric parameters (\AA , $\text{^{\circ}}$)

C1—C6	1.343 (5)	C9—N2	1.272 (4)
C1—C2	1.375 (5)	C9—C10	1.471 (5)
C1—H1	0.9300	C9—H9	0.9300
C2—C3	1.346 (5)	C10—C11	1.373 (4)
C2—H2	0.9300	C10—C15	1.384 (5)
C3—C4	1.332 (5)	C11—C12	1.380 (5)
C3—H3	0.9300	C11—H11	0.9300
C4—C5	1.381 (5)	C12—C13	1.368 (5)
C4—H4	0.9300	C12—N3	1.489 (5)
C5—C6	1.355 (5)	C13—C14	1.384 (5)
C5—H5	0.9300	C13—H13	0.9300
C6—C7	1.514 (5)	C14—C15	1.375 (5)
C7—S1	1.823 (5)	C14—H14	0.9300
C7—H7A	0.95 (4)	C15—H15	0.9300
C7—H7B	0.91 (4)	N1—N2	1.383 (4)
C8—N1	1.324 (4)	N1—H1A	0.93 (3)
C8—S2	1.676 (4)	N3—O1	1.208 (4)
C8—S1	1.735 (4)	N3—O2	1.212 (4)
C6—C1—C2	121.3 (4)	N2—C9—H9	119.6
C6—C1—H1	119.3	C10—C9—H9	119.6
C2—C1—H1	119.3	C11—C10—C15	119.7 (4)
C3—C2—C1	120.2 (5)	C11—C10—C9	118.9 (4)
C3—C2—H2	119.9	C15—C10—C9	121.3 (4)
C1—C2—H2	119.9	C10—C11—C12	118.9 (4)
C4—C3—C2	119.3 (5)	C10—C11—H11	120.6
C4—C3—H3	120.3	C12—C11—H11	120.6
C2—C3—H3	120.3	C13—C12—C11	122.5 (4)
C3—C4—C5	120.5 (5)	C13—C12—N3	118.9 (4)
C3—C4—H4	119.8	C11—C12—N3	118.5 (4)
C5—C4—H4	119.8	C12—C13—C14	117.8 (4)
C6—C5—C4	120.8 (4)	C12—C13—H13	121.1
C6—C5—H5	119.6	C14—C13—H13	121.1
C4—C5—H5	119.6	C15—C14—C13	120.8 (4)
C1—C6—C5	117.9 (4)	C15—C14—H14	119.6
C1—C6—C7	120.6 (5)	C13—C14—H14	119.6
C5—C6—C7	121.6 (5)	C14—C15—C10	120.2 (4)
C6—C7—S1	106.9 (3)	C14—C15—H15	119.9
C6—C7—H7A	113 (2)	C10—C15—H15	119.9
S1—C7—H7A	104 (2)	C8—N1—N2	119.6 (4)
C6—C7—H7B	116 (3)	C8—N1—H1A	121 (2)
S1—C7—H7B	104 (3)	N2—N1—H1A	119 (2)
H7A—C7—H7B	113 (4)	C9—N2—N1	116.5 (4)
N1—C8—S2	120.9 (3)	O1—N3—O2	124.9 (5)

N1—C8—S1	114.5 (3)	O1—N3—C12	117.6 (4)
S2—C8—S1	124.6 (3)	O2—N3—C12	117.5 (5)
N2—C9—C10	120.9 (4)	C8—S1—C7	102.0 (2)
C6—C1—C2—C3	0.8 (8)	N3—C12—C13—C14	179.9 (4)
C1—C2—C3—C4	-0.1 (8)	C12—C13—C14—C15	0.1 (7)
C2—C3—C4—C5	-1.0 (8)	C13—C14—C15—C10	-0.4 (6)
C3—C4—C5—C6	1.4 (8)	C11—C10—C15—C14	1.7 (6)
C2—C1—C6—C5	-0.4 (7)	C9—C10—C15—C14	-178.3 (3)
C2—C1—C6—C7	178.1 (4)	S2—C8—N1—N2	177.2 (2)
C4—C5—C6—C1	-0.7 (7)	S1—C8—N1—N2	-2.9 (4)
C4—C5—C6—C7	-179.2 (4)	C10—C9—N2—N1	177.3 (3)
C1—C6—C7—S1	96.1 (5)	C8—N1—N2—C9	178.3 (3)
C5—C6—C7—S1	-85.4 (5)	C13—C12—N3—O1	174.2 (4)
N2—C9—C10—C11	179.7 (3)	C11—C12—N3—O1	-4.7 (6)
N2—C9—C10—C15	-0.3 (5)	C13—C12—N3—O2	-6.3 (6)
C15—C10—C11—C12	-2.8 (5)	C11—C12—N3—O2	174.8 (4)
C9—C10—C11—C12	177.2 (3)	N1—C8—S1—C7	-177.5 (3)
C10—C11—C12—C13	2.6 (6)	S2—C8—S1—C7	2.4 (3)
C10—C11—C12—N3	-178.5 (3)	C6—C7—S1—C8	173.1 (4)
C11—C12—C13—C14	-1.3 (6)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1A···S2 ⁱ	0.93 (3)	2.45 (4)	3.365 (4)	170 (3)

Symmetry code: (i) $-x+2, -y+1, -z+1$.