

(3*R*,4*R*,5*S*)-4-Hydroxy-3-methyl-5-[(2*S*,3*R*)-3-methylpent-4-en-2-yl]-4,5-dihydrofuran-2(3*H*)-one

Annika Gille, Markus Schürmann, Hans Preut* and Martin Hiersemann

Fakultät Chemie, Technische Universität Dortmund, Otto-Hahn-Strasse 6, 44221 Dortmund, Germany
Correspondence e-mail: hans.preut@udo.edu


Received 18 November 2009; accepted 23 November 2009

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.035; wR factor = 0.048; data-to-parameter ratio = 8.6.

The relative configuration of the title compound, $\text{C}_{11}\text{H}_{18}\text{O}_3$, which was synthesized using a catalytic asymmetric Gosteli–Claisen rearrangement, a diastereoselective reduction with K-Selectride and an Evans aldol addition, was corroborated by single-crystal X-ray diffraction analysis. The five-membered ring has an envelope conformation with a dihedral angle of $29.46(16)^\circ$ between the coplanar part and the flap (the hydroxy-bearing ring C atom). In the crystal, molecules are connected via bifurcated $\text{O}-\text{H}\cdots(\text{O},\text{O})$ hydrogen bonds, generating [010] chains.

Related literature

For further synthetic details, see: Abraham *et al.* (2001, 2004); Brown (1973); Evans *et al.* (1981, 1999); Otera *et al.* (1992). For the structure of the major diastereoisomer arising from the same reaction, see: Gille *et al.* (2008).



Experimental

Crystal data



$M_r = 198.25$

Monoclinic, $P2_1$
 $a = 7.7265(10)\text{ \AA}$
 $b = 6.4798(8)\text{ \AA}$
 $c = 11.0598(16)\text{ \AA}$
 $\beta = 92.563(14)^\circ$
 $V = 553.17(13)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.50 \times 0.18 \times 0.04\text{ mm}$

Data collection

Oxford Diffraction Xcalibur S CCD diffractometer
Absorption correction: none
3255 measured reflections

1129 independent reflections
737 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.040$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.048$
 $S = 0.99$
1129 reflections
131 parameters

1 restraint
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.14\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.15\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}3-\text{H}3\cdots\text{O}1^i$	0.84	2.52	3.023 (2)	120
$\text{O}3-\text{H}3\cdots\text{O}2^i$	0.84	2.10	2.931 (3)	171

Symmetry code: (i) $x, y + 1, z$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2008); cell refinement: *CrysAlis CCD*; data reduction: *CrysAlis CCD*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL-Plus* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5238).

References

- Abraham, L., Czerwonka, R. & Hiersemann, M. (2001). *Angew. Chem. Int. Ed.* **40**, 4700–4703.
- Abraham, L., Körner, M., Schwab, P. & Hiersemann, M. (2004). *Adv. Synth. Catal.* **346**, 1281–1294.
- Brown, C. A. (1973). *J. Am. Chem. Soc.* **95**, 4100–4102.
- Evans, D. A., Bartroli, J. & Shih, T. L. (1981). *J. Am. Chem. Soc.* **103**, 2127–2129.
- Evans, D. A., Miller, S. J., Lectka, T. & Matt, V. P. (1999). *J. Am. Chem. Soc.* **121**, 7559–7573.
- Gille, A., Schürmann, M., Preut, H. & Hiersemann, M. (2008). *Acta Cryst. E64*, o1835.
- Otera, J., Niibo, Y. & Nozaki, H. (1992). *Tetrahedron Lett.* **33**, 3655–3658.
- Oxford Diffraction (2008). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supporting information

Acta Cryst. (2009). E65, o3274 [doi:10.1107/S1600536809050399]

(3*R*,4*R*,5*S*)-4-Hydroxy-3-methyl-5-[(2*S*,3*R*)-3-methylpent-4-en-2-yl]-4,5-dihydrofuran-2(3*H*)-one

Annika Gille, Markus Schürmann, Hans Preut and Martin Hiersemann

S1. Comment

The title compound, (I), was synthesized using a catalytic asymmetric Claisen rearrangement (Abraham *et al.*, 2001; Abraham *et al.*, 2004), a diastereoselective reduction with K-Selectride (Brown, 1973) and an Evans aldol addition (Evans *et al.*, 1981). In order to verify the relative configuration of the obtained diastereomeric aldol adducts, 4-(*tert*-butyldimethylsilyloxy)-3-hydroxy-2,5,6-trimethyloct-7-enoyl-4-isopropylloxazolidin-2-one, (III), the γ -lactones (II) and (I) were prepared by removal of the silyl protecting group (Otera *et al.*, 1992) and subsequent *in situ* lactonization. The diastereomeric mixture of the γ -lactones could be separated by column chromatography. An X-ray crystal structure analysis of the major diastereomer (II) has already been published (Gille *et al.*, 2008). Fig. 1 depicts the structure of the isolated minor diastereomer (I). The configuration of the chiral C atoms in (I) can be attributed to the stereochemical course of the Evans aldol addition (C3 *R* and C4 *R*), the diastereoselective reduction with K-Selectride (C5 *S*) and the catalytic asymmetric Claisen rearrangement (C[2] *S* and C[3] *R*) using the chiral Lewis acid [Cu{(*S,S*)-*tert*-butyl-box}] (H_2O)₂(SbF₆)₂ (Evans *et al.*, 1999).

S2. Experimental

The title compound, (I), was synthesized from the corresponding aldol adduct, (III), using tetrabutylammonium fluoride (TBAF) in the presence of acetic acid (Otera *et al.*, 1992) to remove the silyl protecting group. The subsequent lactonization proceeded *in situ*.

To an ice-cooled solution of crude (III) (dr = 49/51, 0.04 g, 0.10 mmol, 1 eq) in THF (1 ml, 11 ml/mmol III) was added a solution of AcOH (0.5 μ l, 0.010 mmol, 0.1 eq) in THF (0.1 ml, 1.1 ml/mmol III) and TBAF (1 M in THF, 0.11 ml, 0.11 mmol, 1.1 eq). After 15 min at 273 K, the reaction mixture was diluted by the addition of saturated aqueous NH₄Cl solution. The aqueous layer was extracted with CH₂Cl₂ (4x) and the combined organic phases were dried (MgSO₄) and concentrated under reduced pressure. Purification by flash chromatography (crude product charged on silica gel, cyclohexane/ethyl acetate 10/1 to 5/1) afforded lactone (I) (0.006 g, 0.03 mmol, 30%) as a single diastereomer and additionally a mixture of (I) and the diastereomer (II) (0.013 g, 0.07 mmol, 69%, dr = 70/30) as colourless crystals. Subsequent recrystallization of (I) by vapor diffusion technique from isohexane and ethyl acetate provided a colourless plate of (I) single-crystal suitable for an X-ray crystal structure analysis. R_f 0.35 (cyclohexane/ethyl acetate 2/1); mp 378 K; ¹H NMR (CDCl₃, 400 MHz, δ): 0.92 (d, ³J = 7.1 Hz, 3H), 1.00 (d, ³J = 7.0 Hz, 3H), 1.32 (d, ³J = 7.2 Hz, 3H), 1.85 (dq, ³J = 8.2, 7.1, 4.9 Hz, 1H), 2.1 (br. s, 1H), 2.51–2.63 (m, 1H), 2.63 (dq, ³J = 8.4, 7.2 Hz, 1H), 3.92 (dd, ³J = 7.4, 8.2 Hz, 1H), 4.07 (dd, ³J = 8.4, 7.4 Hz, 1H), 5.04 (dd, ³J(E) = 17.8 Hz, ²J = 1.3 Hz, 1H), 5.05 (dd, ³J(Z) = 10.0 Hz, ²J = 1.3 Hz, 1H), 5.81 (ddd, ³J(E) = 17.8 Hz, ³J(Z) = 10.0 Hz, ³J = 6.8 Hz, 1H); ¹³C NMR (CDCl₃, 101 MHz, δ): 10.1 (CH₃), 12.9 (CH₃), 13.7 (CH₃), 37.8 (CH), 41.4 (CH), 44.4 (CH), 78.0 (CH), 84.7 (CH), 114.5 (CH₂), 142.6 (CH), 176.4 (C); IR (cm⁻¹): 3400(br,s) (ν O—H, OH in H-bridges), 3085(w) (ν C—H, olefin), 2970(s) 2925(s) 2890(m) 2855(m) (ν C—H, CH, CH₃), 1735(s) (ν

C=O, lactone), 1640(w) (ν C=C), 1455(m) (δ_{as} C—H, CH₂), 1380(m) (δ_s C—H, CH₃), 1095(s) (ν C—O, alcohol), 1010(s) 910(s) (δ C—H, olefin); Anal. Calcd. for C₁₁H₁₈O₃: C, 66.6; H, 9.2; Found: C, 66.6; H, 9.4; [α]_D²⁰ +16.0 (c 0.6, CHCl₃); C₁₁H₁₈O₃, $M = 198.26$ g/mol.




Figure 1

The molecular structure of (I) with displacement ellipsoids shown at the 30% probability level.

(3*R*,4*R*,5*S*)-4-Hydroxy-3-methyl-5-[(2*S*,3*R*)-3-methylpent-4-en-2-yl]-4,5-dihydrofuran-2(3*H*)-one

Crystal data

C₁₁H₁₈O₃
 $M_r = 198.25$
Monoclinic, P2₁
Hall symbol: P 2yb
 $a = 7.7265$ (10) Å
 $b = 6.4798$ (8) Å
 $c = 11.0598$ (16) Å
 $\beta = 92.563$ (14)°
 $V = 553.17$ (13) Å³
 $Z = 2$

$F(000) = 216$
 $D_x = 1.190 \text{ Mg m}^{-3}$
Mo K α radiation, $\lambda = 0.71073$ Å
Cell parameters from 1159 reflections
 $\theta = 2.6\text{--}29.1^\circ$
 $\mu = 0.09 \text{ mm}^{-1}$
 $T = 173 \text{ K}$
Plate, colourless
0.50 × 0.18 × 0.04 mm

Data collection

Oxford Diffraction Xcalibur S CCD
diffractometer
Radiation source: Enhance (Mo) X-ray Source
Graphite monochromator
Detector resolution: 16.0560 pixels mm⁻¹
 ω scans
3255 measured reflections

1129 independent reflections
737 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.040$
 $\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 2.6^\circ$
 $h = -10 \rightarrow 10$
 $k = -7 \rightarrow 8$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.048$
 $S = 0.99$
1129 reflections
131 parameters
1 restraint

Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.011P)^2]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.14 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.15 \text{ e \AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.2173 (2)	0.1438 (2)	0.07628 (15)	0.0251 (5)
O2	0.2731 (2)	0.0175 (3)	-0.10413 (17)	0.0328 (6)
O3	0.1599 (2)	0.6838 (3)	0.05289 (17)	0.0370 (6)
H3	0.2032	0.7730	0.0082	0.056*
C1	0.2453 (3)	0.1658 (5)	-0.0428 (2)	0.0267 (8)
C2	0.2329 (4)	0.3900 (4)	-0.0771 (2)	0.0251 (8)
H2	0.1128	0.4168	-0.1109	0.030*
C3	0.2533 (3)	0.4969 (4)	0.0446 (2)	0.0275 (8)
H3A	0.3790	0.5230	0.0645	0.033*
C4	0.1847 (4)	0.3431 (4)	0.1331 (3)	0.0251 (7)
H4	0.0569	0.3630	0.1382	0.030*
C5	0.2674 (4)	0.3375 (4)	0.2598 (2)	0.0293 (8)
H5	0.3954	0.3280	0.2512	0.035*
C6	0.2135 (4)	0.1458 (4)	0.3322 (2)	0.0314 (8)
H6	0.2292	0.0238	0.2784	0.038*
C7	0.3345 (4)	0.1158 (5)	0.4405 (3)	0.0409 (9)
H7	0.4546	0.1147	0.4255	0.049*
C8	0.2950 (4)	0.0912 (5)	0.5523 (2)	0.0487 (10)
H8A	0.1770	0.0910	0.5730	0.058*
H8B	0.3841	0.0735	0.6134	0.058*
C9	0.3607 (3)	0.4531 (4)	-0.1730 (2)	0.0350 (9)
H9A	0.4789	0.4182	-0.1443	0.053*
H9B	0.3523	0.6022	-0.1872	0.053*
H9C	0.3323	0.3793	-0.2487	0.053*
C10	0.2329 (4)	0.5418 (4)	0.3243 (2)	0.0401 (9)
H10A	0.2922	0.6541	0.2836	0.060*
H10B	0.2765	0.5329	0.4087	0.060*
H10C	0.1080	0.5689	0.3218	0.060*
C11	0.0231 (3)	0.1493 (5)	0.3622 (2)	0.0434 (9)
H11A	-0.0082	0.0167	0.3979	0.065*
H11B	-0.0484	0.1728	0.2881	0.065*
H11C	0.0033	0.2605	0.4201	0.065*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0324 (13)	0.0134 (13)	0.0295 (11)	-0.0005 (11)	0.0026 (10)	-0.0004 (11)
O2	0.0385 (14)	0.0195 (12)	0.0403 (14)	-0.0003 (11)	0.0013 (11)	-0.0058 (12)
O3	0.0427 (14)	0.0116 (11)	0.0573 (15)	0.0053 (11)	0.0087 (11)	0.0052 (12)
C1	0.0145 (16)	0.0214 (18)	0.044 (2)	-0.0032 (17)	-0.0048 (15)	0.003 (2)
C2	0.0219 (17)	0.0155 (18)	0.038 (2)	-0.0004 (14)	0.0015 (15)	0.0037 (15)
C3	0.0222 (18)	0.0125 (16)	0.048 (2)	0.0004 (16)	0.0007 (16)	0.0048 (16)
C4	0.0229 (17)	0.0143 (16)	0.0380 (19)	0.0038 (15)	0.0020 (15)	-0.0005 (16)
C5	0.0261 (17)	0.0224 (19)	0.039 (2)	0.0040 (16)	-0.0001 (16)	-0.0068 (17)
C6	0.048 (2)	0.0133 (19)	0.0330 (18)	0.0029 (17)	0.0011 (16)	-0.0060 (17)
C7	0.049 (2)	0.038 (2)	0.0359 (17)	0.0078 (19)	0.0001 (17)	0.0032 (18)
C8	0.054 (3)	0.052 (2)	0.039 (2)	-0.003 (2)	-0.005 (2)	0.0019 (19)
C9	0.0337 (19)	0.0271 (19)	0.045 (2)	-0.0014 (17)	0.0090 (16)	0.0070 (17)
C10	0.046 (2)	0.025 (2)	0.049 (2)	-0.0045 (17)	0.0060 (18)	-0.0113 (17)
C11	0.049 (2)	0.037 (2)	0.0442 (19)	-0.010 (2)	0.0005 (17)	0.007 (2)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.352 (3)	C6—C7	1.499 (3)
O1—C4	1.463 (3)	C6—C11	1.523 (3)
O2—C1	1.201 (3)	C6—H6	1.0000
O3—C3	1.415 (3)	C7—C8	1.296 (3)
O3—H3	0.8400	C7—H7	0.9500
C1—C2	1.504 (4)	C8—H8A	0.9500
C2—C3	1.516 (3)	C8—H8B	0.9500
C2—C9	1.536 (3)	C9—H9A	0.9800
C2—H2	1.0000	C9—H9B	0.9800
C3—C4	1.510 (3)	C9—H9C	0.9800
C3—H3A	1.0000	C10—H10A	0.9800
C4—C5	1.515 (3)	C10—H10B	0.9800
C4—H4	1.0000	C10—H10C	0.9800
C5—C10	1.533 (3)	C11—H11A	0.9800
C5—C6	1.544 (4)	C11—H11B	0.9800
C5—H5	1.0000	C11—H11C	0.9800
C1—O1—C4	111.3 (2)	C7—C6—C5	110.2 (2)
C3—O3—H3	109.5	C11—C6—C5	112.7 (2)
O2—C1—O1	120.4 (3)	C7—C6—H6	106.6
O2—C1—C2	129.9 (2)	C11—C6—H6	106.6
O1—C1—C2	109.7 (3)	C5—C6—H6	106.6
C1—C2—C3	102.4 (2)	C8—C7—C6	127.8 (3)
C1—C2—C9	113.4 (2)	C8—C7—H7	116.1
C3—C2—C9	116.5 (2)	C6—C7—H7	116.1
C1—C2—H2	108.1	C7—C8—H8A	120.0
C3—C2—H2	108.1	C7—C8—H8B	120.0
C9—C2—H2	108.1	H8A—C8—H8B	120.0

O3—C3—C4	109.1 (2)	C2—C9—H9A	109.5
O3—C3—C2	114.6 (2)	C2—C9—H9B	109.5
C4—C3—C2	104.4 (2)	H9A—C9—H9B	109.5
O3—C3—H3A	109.5	C2—C9—H9C	109.5
C4—C3—H3A	109.5	H9A—C9—H9C	109.5
C2—C3—H3A	109.5	H9B—C9—H9C	109.5
O1—C4—C3	103.4 (2)	C5—C10—H10A	109.5
O1—C4—C5	107.6 (2)	C5—C10—H10B	109.5
C3—C4—C5	118.0 (2)	H10A—C10—H10B	109.5
O1—C4—H4	109.1	C5—C10—H10C	109.5
C3—C4—H4	109.1	H10A—C10—H10C	109.5
C5—C4—H4	109.1	H10B—C10—H10C	109.5
C4—C5—C10	109.6 (2)	C6—C11—H11A	109.5
C4—C5—C6	112.7 (2)	C6—C11—H11B	109.5
C10—C5—C6	113.4 (2)	H11A—C11—H11B	109.5
C4—C5—H5	106.9	C6—C11—H11C	109.5
C10—C5—H5	106.9	H11A—C11—H11C	109.5
C6—C5—H5	106.9	H11B—C11—H11C	109.5
C7—C6—C11	113.6 (2)		
C4—O1—C1—O2	-179.5 (2)	C2—C3—C4—O1	28.3 (3)
C4—O1—C1—C2	-0.1 (3)	O3—C3—C4—C5	-90.1 (3)
O2—C1—C2—C3	-162.7 (3)	C2—C3—C4—C5	147.0 (2)
O1—C1—C2—C3	18.0 (3)	O1—C4—C5—C10	-178.7 (2)
O2—C1—C2—C9	-36.3 (4)	C3—C4—C5—C10	64.9 (3)
O1—C1—C2—C9	144.4 (2)	O1—C4—C5—C6	-51.4 (3)
C1—C2—C3—O3	-147.4 (2)	C3—C4—C5—C6	-167.8 (2)
C9—C2—C3—O3	88.3 (3)	C4—C5—C6—C7	163.9 (2)
C1—C2—C3—C4	-28.1 (3)	C10—C5—C6—C7	-70.9 (3)
C9—C2—C3—C4	-152.4 (2)	C4—C5—C6—C11	-68.0 (3)
C1—O1—C4—C3	-18.0 (3)	C10—C5—C6—C11	57.3 (3)
C1—O1—C4—C5	-143.6 (2)	C11—C6—C7—C8	0.8 (5)
O3—C3—C4—O1	151.2 (2)	C5—C6—C7—C8	128.5 (4)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O3—H3···O1 ⁱ	0.84	2.52	3.023 (2)	120
O3—H3···O2 ⁱ	0.84	2.10	2.931 (3)	171

Symmetry code: (i) $x, y+1, z$.