

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

**Poly[diacquabis( $\mu_3$ -1*H*-benzimidazole-5,6-dicarboxylato- $\kappa^4$ N<sup>3</sup>:O<sup>5</sup>,O<sup>5'</sup>:O<sup>6</sup>)bis-( $\mu_2$ -1*H*,3*H*-benzimidazolium-5,6-dicarboxylato- $\kappa^3$ O<sup>5</sup>,O<sup>5'</sup>:O<sup>6</sup>)-digadolinium(III)]**

Jie-Xuan Huang,<sup>a</sup> Yi-Yi Wu,<sup>a</sup> Chun-De Huang,<sup>a</sup> Qing-Yang Lian<sup>a</sup> and Rong-Hua Zeng<sup>a,b\*</sup>

<sup>a</sup>School of Chemistry and Environment, South China Normal University, Guangzhou 510006, People's Republic of China, and <sup>b</sup>Key Laboratory of Technology of Electrochemical Energy Storage and Power Generation in Guangdong Universities, South China Normal University, Guangzhou 510006, People's Republic of China  
Correspondence e-mail: zrh321@yahoo.com.cn


Received 19 October 2009; accepted 5 November 2009

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.020;  $wR$  factor = 0.049; data-to-parameter ratio = 10.7.

In the title complex,  $[\text{Gd}_2(\text{C}_9\text{H}_4\text{N}_2\text{O}_4)_2(\text{C}_9\text{H}_5\text{N}_2\text{O}_4)_2(\text{H}_2\text{O})_2]_n$ , two of the benzimidazole-5,6-dicarboxylate ligands are protonated at the imidazole groups. Each  $\text{Gd}^{\text{III}}$  ion is coordinated by six O atoms and one N atom from five ligands and one water molecule, displaying a distorted bicapped trigonal-prismatic geometry. The  $\text{Gd}^{\text{III}}$  ions are linked by the carboxylate groups and imidazole N atoms, forming a layer parallel to (001). These layers are further connected by O—H...O and N—H...O hydrogen bonds into a three-dimensional supramolecular network.

## Related literature

For related structures, see: Gao *et al.* (2008); Lo *et al.* (2007); Wei *et al.* (2008); Yao *et al.* (2008).



## Experimental

## Crystal data

$[\text{Gd}_2(\text{C}_9\text{H}_4\text{N}_2\text{O}_4)_2(\text{C}_9\text{H}_5\text{N}_2\text{O}_4)_2(\text{H}_2\text{O})_2]$   
 $M_r = 1169.12$   
Monoclinic,  $P2_1/c$   
 $a = 18.7856$  (11) Å  
 $b = 12.7745$  (7) Å  
 $c = 15.4776$  (9) Å

$\beta = 108.010$  (1)°  
 $V = 3532.3$  (3) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 3.82$  mm<sup>-1</sup>  
 $T = 296$  K  
 $0.25 \times 0.24 \times 0.21$  mm

## Data collection

Bruker APEXII CCD diffractometer  
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\text{min}} = 0.448$ ,  $T_{\text{max}} = 0.501$

24827 measured reflections  
6338 independent reflections  
5929 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.027$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.020$   
 $wR(F^2) = 0.049$   
 $S = 1.08$   
6338 reflections  
595 parameters  
13 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.63$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.55$  e Å<sup>-3</sup>

Table 1

Selected bond lengths (Å).

Gd1—O1	2.338 (2)	Gd2—O9	2.312 (2)
Gd1—O2	2.526 (2)	Gd2—O11 <sup>iii</sup>	2.539 (2)
Gd1—O3 <sup>i</sup>	2.350 (2)	Gd2—O12 <sup>iii</sup>	2.342 (2)
Gd1—O5 <sup>ii</sup>	2.461 (2)	Gd2—O13 <sup>iv</sup>	2.267 (2)
Gd1—O6 <sup>ii</sup>	2.499 (2)	Gd2—O15	2.449 (2)
Gd1—O8	2.314 (2)	Gd2—O16	2.520 (2)
Gd1—N6	2.617 (3)	Gd2—N1 <sup>iii</sup>	2.612 (3)
Gd1—O1W	2.374 (2)	Gd2—O2W	2.384 (2)

Symmetry codes: (i)  $-x, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (iii)  $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (iv)  $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$ .

**Table 2**  
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1W—H1W $\cdots$ O7 <sup>v</sup>	0.84	1.80	2.629 (3)	167
O1W—H2W $\cdots$ O4 <sup>i</sup>	0.82	1.92	2.670 (3)	153
O2W—H3W $\cdots$ O10 <sup>vi</sup>	0.86	1.74	2.590 (3)	172
O2W—H4WA $\cdots$ O15	0.82	2.14	2.716 (5)	127
O2W—H4WB $\cdots$ O2W <sup>vi</sup>	0.82	1.91	2.723 (4)	170
N2—H2A $\cdots$ O14 <sup>viii</sup>	0.82 (4)	1.94 (4)	2.742 (3)	169 (3)
N3—H3A $\cdots$ O10 <sup>vii</sup>	0.81 (4)	1.94 (4)	2.731 (4)	166 (4)
N4—H4A $\cdots$ O16	0.84 (4)	1.97 (4)	2.801 (4)	170 (4)
N5—H5A $\cdots$ O4 <sup>v</sup>	0.83 (4)	1.96 (4)	2.772 (3)	165 (4)
N7—H7A $\cdots$ O5	0.77 (4)	2.08 (4)	2.845 (3)	171 (4)
N8—H8A $\cdots$ O7 <sup>viii</sup>	0.80 (4)	1.95 (4)	2.747 (3)	172 (4)

Symmetry codes: (i)  $-x, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (v)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (vi)  $-x + 1, -y, -z$ ; (vii)  $x, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (viii)  $x, -y - \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINTE* (Bruker, 2007); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors acknowledge the Students' Extracurricular Scientific Research Projects of South China Normal University and the Key Laboratory of Electrochemical Technology of Energy Storage and Power Generation in Guangdong Universities for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2247).

## References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.  
 Bruker (2007). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Gao, Q., Gao, W.-H., Zhang, C.-Y. & Xie, Y.-B. (2008). *Acta Cryst.* **E64**, m928.  
 Lo, Y.-L., Wang, W.-C., Lee, G.-A. & Liu, Y.-H. (2007). *Acta Cryst.* **E63**, m2657–m2658.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Wei, Y.-Q., Yu, Y.-F. & Wu, K.-C. (2008). *Cryst. Growth Des.* **8**, 2087–2089.  
 Yao, Y.-L., Che, Y.-X. & Zheng, J.-M. (2008). *Cryst. Growth Des.* **8**, 2299–2306.

**supplementary materials**

*Acta Cryst.* (2009). E65, m1566-m1567 [ doi:10.1107/S1600536809046819 ]

**Poly[*diaquabis*( $\mu_3$ -1*H*-benzimidazole-5,6-dicarboxylato- $\kappa^4$ N<sup>3</sup>:O<sup>5</sup>,O<sup>5'</sup>:O<sup>6</sup>)bis( $\mu_2$ -1*H*,3*H*-benzimidazolium-5,6-dicarboxylato- $\kappa^3$ O<sup>5</sup>,O<sup>5'</sup>:O<sup>6</sup>)digadolinium(III)]**

**J.-X. Huang, Y.-Y. Wu, C.-D. Huang, Q.-Y. Lian and R.-H. Zeng**

### Comment

In recent years, research on coordination polymers has made considerable progress in the fields of supramolecular chemistry and crystal engineering, because of their intriguing structural motifs and functional properties, such as molecular adsorption, magnetism and luminescence. In general, the structural motifs of these hybrid compounds are closely related to the geometries of metal centers and the number of coordination sites provided by multidentate ligands. On the other hand, the supramolecular interactions such as hydrogen-bonding,  $\pi$ - $\pi$  stacking and metallophilic interactions also play the key roles in the recognition process forming final three-dimensional architectures. As a building block, benzimidazole-5,6-dicarboxylic acid is a good ligand with multifunctional coordination sites providing intriguing architectures and topologies (Gao *et al.*, 2008; Lo *et al.*, 2007; Wei *et al.*, 2008; Yao *et al.*, 2008). Recently, we obtained the title coordination polymer, which was synthesized under hydrothermal conditions.

In the title compound (Fig. 1), two of the benzimidazole-5,6-dicarboxylate ligands are protonated at the imidazole groups. Each Gd<sup>III</sup> ion is eight-coordinated by six O atoms one N atom from five ligands and one water molecule. The coordination geometry can be described as distorted bicapped trigonal prismatic, with Gd—O distances and O—Gd—O angles ranging from 2.267 (2) to 2.539 (2) Å (Table 1) and 52.14 (7) to 156.82 (8)°, respectively. The benzimidazole-5,6-dicarboxylate ligands acting as bridging ligands link the Gd<sup>III</sup> centers into a layer parallel to the (0 0 1) plane. O—H...O and N—H...O hydrogen bonds (Table 2) connect the layers into a three-dimensional supramolecular motif (Fig. 2). Within the layer, the  $\pi$ - $\pi$  stacking interactions between neighboring imidazole and benzene rings [centroid-centroid distances = 3.629 (3), 3.755 (4), 3.656 (3) and 3.606 (3) Å] enhance the stability of the crystal structure.

### Experimental

A mixture of Gd<sub>2</sub>O<sub>3</sub> (0.363 g, 1 mmol), benzimidazole-5,6-dicarboxylic acid (0.206 g, 1 mmol), water (10 ml) in the presence of HClO<sub>4</sub> (0.039 g, 0.385 mmol) was stirred vigorously for 30 min and then sealed in a Teflon-lined stainless-steel autoclave (20 ml capacity). The autoclave was heated and maintained at 433 K for 3 d, and then cooled to room temperature at 5 K h<sup>-1</sup>. The colorless block crystals were obtained.

### Refinement

Water H atoms were tentatively located in difference Fourier maps and fixed in refinements, with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ . One of the H atoms of O2W is disordered over two sites (O4WA and O4WB), each with an occupancy factor of 0.5. H atoms on N atoms were tentatively located in difference Fourier maps and refined with distance restraints of N—H = 0.82 (1) Å and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ . H atoms attached to C atoms were positioned geometrically and refined as riding, with C—H = 0.93 Å and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

## Figures




Fig. 1. The asymmetric unit of the title compound. Displacement ellipsoids are shown at the 50% probability level. H atoms have been omitted for clarity. [Symmetry codes: (i)  $-x, -1/2+y, 1/2-z$ ; (ii)  $-x, 1/2+y, 1/2-z$ ; (iii)  $1-x, -1/2+y, 1/2-z$ ; (iv)  $1-x, 1/2+y, 1/2-z$ .]




Fig. 2. A view of the three-dimensional supramolecular network. Hydrogen bonds are shown as dashed lines.

## Poly[diabiquis( $\mu_3$ -1*H*-benzimidazole-5,6-dicarboxylato- $\kappa^4 N^3:O^5,O^{5'}:O^6$ )bis( $\mu_2$ - 1*H*,3*H*-benzimidazolium-5,6-dicarboxylato- $\kappa^3 O^5,O^{5'}:O^6$ )digadolinium(III)]

### Crystal data

$[\text{Gd}_2(\text{C}_9\text{H}_4\text{N}_2\text{O}_4)_2(\text{C}_9\text{H}_5\text{N}_2\text{O}_4)_2(\text{H}_2\text{O})_2]$	$F_{000} = 2264$
$M_r = 1169.12$	$D_x = 2.198 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: $-P\ 2ybc$	Cell parameters from 7234 reflections
$a = 18.7856 (11) \text{ \AA}$	$\theta = 2.6\text{--}28.2^\circ$
$b = 12.7745 (7) \text{ \AA}$	$\mu = 3.82 \text{ mm}^{-1}$
$c = 15.4776 (9) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 108.010 (1)^\circ$	Block, colorless
$V = 3532.3 (3) \text{ \AA}^3$	$0.25 \times 0.24 \times 0.21 \text{ mm}$
$Z = 4$	

### Data collection

Bruker APEXII CCD diffractometer	6338 independent reflections
Radiation source: fine-focus sealed tube	5929 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
$T = 296 \text{ K}$	$\theta_{\text{max}} = 25.2^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -21 \rightarrow 22$
$T_{\text{min}} = 0.448, T_{\text{max}} = 0.501$	$k = -15 \rightarrow 15$
24827 measured reflections	$l = -18 \rightarrow 18$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.020$	H atoms treated by a mixture of

independent and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.021P)^2 + 3.3986P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.002$   
 $\Delta\rho_{\max} = 0.63 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.55 \text{ e } \text{\AA}^{-3}$   
 Extinction correction: none

$wR(F^2) = 0.049$   
 $S = 1.08$   
 6338 reflections  
 595 parameters  
 13 restraints  
 Primary atom site location: structure-invariant direct methods

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Gd1	0.028568 (8)	0.296570 (11)	0.225244 (9)	0.01343 (5)	
Gd2	0.524307 (8)	-0.054993 (10)	0.209686 (9)	0.01317 (5)	
N6	0.10953 (14)	0.1586 (2)	0.17264 (18)	0.0208 (6)	
N5	0.12241 (15)	0.0250 (2)	0.08627 (19)	0.0251 (6)	
H5A	0.111 (2)	-0.029 (3)	0.056 (2)	0.030*	
C23	0.07832 (18)	0.0751 (2)	0.1265 (2)	0.0218 (7)	
H23	0.0302	0.0529	0.1222	0.026*	
C25	0.24026 (17)	0.2316 (2)	0.1949 (2)	0.0220 (7)	
H25	0.2367	0.2863	0.2331	0.026*	
C21	0.25303 (17)	0.0637 (2)	0.0817 (2)	0.0215 (7)	
H21	0.2569	0.0077	0.0449	0.026*	
C24	0.18074 (16)	0.1631 (2)	0.1607 (2)	0.0181 (6)	
C22	0.18850 (17)	0.0794 (2)	0.1059 (2)	0.0197 (6)	
C20	0.31133 (16)	0.1340 (2)	0.11398 (19)	0.0166 (6)	
C26	0.30518 (17)	0.2172 (2)	0.1711 (2)	0.0178 (6)	
N3	0.36219 (16)	0.1575 (2)	0.4264 (2)	0.0312 (7)	
H3A	0.375 (2)	0.212 (3)	0.453 (3)	0.037*	
C13	0.28840 (17)	0.0294 (2)	0.3540 (2)	0.0199 (7)	
N4	0.35934 (15)	0.0200 (2)	0.3445 (2)	0.0256 (6)	
H4A	0.372 (2)	-0.029 (3)	0.316 (2)	0.031*	
C15	0.29033 (17)	0.1190 (2)	0.4058 (2)	0.0195 (6)	
C14	0.40105 (19)	0.0970 (3)	0.3890 (2)	0.0334 (8)	
H14	0.4511	0.1073	0.3934	0.040*	
C27	0.36712 (17)	0.2947 (2)	0.2069 (2)	0.0207 (7)	
C19	0.38033 (17)	0.1168 (2)	0.0861 (2)	0.0181 (6)	
O11	0.41746 (14)	0.30726 (19)	0.17120 (17)	0.0342 (6)	
O10	0.37957 (14)	0.15189 (19)	0.01034 (16)	0.0338 (6)	
C17	0.16278 (17)	0.0978 (2)	0.39339 (19)	0.0163 (6)	
C12	0.22389 (17)	-0.0299 (2)	0.3229 (2)	0.0206 (7)	
H12	0.2230	-0.0912	0.2902	0.025*	
C16	0.22774 (17)	0.1551 (2)	0.4263 (2)	0.0199 (7)	
H16	0.2295	0.2153	0.4607	0.024*	
C11	0.16054 (17)	0.0050 (2)	0.3421 (2)	0.0177 (6)	
C10	0.08996 (17)	-0.0570 (2)	0.3073 (2)	0.0185 (7)	
C18	0.09517 (16)	0.1407 (2)	0.41501 (19)	0.0161 (6)	

## supplementary materials

---

O8	0.05302 (12)	0.20217 (15)	0.35883 (14)	0.0207 (5)
O7	0.08767 (13)	0.11675 (18)	0.48980 (14)	0.0269 (5)
O5	0.08938 (12)	-0.13723 (16)	0.25772 (16)	0.0266 (5)
O6	0.03295 (12)	-0.03134 (16)	0.32721 (15)	0.0246 (5)
O12	0.36722 (13)	0.3474 (2)	0.27665 (18)	0.0391 (7)
C36	0.40339 (17)	-0.1991 (2)	0.1826 (2)	0.0185 (6)
C33	0.20968 (17)	-0.2999 (2)	0.1476 (2)	0.0192 (6)
C35	0.33593 (17)	-0.2677 (2)	0.1548 (2)	0.0175 (6)
C34	0.27150 (17)	-0.2345 (2)	0.1727 (2)	0.0200 (7)
H34	0.2699	-0.1705	0.2006	0.024*
O16	0.40712 (12)	-0.12307 (16)	0.23682 (15)	0.0246 (5)
O15	0.45572 (13)	-0.21597 (18)	0.15068 (17)	0.0320 (6)
O9	0.43302 (13)	0.06471 (18)	0.13734 (16)	0.0294 (5)
N7	0.14022 (15)	-0.2953 (2)	0.16138 (19)	0.0242 (6)
H7A	0.127 (2)	-0.248 (3)	0.184 (2)	0.029*
C28	0.40817 (17)	-0.4085 (2)	0.09601 (19)	0.0163 (6)
C29	0.33774 (16)	-0.3652 (2)	0.11096 (19)	0.0159 (6)
C32	0.10256 (17)	-0.3811 (3)	0.1275 (2)	0.0233 (7)
H32	0.0547	-0.3968	0.1292	0.028*
O14	0.41716 (12)	-0.40122 (17)	0.02045 (14)	0.0237 (5)
N8	0.14248 (15)	-0.4412 (2)	0.09098 (19)	0.0219 (6)
H8A	0.126 (2)	-0.495 (3)	0.066 (2)	0.026*
C31	0.21123 (16)	-0.3943 (2)	0.1031 (2)	0.0168 (6)
C30	0.27505 (17)	-0.4281 (2)	0.0840 (2)	0.0192 (6)
H30	0.2755	-0.4911	0.0539	0.023*
C1	0.13784 (17)	0.4457 (2)	0.3037 (2)	0.0188 (7)
C2	0.19826 (17)	0.5268 (2)	0.3348 (2)	0.0176 (6)
O2	0.09162 (12)	0.42965 (17)	0.34523 (15)	0.0249 (5)
O1	0.13579 (12)	0.39355 (17)	0.23230 (16)	0.0287 (5)
C8	0.18820 (16)	0.6163 (2)	0.38430 (19)	0.0158 (6)
C7	0.24484 (17)	0.6887 (2)	0.4144 (2)	0.0185 (6)
H7	0.2389	0.7474	0.4471	0.022*
C4	0.32222 (16)	0.5828 (2)	0.3467 (2)	0.0175 (6)
C6	0.31138 (16)	0.6714 (2)	0.39437 (19)	0.0172 (6)
C3	0.26477 (16)	0.5098 (2)	0.3167 (2)	0.0193 (7)
H3	0.2711	0.4506	0.2849	0.023*
N1	0.39340 (14)	0.5868 (2)	0.33531 (17)	0.0190 (5)
N2	0.37599 (15)	0.7294 (2)	0.41004 (18)	0.0217 (6)
H2A	0.385 (2)	0.785 (3)	0.437 (2)	0.026*
C5	0.42138 (17)	0.6759 (2)	0.3735 (2)	0.0208 (7)
H5	0.4685	0.6999	0.3752	0.025*
C9	0.11320 (17)	0.6424 (2)	0.3956 (2)	0.0165 (6)
O4	0.10804 (12)	0.64817 (17)	0.47426 (14)	0.0242 (5)
O13	0.45307 (12)	-0.45613 (17)	0.16304 (14)	0.0247 (5)
O3	0.06077 (11)	0.66278 (16)	0.32358 (14)	0.0203 (5)
O1W	-0.00880 (13)	0.30447 (17)	0.06416 (14)	0.0264 (5)
H1W	0.0184	0.3242	0.0331	0.040*
H2W	-0.0314	0.2519	0.0407	0.040*
O2W	0.51762 (14)	-0.0852 (2)	0.05540 (15)	0.0346 (6)

H3W	0.5544	-0.1080	0.0390	0.052*	
H4WA	0.4879	-0.1341	0.0480	0.052*	0.50
H4WB	0.5025	-0.0331	0.0237	0.052*	0.50

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Gd1	0.01015 (8)	0.01374 (8)	0.01696 (8)	-0.00059 (5)	0.00501 (6)	0.00056 (5)
Gd2	0.01062 (8)	0.01412 (8)	0.01541 (8)	-0.00007 (5)	0.00497 (6)	-0.00022 (5)
N6	0.0124 (13)	0.0236 (14)	0.0276 (14)	-0.0025 (11)	0.0079 (11)	-0.0044 (11)
N5	0.0183 (15)	0.0242 (14)	0.0323 (16)	-0.0062 (12)	0.0071 (12)	-0.0130 (12)
C23	0.0146 (16)	0.0244 (16)	0.0278 (17)	-0.0023 (13)	0.0084 (14)	-0.0022 (13)
C25	0.0183 (17)	0.0186 (15)	0.0294 (17)	-0.0016 (13)	0.0077 (14)	-0.0101 (13)
C21	0.0188 (17)	0.0214 (16)	0.0252 (17)	-0.0004 (13)	0.0080 (14)	-0.0090 (13)
C24	0.0115 (15)	0.0204 (15)	0.0233 (16)	0.0007 (12)	0.0067 (12)	-0.0028 (12)
C22	0.0152 (16)	0.0202 (15)	0.0232 (16)	-0.0026 (12)	0.0052 (13)	-0.0044 (13)
C20	0.0123 (15)	0.0210 (15)	0.0155 (14)	0.0035 (12)	0.0029 (12)	0.0036 (12)
C26	0.0141 (16)	0.0166 (15)	0.0235 (16)	-0.0007 (12)	0.0067 (13)	-0.0023 (12)
N3	0.0182 (16)	0.0345 (17)	0.0425 (18)	-0.0092 (13)	0.0116 (13)	-0.0153 (14)
C13	0.0139 (16)	0.0242 (16)	0.0230 (16)	0.0032 (13)	0.0077 (13)	-0.0003 (13)
N4	0.0163 (15)	0.0311 (16)	0.0328 (16)	0.0008 (12)	0.0126 (12)	-0.0051 (13)
C15	0.0126 (16)	0.0225 (16)	0.0221 (16)	-0.0037 (12)	0.0036 (13)	-0.0016 (13)
C14	0.0156 (18)	0.043 (2)	0.044 (2)	-0.0055 (15)	0.0122 (16)	-0.0085 (17)
C27	0.0136 (16)	0.0185 (15)	0.0300 (18)	0.0017 (12)	0.0068 (14)	-0.0032 (13)
C19	0.0158 (16)	0.0151 (14)	0.0239 (16)	0.0016 (12)	0.0066 (13)	-0.0024 (12)
O11	0.0307 (14)	0.0418 (15)	0.0373 (14)	-0.0177 (11)	0.0211 (12)	-0.0165 (11)
O10	0.0385 (15)	0.0404 (14)	0.0280 (13)	0.0139 (12)	0.0184 (11)	0.0108 (11)
C17	0.0162 (16)	0.0181 (15)	0.0153 (14)	0.0015 (12)	0.0059 (12)	0.0001 (12)
C12	0.0199 (17)	0.0171 (15)	0.0272 (17)	-0.0006 (13)	0.0108 (14)	-0.0045 (13)
C16	0.0193 (17)	0.0188 (15)	0.0232 (16)	-0.0021 (12)	0.0090 (13)	-0.0043 (13)
C11	0.0144 (16)	0.0189 (15)	0.0205 (15)	-0.0001 (12)	0.0065 (13)	0.0003 (12)
C10	0.0178 (17)	0.0150 (15)	0.0227 (16)	-0.0015 (12)	0.0062 (13)	0.0010 (12)
C18	0.0144 (16)	0.0163 (14)	0.0185 (15)	-0.0019 (12)	0.0062 (12)	-0.0018 (12)
O8	0.0202 (12)	0.0202 (11)	0.0221 (11)	0.0063 (9)	0.0070 (9)	0.0064 (9)
O7	0.0288 (13)	0.0343 (13)	0.0225 (12)	0.0150 (10)	0.0151 (10)	0.0088 (10)
O5	0.0197 (12)	0.0208 (11)	0.0433 (14)	-0.0069 (9)	0.0155 (11)	-0.0133 (10)
O6	0.0162 (12)	0.0223 (11)	0.0380 (13)	-0.0038 (9)	0.0124 (10)	-0.0065 (10)
O12	0.0252 (14)	0.0405 (15)	0.0588 (17)	-0.0157 (11)	0.0233 (13)	-0.0302 (13)
C36	0.0173 (17)	0.0195 (15)	0.0186 (15)	-0.0020 (12)	0.0054 (13)	0.0015 (12)
C33	0.0130 (16)	0.0244 (16)	0.0213 (16)	0.0017 (12)	0.0066 (13)	-0.0002 (13)
C35	0.0150 (16)	0.0185 (15)	0.0189 (15)	-0.0016 (12)	0.0053 (13)	-0.0006 (12)
C34	0.0199 (17)	0.0160 (15)	0.0249 (16)	-0.0002 (12)	0.0083 (14)	-0.0042 (12)
O16	0.0219 (12)	0.0203 (11)	0.0351 (13)	-0.0064 (9)	0.0136 (10)	-0.0118 (10)
O15	0.0281 (14)	0.0341 (13)	0.0422 (15)	-0.0139 (11)	0.0232 (12)	-0.0182 (11)
O9	0.0184 (13)	0.0341 (13)	0.0370 (14)	0.0092 (10)	0.0106 (11)	0.0128 (11)
N7	0.0180 (15)	0.0282 (16)	0.0287 (16)	0.0006 (12)	0.0104 (12)	-0.0061 (12)
C28	0.0180 (16)	0.0127 (14)	0.0181 (15)	-0.0019 (12)	0.0052 (13)	-0.0002 (12)
C29	0.0155 (16)	0.0190 (15)	0.0146 (14)	-0.0001 (12)	0.0069 (12)	0.0003 (12)

## supplementary materials

---

C32	0.0122 (16)	0.0313 (18)	0.0259 (17)	-0.0018 (13)	0.0053 (13)	0.0008 (14)
O14	0.0258 (13)	0.0289 (12)	0.0184 (11)	0.0076 (10)	0.0097 (10)	0.0039 (9)
N8	0.0121 (14)	0.0244 (15)	0.0276 (15)	-0.0051 (11)	0.0039 (12)	-0.0046 (12)
C31	0.0120 (15)	0.0181 (15)	0.0187 (15)	-0.0029 (12)	0.0024 (12)	0.0000 (12)
C30	0.0177 (17)	0.0180 (15)	0.0216 (16)	-0.0004 (12)	0.0055 (13)	-0.0032 (12)
C1	0.0121 (16)	0.0158 (15)	0.0268 (17)	0.0017 (12)	0.0033 (13)	-0.0022 (12)
C2	0.0141 (16)	0.0158 (15)	0.0229 (16)	0.0003 (12)	0.0057 (13)	0.0001 (12)
O2	0.0235 (13)	0.0260 (12)	0.0282 (12)	-0.0070 (9)	0.0122 (10)	-0.0035 (10)
O1	0.0207 (13)	0.0270 (12)	0.0441 (14)	-0.0100 (10)	0.0186 (11)	-0.0172 (11)
C8	0.0141 (16)	0.0158 (14)	0.0165 (14)	0.0027 (12)	0.0033 (12)	0.0010 (11)
C7	0.0179 (16)	0.0194 (15)	0.0186 (15)	0.0029 (12)	0.0064 (13)	-0.0032 (12)
C4	0.0117 (16)	0.0212 (15)	0.0188 (15)	0.0003 (12)	0.0035 (12)	-0.0029 (12)
C6	0.0144 (16)	0.0192 (15)	0.0166 (15)	-0.0001 (12)	0.0029 (12)	-0.0004 (12)
C3	0.0138 (16)	0.0159 (15)	0.0276 (17)	0.0003 (12)	0.0057 (13)	-0.0060 (12)
N1	0.0104 (13)	0.0228 (13)	0.0229 (13)	-0.0027 (10)	0.0038 (11)	-0.0038 (11)
N2	0.0166 (14)	0.0196 (14)	0.0282 (15)	-0.0033 (11)	0.0060 (12)	-0.0092 (12)
C5	0.0154 (16)	0.0241 (16)	0.0240 (16)	-0.0015 (13)	0.0075 (13)	-0.0015 (13)
C9	0.0161 (16)	0.0110 (14)	0.0235 (16)	0.0036 (11)	0.0077 (13)	0.0013 (12)
O4	0.0261 (13)	0.0288 (12)	0.0214 (11)	0.0088 (10)	0.0128 (10)	0.0055 (9)
O13	0.0208 (12)	0.0322 (13)	0.0224 (12)	0.0071 (10)	0.0084 (10)	0.0112 (10)
O3	0.0131 (11)	0.0248 (11)	0.0218 (11)	0.0054 (9)	0.0038 (9)	0.0013 (9)
O1W	0.0313 (14)	0.0279 (12)	0.0216 (12)	-0.0129 (10)	0.0108 (10)	0.0001 (9)
O2W	0.0307 (14)	0.0532 (16)	0.0215 (12)	-0.0001 (12)	0.0108 (11)	-0.0028 (11)

### *Geometric parameters (Å, °)*

Gd1—O1	2.338 (2)	C16—H16	0.9300
Gd1—O2	2.526 (2)	C11—C10	1.495 (4)
Gd1—O3 <sup>i</sup>	2.350 (2)	C10—O6	1.247 (4)
Gd1—O5 <sup>ii</sup>	2.461 (2)	C10—O5	1.278 (4)
Gd1—O6 <sup>ii</sup>	2.499 (2)	C18—O7	1.247 (4)
Gd1—O8	2.314 (2)	C18—O8	1.255 (3)
Gd1—N6	2.617 (3)	C36—O15	1.248 (4)
Gd1—O1W	2.374 (2)	C36—O16	1.271 (4)
Gd2—O9	2.312 (2)	C36—C35	1.491 (4)
Gd2—O11 <sup>iii</sup>	2.539 (2)	C33—C34	1.385 (4)
Gd2—O12 <sup>iii</sup>	2.342 (2)	C33—N7	1.387 (4)
Gd2—O13 <sup>iv</sup>	2.267 (2)	C33—C31	1.393 (4)
Gd2—O15	2.449 (2)	C35—C34	1.388 (4)
Gd2—O16	2.520 (2)	C35—C29	1.424 (4)
Gd2—N1 <sup>iii</sup>	2.612 (3)	C34—H34	0.9300
Gd2—O2W	2.384 (2)	N7—C32	1.323 (4)
N6—C23	1.316 (4)	N7—H7A	0.77 (4)
N6—C24	1.407 (4)	C28—O14	1.236 (4)
N5—C23	1.343 (4)	C28—O13	1.272 (4)
N5—C22	1.372 (4)	C28—C29	1.516 (4)
N5—H5A	0.83 (4)	C29—C30	1.380 (4)
C23—H23	0.9300	C32—N8	1.316 (4)

C25—C24	1.389 (4)	C32—H32	0.9300
C25—C26	1.391 (4)	N8—C31	1.382 (4)
C25—H25	0.9300	N8—H8A	0.80 (4)
C21—C20	1.385 (4)	C31—C30	1.390 (4)
C21—C22	1.390 (4)	C30—H30	0.9300
C21—H21	0.9300	C1—O2	1.247 (4)
C24—C22	1.400 (4)	C1—O1	1.281 (4)
C20—C26	1.411 (4)	C1—C2	1.502 (4)
C20—C19	1.504 (4)	C2—C3	1.380 (4)
C26—C27	1.498 (4)	C2—C8	1.420 (4)
N3—C14	1.314 (4)	C8—C7	1.378 (4)
N3—C15	1.378 (4)	C8—C9	1.510 (4)
N3—H3A	0.81 (4)	C7—C6	1.396 (4)
C13—C12	1.383 (4)	C7—H7	0.9300
C13—N4	1.391 (4)	C4—C3	1.392 (4)
C13—C15	1.392 (4)	C4—C6	1.400 (4)
N4—C14	1.313 (4)	C4—N1	1.403 (4)
N4—H4A	0.84 (4)	C6—N2	1.378 (4)
C15—C16	1.387 (4)	C3—H3	0.9300
C14—H14	0.9300	N1—C5	1.315 (4)
C27—O11	1.245 (4)	N2—C5	1.345 (4)
C27—O12	1.270 (4)	N2—H2A	0.82 (4)
C19—O10	1.251 (4)	C5—H5	0.9300
C19—O9	1.251 (4)	C9—O4	1.252 (4)
O11—Gd2 <sup>iv</sup>	2.539 (2)	C9—O3	1.266 (4)
C17—C16	1.378 (4)	O1W—H1W	0.84
C17—C11	1.420 (4)	O1W—H2W	0.82
C17—C18	1.513 (4)	O2W—H3W	0.86
C12—C11	1.386 (4)	O2W—H4WA	0.82
C12—H12	0.9300	O2W—H4WB	0.82
O8—Gd1—O1	107.71 (8)	N3—C14—H14	124.8
O8—Gd1—O3 <sup>i</sup>	80.39 (7)	O11—C27—O12	120.5 (3)
O1—Gd1—O3 <sup>i</sup>	156.84 (7)	O11—C27—C26	122.0 (3)
O8—Gd1—O1W	150.83 (7)	O12—C27—C26	117.5 (3)
O1—Gd1—O1W	89.84 (8)	O11—C27—Gd2 <sup>iv</sup>	64.70 (17)
O3 <sup>i</sup> —Gd1—O1W	75.13 (7)	O12—C27—Gd2 <sup>iv</sup>	55.80 (16)
O8—Gd1—O5 <sup>ii</sup>	91.14 (7)	C26—C27—Gd2 <sup>iv</sup>	173.3 (2)
O1—Gd1—O5 <sup>ii</sup>	127.16 (7)	O10—C19—O9	124.2 (3)
O3 <sup>i</sup> —Gd1—O5 <sup>ii</sup>	72.98 (7)	O10—C19—C20	117.1 (3)
O1W—Gd1—O5 <sup>ii</sup>	96.56 (8)	O9—C19—C20	118.6 (3)
O8—Gd1—O6 <sup>ii</sup>	133.85 (7)	C27—O11—Gd2 <sup>iv</sup>	88.98 (18)
O1—Gd1—O6 <sup>ii</sup>	81.62 (8)	C16—C17—C11	121.3 (3)
O3 <sup>i</sup> —Gd1—O6 <sup>ii</sup>	108.82 (7)	C16—C17—C18	115.8 (3)
O1W—Gd1—O6 <sup>ii</sup>	70.40 (8)	C11—C17—C18	123.0 (3)
O5 <sup>ii</sup> —Gd1—O6 <sup>ii</sup>	52.51 (7)	C13—C12—C11	117.9 (3)
O8—Gd1—O2	77.02 (7)	C13—C12—H12	121.1

## supplementary materials

---

O1—Gd1—O2	53.69 (7)	C11—C12—H12	121.1
O3 <sup>i</sup> —Gd1—O2	148.38 (7)	C17—C16—C15	117.1 (3)
O1W—Gd1—O2	131.50 (7)	C17—C16—H16	121.4
O5 <sup>ii</sup> —Gd1—O2	85.58 (8)	C15—C16—H16	121.4
O6 <sup>ii</sup> —Gd1—O2	73.23 (7)	C12—C11—C17	120.6 (3)
O8—Gd1—N6	87.31 (8)	C12—C11—C10	118.5 (3)
O1—Gd1—N6	78.24 (8)	C17—C11—C10	120.9 (3)
O3 <sup>i</sup> —Gd1—N6	80.57 (8)	O6—C10—O5	120.7 (3)
O1W—Gd1—N6	73.35 (8)	O6—C10—C11	120.3 (3)
O5 <sup>ii</sup> —Gd1—N6	153.38 (8)	O5—C10—C11	118.9 (3)
O6 <sup>ii</sup> —Gd1—N6	138.26 (8)	O6—C10—Gd1 <sup>i</sup>	61.93 (15)
O2—Gd1—N6	119.76 (8)	O5—C10—Gd1 <sup>i</sup>	60.23 (15)
O8—Gd1—C10 <sup>ii</sup>	115.53 (8)	C11—C10—Gd1 <sup>i</sup>	167.3 (2)
O1—Gd1—C10 <sup>ii</sup>	106.09 (8)	O7—C18—O8	124.7 (3)
O3 <sup>i</sup> —Gd1—C10 <sup>ii</sup>	88.80 (8)	O7—C18—C17	117.4 (3)
O1W—Gd1—C10 <sup>ii</sup>	79.89 (8)	O8—C18—C17	117.8 (3)
O5 <sup>ii</sup> —Gd1—C10 <sup>ii</sup>	26.78 (8)	C18—O8—Gd1	148.6 (2)
O6 <sup>ii</sup> —Gd1—C10 <sup>ii</sup>	26.11 (8)	C10—O5—Gd1 <sup>i</sup>	92.99 (18)
O2—Gd1—C10 <sup>ii</sup>	81.39 (8)	C10—O6—Gd1 <sup>i</sup>	91.96 (17)
N6—Gd1—C10 <sup>ii</sup>	152.92 (8)	C27—O12—Gd2 <sup>iv</sup>	97.54 (19)
O13 <sup>iv</sup> —Gd2—O9	87.94 (8)	O15—C36—O16	120.3 (3)
O13 <sup>iv</sup> —Gd2—O12 <sup>iii</sup>	107.11 (9)	O15—C36—C35	119.2 (3)
O9—Gd2—O12 <sup>iii</sup>	152.90 (8)	O16—C36—C35	120.5 (3)
O13 <sup>iv</sup> —Gd2—O2W	153.97 (8)	C34—C33—N7	132.7 (3)
O9—Gd2—O2W	79.66 (9)	C34—C33—C31	121.1 (3)
O12 <sup>iii</sup> —Gd2—O2W	77.24 (9)	N7—C33—C31	106.1 (3)
O13 <sup>iv</sup> —Gd2—O15	136.70 (8)	C34—C35—C29	120.9 (3)
O9—Gd2—O15	98.56 (8)	C34—C35—C36	118.6 (3)
O12 <sup>iii</sup> —Gd2—O15	85.94 (9)	C29—C35—C36	120.6 (3)
O2W—Gd2—O15	68.38 (8)	C33—C34—C35	117.8 (3)
O13 <sup>iv</sup> —Gd2—O16	89.20 (7)	C33—C34—H34	121.1
O9—Gd2—O16	75.74 (8)	C35—C34—H34	121.1
O12 <sup>iii</sup> —Gd2—O16	125.56 (8)	C36—O16—Gd2	91.50 (18)
O2W—Gd2—O16	109.44 (8)	C36—O15—Gd2	95.47 (18)
O15—Gd2—O16	52.14 (7)	C19—O9—Gd2	168.9 (2)
O13 <sup>iv</sup> —Gd2—O11 <sup>iii</sup>	80.34 (8)	C32—N7—C33	108.5 (3)
O9—Gd2—O11 <sup>iii</sup>	153.99 (8)	C32—N7—H7A	129 (3)
O12 <sup>iii</sup> —Gd2—O11 <sup>iii</sup>	52.97 (8)	C33—N7—H7A	123 (3)
O2W—Gd2—O11 <sup>iii</sup>	119.51 (8)	O14—C28—O13	123.9 (3)
O15—Gd2—O11 <sup>iii</sup>	75.38 (9)	O14—C28—C29	119.9 (3)
O16—Gd2—O11 <sup>iii</sup>	80.92 (8)	O13—C28—C29	116.1 (3)
O13 <sup>iv</sup> —Gd2—N1 <sup>iii</sup>	82.99 (8)	C30—C29—C35	120.7 (3)

O9—Gd2—N1 <sup>iii</sup>	79.84 (8)	C30—C29—C28	116.4 (3)
O12 <sup>iii</sup> —Gd2—N1 <sup>iii</sup>	79.83 (8)	C35—C29—C28	122.8 (3)
O2W—Gd2—N1 <sup>iii</sup>	72.39 (8)	N8—C32—N7	110.4 (3)
O15—Gd2—N1 <sup>iii</sup>	140.32 (8)	N8—C32—H32	124.8
O16—Gd2—N1 <sup>iii</sup>	154.59 (7)	N7—C32—H32	124.8
O11 <sup>iii</sup> —Gd2—N1 <sup>iii</sup>	121.11 (8)	C32—N8—C31	108.8 (3)
O13 <sup>iv</sup> —Gd2—C27 <sup>iii</sup>	93.85 (9)	C32—N8—H8A	122 (3)
O9—Gd2—C27 <sup>iii</sup>	178.09 (9)	C31—N8—H8A	130 (3)
O12 <sup>iii</sup> —Gd2—C27 <sup>iii</sup>	26.65 (9)	N8—C31—C30	131.9 (3)
O2W—Gd2—C27 <sup>iii</sup>	98.94 (9)	N8—C31—C33	106.3 (3)
O15—Gd2—C27 <sup>iii</sup>	79.68 (9)	C30—C31—C33	121.7 (3)
O16—Gd2—C27 <sup>iii</sup>	103.62 (8)	C29—C30—C31	117.8 (3)
O11 <sup>iii</sup> —Gd2—C27 <sup>iii</sup>	26.32 (8)	C29—C30—H30	121.1
N1 <sup>iii</sup> —Gd2—C27 <sup>iii</sup>	101.01 (8)	C31—C30—H30	121.1
C23—N6—C24	104.1 (3)	O2—C1—O1	121.3 (3)
C23—N6—Gd1	120.7 (2)	O2—C1—C2	121.7 (3)
C24—N6—Gd1	132.91 (19)	O1—C1—C2	117.0 (3)
C23—N5—C22	107.6 (3)	C3—C2—C8	121.1 (3)
C23—N5—H5A	126 (3)	C3—C2—C1	117.9 (3)
C22—N5—H5A	127 (3)	C8—C2—C1	121.0 (3)
N6—C23—N5	113.7 (3)	C1—O2—Gd1	88.55 (18)
N6—C23—H23	123.1	C1—O1—Gd1	96.42 (18)
N5—C23—H23	123.1	C7—C8—C2	120.3 (3)
C24—C25—C26	119.0 (3)	C7—C8—C9	117.6 (3)
C24—C25—H25	120.5	C2—C8—C9	121.6 (3)
C26—C25—H25	120.5	C8—C7—C6	118.0 (3)
C20—C21—C22	117.9 (3)	C8—C7—H7	121.0
C20—C21—H21	121.1	C6—C7—H7	121.0
C22—C21—H21	121.1	C3—C4—C6	119.5 (3)
C25—C24—C22	119.3 (3)	C3—C4—N1	130.8 (3)
C25—C24—N6	131.4 (3)	C6—C4—N1	109.6 (3)
C22—C24—N6	109.2 (3)	N2—C6—C7	132.6 (3)
N5—C22—C21	132.2 (3)	N2—C6—C4	105.3 (3)
N5—C22—C24	105.3 (3)	C7—C6—C4	122.1 (3)
C21—C22—C24	122.4 (3)	C2—C3—C4	118.9 (3)
C21—C20—C26	120.4 (3)	C2—C3—H3	120.5
C21—C20—C19	117.2 (3)	C4—C3—H3	120.5
C26—C20—C19	122.3 (3)	C5—N1—C4	103.9 (2)
C25—C26—C20	120.9 (3)	C5—N1—Gd2 <sup>iv</sup>	122.2 (2)
C25—C26—C27	117.3 (3)	C4—N1—Gd2 <sup>iv</sup>	132.34 (19)
C20—C26—C27	121.7 (3)	C5—N2—C6	106.9 (3)
C14—N3—C15	109.0 (3)	C5—N2—H2A	127 (3)
C14—N3—H3A	128 (3)	C6—N2—H2A	126 (3)
C15—N3—H3A	123 (3)	N1—C5—N2	114.3 (3)
C12—C13—N4	132.9 (3)	N1—C5—H5	122.9
C12—C13—C15	121.0 (3)	N2—C5—H5	122.9

## supplementary materials

---

N4—C13—C15	106.1 (3)	O4—C9—O3	124.8 (3)
C14—N4—C13	108.4 (3)	O4—C9—C8	118.8 (3)
C14—N4—H4A	127 (3)	O3—C9—C8	116.3 (3)
C13—N4—H4A	124 (3)	C28—O13—Gd2 <sup>iii</sup>	151.0 (2)
N3—C15—C16	131.9 (3)	C9—O3—Gd1 <sup>ii</sup>	135.77 (18)
N3—C15—C13	106.0 (3)	H1W—O1W—H2W	108
C16—C15—C13	122.1 (3)	H3W—O2W—H4WA	106
N4—C14—N3	110.4 (3)	H3W—O2W—H4WB	105
N4—C14—H14	124.8		

Symmetry codes: (i)  $-x, y-1/2, -z+1/2$ ; (ii)  $-x, y+1/2, -z+1/2$ ; (iii)  $-x+1, y-1/2, -z+1/2$ ; (iv)  $-x+1, y+1/2, -z+1/2$ .

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1W—H1W <sup>v</sup> ...O7 <sup>v</sup>	0.84	1.80	2.629 (3)	167
O1W—H2W...O4 <sup>i</sup>	0.82	1.92	2.670 (3)	153
O2W—H3W...O10 <sup>vi</sup>	0.86	1.74	2.590 (3)	172
O2W—H4WA...O15	0.82	2.14	2.716 (5)	127
O2W—H4WB...O2W <sup>vi</sup>	0.82	1.91	2.723 (4)	170
N2—H2A...O14 <sup>vii</sup>	0.82 (4)	1.94 (4)	2.742 (3)	169 (3)
N3—H3A...O10 <sup>vii</sup>	0.81 (4)	1.94 (4)	2.731 (4)	166 (4)
N4—H4A...O16	0.84 (4)	1.97 (4)	2.801 (4)	170 (4)
N5—H5A...O4 <sup>v</sup>	0.83 (4)	1.96 (4)	2.772 (3)	165 (4)
N7—H7A...O5	0.77 (4)	2.08 (4)	2.845 (3)	171 (4)
N8—H8A...O7 <sup>viii</sup>	0.80 (4)	1.95 (4)	2.747 (3)	172 (4)

Symmetry codes: (v)  $x, -y+1/2, z-1/2$ ; (i)  $-x, y-1/2, -z+1/2$ ; (vi)  $-x+1, -y, -z$ ; (vii)  $x, -y+1/2, z+1/2$ ; (viii)  $x, -y-1/2, z-1/2$ .



Fig. 2

