

# Poly[[tetraaquabis( $\mu_3$ -1*H*-benzimidazole-5,6-dicarboxylato)dicobalt(II)] trihydrate]

Jun-Dan Fu, Zhi-Wei Tang, Ming-Yue Yuan and Yi-Hang Wen\*

Zhejiang Key Laboratory for Reactive Chemistry on Solid Surfaces, Institute of Physical Chemistry, Zhejiang Normal University, Jinhua, Zhejiang 321004, People's Republic of China

Correspondence e-mail: wyh@zjnu.edu.cn


Received 5 November 2009; accepted 18 November 2009

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.029;  $wR$  factor = 0.087; data-to-parameter ratio = 15.3.

The title complex,  $\{[\text{Co}_2(\text{C}_9\text{H}_4\text{N}_2\text{O}_4)_2(\text{H}_2\text{O})_4]\cdot 3\text{H}_2\text{O}\}_n$ , was synthesized hydrothermally. The unique  $\text{Co}^{\text{II}}$  ion is coordinated in a distorted octahedral coordination environment by two water molecules and three symmetry-related 1*H*-benzimidazole-5,6-dicarboxylate (Hbdc) ligands. The Hbdc ligands coordinate *via* a bis-chelating and mono-chelating carboxylate group and by an imidazole group N atom, bridging the  $\text{Co}^{\text{II}}$  ions and forming an extended two-dimensional structure in the *ab* plane. In the crystal structure, intermolecular  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds connect complex and solvent water molecules, forming a three-dimensional supermolecular network. One of the solvent water molecules lies on a twofold rotation axis.

## Related literature

For background information on carboxylate ligands in coordination chemistry, see: Laduca (2009); Grodzicki *et al.* (2005). For the isostructural Ni(II) complex, see: Yao *et al.* (2008). For related structures, see: Wei *et al.* (2008); Xu & Yu (2009).



## Experimental

### Crystal data

$[\text{Co}_2(\text{C}_9\text{H}_4\text{N}_2\text{O}_4)_2(\text{H}_2\text{O})_4]\cdot 3\text{H}_2\text{O}$	$V = 2302.2(3) \text{ \AA}^3$
$M_r = 652.26$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 22.4085(18) \text{ \AA}$	$\mu = 1.53 \text{ mm}^{-1}$
$b = 9.1564(7) \text{ \AA}$	$T = 296 \text{ K}$
$c = 13.0907(10) \text{ \AA}$	$0.43 \times 0.25 \times 0.07 \text{ mm}$
$\beta = 121.006(4)^\circ$	

### Data collection

Bruker APEXII diffractometer	9315 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	2656 independent reflections
$T_{\text{min}} = 0.63$ , $T_{\text{max}} = 0.90$	2402 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.018$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$	6 restraints
$wR(F^2) = 0.087$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.81 \text{ e \AA}^{-3}$
2656 reflections	$\Delta\rho_{\text{min}} = -0.63 \text{ e \AA}^{-3}$
174 parameters	

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O3W}-\text{H3WA}\cdots\text{O1}^{\text{i}}$	0.84	1.97	2.798 (2)	170
$\text{N1}-\text{H1A}\cdots\text{O1W}^{\text{i}}$	0.86	2.45	3.130 (2)	136
$\text{O2W}-\text{H2WA}\cdots\text{O4}^{\text{i}}$	0.84	1.99	2.786	157
$\text{O3W}-\text{H3WB}\cdots\text{O3}^{\text{ii}}$	0.84	1.85	2.641 (2)	158
$\text{O4W}-\text{H4WB}\cdots\text{O3}^{\text{iii}}$	0.84	2.60	3.095 (2)	119
$\text{O4W}-\text{H4WA}\cdots\text{O2}^{\text{iii}}$	0.84	1.86	2.679 (2)	165
$\text{O4W}-\text{H4WB}\cdots\text{O2W}^{\text{iv}}$	0.84	2.02	2.769	148
$\text{N1}-\text{H1A}\cdots\text{O2W}$	0.86	2.31	2.899	125
$\text{O1W}-\text{H1W}\cdots\text{O3}^{\text{v}}$	0.84	1.98	2.8218 (18)	180
$\text{O2W}-\text{H2WB}\cdots\text{O3W}^{\text{vi}}$	0.84	2.16	2.949	157

Symmetry codes: (i)  $-x, -y + 1, -z$ ; (ii)  $-x, -y, -z$ ; (iii)  $-x, y, -z - \frac{1}{2}$ ; (iv)  $x, y - 1, z$ ; (v)  $x, -y, z + \frac{1}{2}$ ; (vi)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2948).

## References

- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Grodzicki, A., Lakomski, I., Piszczek, P., Szymanska, I. & Szlyk, Z. (2005). *Coord. Chem. Rev.* **249**, 2232–2258.
- Laduca, R. L. (2009). *Coord. Chem. Rev.* **253**, 1759–1792.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wei, Y. Q., Yu, Y. F. & Wu, K. C. (2008). *Cryst. Growth Des.* **8**, 2087–2089.
- Xu, K. & Yu, L.-P. (2009). *Acta Cryst.* **E65**, m295.
- Yao, Y. L., Che, Y. X. & Zheng, J. M. (2008). *Cryst. Growth Des.* **8**, 2299–2306.

**supplementary materials**

*Acta Cryst.* (2009). E65, m1657 [ doi:10.1107/S1600536809049083 ]

## Poly[[tetraaquabis( $\mu_3$ -1*H*-benzimidazole-5,6-dicarboxylato)dicobalt(II)] trihydrate]

J.-D. Fu, Z.-W. Tang, M.-Y. Yuan and Y.-H. Wen

### Comment

It is well known that carboxylate ligands play an important role in coordination chemistry (Grodzicki *et al.*, 2005; Laduca, 2009). In recent years, the interaction of Hbdc with several metal ions has been studied, due to its unique ability to form stable chelates in diverse coordination modes such as bidentate, meridian and bridging (Wei *et al.*, 2008; Yao *et al.*, 2008). Herein we report the synthesis and crystal structure of the title two-dimensional complex of Hbdc (I).

Part of the 2-D structure of (I) is shown in Fig.1. The unique Co<sup>II</sup> ion is six-coordinated by one N atom and three O atoms from three symmetry related Hbdc ligands and two oxygen atoms from two water molecules. Each Hbdc ligand coordinates via a chelating carboxylate group and a single oxygen atom of another carboxylate group bridging two Co<sup>II</sup> ions to form a one-dimensional chain along the *b*-axis with a Co $\cdots$ Co separation of 5.4374 (5) Å. In addition, Hbdc ligands coordinate through a N atom to connect the adjacent chains forming a two-dimensional network with chains separated by ca. 7.06 Å (see Fig. 2a). In the crystal structure, intermolecular N-H $\cdots$ O and O-H $\cdots$ O hydrogen bonds connect complex and solvent water molecules to form a three-dimensional supermolecular network (see Table 1 and Fig. 2b).

### Experimental

A mixture of CoSO<sub>4</sub>·7H<sub>2</sub>O(0.141 g, 0.5 mmol), benzimidazole-5,6-dicarboxylic acid(0.103 g, 0.5 mmol), H<sub>2</sub>O(16 ml)and 4-sulfophthalic(1 ml)(solution pH = 5) was sealed in a 25 ml Teflon-lined stainless steel reactor and heated at 393 k for 3 d. On completion of the reaction, the reactor was cooled slowly to room temperature and the mixture was filtered, giving red single crystals suitable for X-ray analysis in 30% yield.

### Refinement

H-atoms were positioned geometrically and included in the refinement using a riding-model approximation [C–H = 0.93, O–H = 0.84 and N–H = 0.86Å] with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C},\text{N})$  or  $1.5U_{\text{eq}}(\text{O})$ .

### Figures




Fig. 1. Part of the 2-D structure of (I), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. [Symmetry codes: (A)-*x*, -*y*, -*z*; (B)*x* + 1/2, -*y* + 1/2, *z* + 1/2; (C)*x* - 1/2, -*y* + 1/2, *z* - 1/2.




Fig. 2. (a) Part of the 2-D structure of (I) viewed along the crystallographic *c*-axis. (b) Part of the crystal structure showing the donor...acceptor atom distances of hydrogen bonds as dashed lines.

**Poly[[tetraaquabis( $\mu_3$ -1*H*-benzimidazole-5,6-dicarboxylato)dicobalt(II)] trihydrate]**

*Crystal data*

[Co<sub>2</sub>(C<sub>9</sub>H<sub>4</sub>N<sub>2</sub>O<sub>4</sub>)<sub>2</sub>(H<sub>2</sub>O)<sub>4</sub>]<sub>3</sub>·3H<sub>2</sub>O

*M<sub>r</sub>* = 652.26

Monoclinic, *C2/c*

Hall symbol: -*C* 2yc

*a* = 22.4085 (18) Å

*b* = 9.1564 (7) Å

*c* = 13.0907 (10) Å

β = 121.006 (4)°

*V* = 2302.2 (3) Å<sup>3</sup>

*Z* = 4

*F*<sub>000</sub> = 1328

*D<sub>x</sub>* = 1.882 Mg m<sup>-3</sup>

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 5695 reflections

θ = 2.1–27.6°

μ = 1.53 mm<sup>-1</sup>

*T* = 296 K

Block, red

0.43 × 0.25 × 0.07 mm

*Data collection*

Bruker APEXII  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

*T* = 296 K

ω scans

Absorption correction: multi-scan  
(SADABS; Sheldrick, 1996)

*T<sub>min</sub>* = 0.63, *T<sub>max</sub>* = 0.90

9315 measured reflections

2656 independent reflections

2402 reflections with *I* > 2σ(*I*)

*R<sub>int</sub>* = 0.018

θ<sub>max</sub> = 27.6°

θ<sub>min</sub> = 2.1°

*h* = -29→27

*k* = -11→11

*l* = -16→17

*Refinement*

Refinement on *F*<sup>2</sup>

Least-squares matrix: full

*R*[*F*<sup>2</sup> > 2σ(*F*<sup>2</sup>)] = 0.029

*wR*(*F*<sup>2</sup>) = 0.087

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

*w* = 1/[σ<sup>2</sup>(*F<sub>o</sub>*<sup>2</sup>) + (0.0533*P*)<sup>2</sup> + 3.4124*P*]

where *P* = (*F<sub>o</sub>*<sup>2</sup> + 2*F<sub>c</sub>*<sup>2</sup>)/3

$S = 1.03$   $(\Delta/\sigma)_{\max} = 0.001$   
 2656 reflections  $\Delta\rho_{\max} = 0.81 \text{ e } \text{\AA}^{-3}$   
 174 parameters  $\Delta\rho_{\min} = -0.63 \text{ e } \text{\AA}^{-3}$   
 6 restraints Extinction correction: none  
 Primary atom site location: structure-invariant direct methods

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.199951 (12)	0.14498 (3)	0.12124 (2)	0.01825 (10)
O1	-0.19686 (7)	0.44398 (15)	-0.25564 (13)	0.0239 (3)
O2	-0.20916 (7)	0.22564 (15)	-0.33254 (12)	0.0216 (3)
O3	-0.10064 (8)	-0.07292 (17)	-0.23544 (16)	0.0335 (4)
O3W	0.20211 (8)	0.25089 (16)	0.26967 (13)	0.0277 (3)
H3WA	0.1974	0.3419	0.2690	0.042*
H3WB	0.1706	0.2071	0.2742	0.042*
O4	-0.14121 (7)	0.02445 (15)	-0.12688 (12)	0.0224 (3)
O4W	0.20838 (8)	0.03724 (17)	-0.01154 (13)	0.0302 (3)
H4WA	0.2019	0.0885	-0.0696	0.045*
H4WB	0.1863	-0.0415	-0.0365	0.045*
N1	0.07455 (8)	0.50513 (18)	-0.06630 (15)	0.0222 (3)
H1A	0.0716	0.5976	-0.0794	0.027*
N2	0.12237 (8)	0.28672 (18)	0.00087 (15)	0.0225 (3)
C1	-0.09320 (9)	0.3057 (2)	-0.19386 (16)	0.0181 (4)
C2	-0.06298 (9)	0.1649 (2)	-0.15733 (16)	0.0175 (4)
C3	0.00868 (10)	0.1468 (2)	-0.09321 (18)	0.0207 (4)
H3A	0.0285	0.0544	-0.0700	0.025*
C4	0.05016 (9)	0.2710 (2)	-0.06466 (17)	0.0195 (4)
C5	0.01982 (10)	0.4091 (2)	-0.10514 (17)	0.0189 (4)
C6	-0.05185 (9)	0.4293 (2)	-0.16866 (17)	0.0195 (4)
H6A	-0.0714	0.5216	-0.1933	0.023*
C7	-0.17032 (9)	0.3262 (2)	-0.26428 (16)	0.0177 (3)
C8	-0.10569 (9)	0.0287 (2)	-0.17727 (16)	0.0194 (4)
C9	0.13324 (10)	0.4278 (2)	-0.00439 (18)	0.0240 (4)
H9A	0.1774	0.4691	0.0315	0.029*

## supplementary materials

---

O1W	0.0000	0.24202 (17)	0.2500	0.1066 (18)
H1W	-0.0300	0.1917	0.2543	0.160*
O2W	0.1643	0.75912 (17)	0.0016	0.0530 (5)
H2WA	0.1492	0.8072	0.0382	0.080*
H2WB	0.2053	0.7403	0.0558	0.080*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Co1	0.01193 (15)	0.01637 (16)	0.02247 (16)	-0.00183 (8)	0.00601 (11)	0.00077 (9)
O1	0.0152 (6)	0.0189 (7)	0.0326 (7)	0.0021 (5)	0.0086 (6)	-0.0031 (6)
O2	0.0145 (6)	0.0201 (7)	0.0250 (7)	-0.0001 (5)	0.0064 (5)	-0.0035 (5)
O3	0.0321 (8)	0.0251 (8)	0.0530 (10)	-0.0099 (6)	0.0289 (8)	-0.0140 (7)
O3W	0.0269 (8)	0.0208 (7)	0.0381 (8)	-0.0053 (6)	0.0186 (7)	-0.0047 (6)
O4	0.0197 (6)	0.0207 (7)	0.0290 (7)	-0.0043 (5)	0.0139 (6)	-0.0005 (5)
O4W	0.0398 (9)	0.0241 (7)	0.0287 (7)	-0.0068 (6)	0.0191 (7)	-0.0012 (6)
N1	0.0166 (8)	0.0159 (8)	0.0313 (9)	-0.0036 (6)	0.0104 (7)	-0.0003 (6)
N2	0.0118 (7)	0.0221 (8)	0.0271 (8)	-0.0023 (6)	0.0054 (6)	0.0023 (7)
C1	0.0134 (8)	0.0187 (9)	0.0201 (8)	0.0002 (7)	0.0072 (7)	0.0008 (7)
C2	0.0138 (8)	0.0164 (9)	0.0202 (8)	-0.0008 (6)	0.0072 (7)	0.0011 (7)
C3	0.0148 (9)	0.0162 (9)	0.0263 (9)	0.0010 (6)	0.0072 (7)	0.0029 (7)
C4	0.0123 (8)	0.0212 (9)	0.0219 (9)	-0.0009 (7)	0.0067 (7)	0.0021 (7)
C5	0.0171 (9)	0.0161 (9)	0.0225 (9)	-0.0024 (7)	0.0095 (7)	0.0000 (7)
C6	0.0163 (8)	0.0159 (9)	0.0244 (9)	0.0015 (7)	0.0091 (7)	0.0023 (7)
C7	0.0135 (8)	0.0176 (8)	0.0208 (9)	0.0006 (7)	0.0079 (7)	0.0026 (7)
C8	0.0119 (8)	0.0173 (9)	0.0237 (9)	-0.0013 (7)	0.0054 (7)	0.0017 (7)
C9	0.0150 (9)	0.0241 (10)	0.0289 (10)	-0.0037 (7)	0.0084 (8)	-0.0001 (8)
O1W	0.103 (3)	0.0399 (18)	0.247 (6)	0.000	0.140 (4)	0.000
O2W	0.0709 (14)	0.0310 (10)	0.0514 (11)	-0.0099 (9)	0.0275 (10)	-0.0037 (8)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Co1—O4 <sup>i</sup>	2.0603 (14)	N1—C5	1.376 (2)
Co1—N2	2.0898 (16)	N1—H1A	0.8598
Co1—O4W	2.0901 (15)	N2—C9	1.322 (3)
Co1—O3W	2.1497 (15)	N2—C4	1.395 (2)
Co1—O2 <sup>ii</sup>	2.1560 (13)	C1—C6	1.390 (3)
Co1—O1 <sup>ii</sup>	2.1837 (14)	C1—C2	1.420 (3)
Co1—C7 <sup>ii</sup>	2.5057 (18)	C1—C7	1.493 (2)
O1—C7	1.265 (2)	C2—C3	1.386 (3)
O1—Co1 <sup>iii</sup>	2.1837 (14)	C2—C8	1.511 (2)
O2—C7	1.266 (2)	C3—C4	1.393 (3)
O2—Co1 <sup>iii</sup>	2.1560 (13)	C3—H3A	0.9300
O3—C8	1.244 (2)	C4—C5	1.404 (3)
O3W—H3WA	0.8399	C5—C6	1.389 (3)
O3W—H3WB	0.8399	C6—H6A	0.9300
O4—C8	1.269 (2)	C7—Co1 <sup>iii</sup>	2.5057 (18)

O4—Co1 <sup>i</sup>	2.0603 (14)	C9—H9A	0.9300
O4W—H4WA	0.8401	O1W—H1W	0.8401
O4W—H4WB	0.8393	O2W—H2WA	0.8400
N1—C9	1.338 (3)	O2W—H2WB	0.8401
O4 <sup>i</sup> —Co1—N2	101.30 (6)	C9—N2—Co1	122.80 (13)
O4 <sup>i</sup> —Co1—O4W	90.43 (6)	C4—N2—Co1	130.58 (13)
N2—Co1—O4W	93.58 (7)	C6—C1—C2	121.02 (16)
O4 <sup>i</sup> —Co1—O3W	91.41 (6)	C6—C1—C7	117.55 (16)
N2—Co1—O3W	91.38 (6)	C2—C1—C7	121.40 (16)
O4W—Co1—O3W	174.27 (6)	C3—C2—C1	121.02 (16)
O4 <sup>i</sup> —Co1—O2 <sup>ii</sup>	158.80 (5)	C3—C2—C8	116.03 (16)
N2—Co1—O2 <sup>ii</sup>	99.73 (6)	C1—C2—C8	122.80 (16)
O4W—Co1—O2 <sup>ii</sup>	90.90 (6)	C2—C3—C4	118.00 (17)
O3W—Co1—O2 <sup>ii</sup>	85.43 (5)	C2—C3—H3A	121.0
O4 <sup>i</sup> —Co1—O1 <sup>ii</sup>	98.39 (5)	C4—C3—H3A	121.0
N2—Co1—O1 <sup>ii</sup>	160.30 (6)	C3—C4—N2	130.70 (18)
O4W—Co1—O1 <sup>ii</sup>	85.57 (6)	C3—C4—C5	120.52 (17)
O3W—Co1—O1 <sup>ii</sup>	88.79 (6)	N2—C4—C5	108.77 (16)
O2 <sup>ii</sup> —Co1—O1 <sup>ii</sup>	60.64 (5)	N1—C5—C6	132.21 (18)
O4 <sup>i</sup> —Co1—C7 <sup>ii</sup>	128.59 (6)	N1—C5—C4	105.67 (16)
N2—Co1—C7 <sup>ii</sup>	130.06 (6)	C6—C5—C4	122.11 (17)
O4W—Co1—C7 <sup>ii</sup>	88.46 (6)	C5—C6—C1	117.23 (17)
O3W—Co1—C7 <sup>ii</sup>	86.15 (6)	C5—C6—H6A	121.4
O2 <sup>ii</sup> —Co1—C7 <sup>ii</sup>	30.33 (6)	C1—C6—H6A	121.4
O1 <sup>ii</sup> —Co1—C7 <sup>ii</sup>	30.31 (6)	O1—C7—O2	119.96 (16)
C7—O1—Co1 <sup>iii</sup>	89.07 (11)	O1—C7—C1	119.97 (16)
C7—O2—Co1 <sup>iii</sup>	90.30 (11)	O2—C7—C1	120.07 (16)
Co1—O3W—H3WA	119.7	O1—C7—Co1 <sup>iii</sup>	60.62 (9)
Co1—O3W—H3WB	102.7	O2—C7—Co1 <sup>iii</sup>	59.36 (9)
H3WA—O3W—H3WB	111.6	C1—C7—Co1 <sup>iii</sup>	178.35 (14)
C8—O4—Co1 <sup>i</sup>	128.68 (13)	O3—C8—O4	125.06 (18)
Co1—O4W—H4WA	116.2	O3—C8—C2	118.29 (17)
Co1—O4W—H4WB	116.2	O4—C8—C2	116.53 (17)
H4WA—O4W—H4WB	109.6	N2—C9—N1	113.50 (17)
C9—N1—C5	107.24 (16)	N2—C9—H9A	123.3
C9—N1—H1A	126.4	N1—C9—H9A	123.3
C5—N1—H1A	126.4	H2WA—O2W—H2WB	102.3
C9—N2—C4	104.80 (16)		
O4 <sup>i</sup> —Co1—N2—C9	158.93 (16)	C3—C4—C5—N1	-177.40 (18)
O4W—Co1—N2—C9	-109.92 (17)	N2—C4—C5—N1	1.9 (2)
O3W—Co1—N2—C9	67.21 (17)	C3—C4—C5—C6	3.7 (3)
O2 <sup>ii</sup> —Co1—N2—C9	-18.39 (18)	N2—C4—C5—C6	-177.00 (18)
O1 <sup>ii</sup> —Co1—N2—C9	-23.1 (3)	N1—C5—C6—C1	179.8 (2)

## supplementary materials

C7 <sup>ii</sup> —Co1—N2—C9	-18.9 (2)	C4—C5—C6—C1	-1.6 (3)
O4 <sup>i</sup> —Co1—N2—C4	-3.25 (19)	C2—C1—C6—C5	-0.9 (3)
O4W—Co1—N2—C4	87.90 (18)	C7—C1—C6—C5	-178.77 (17)
O3W—Co1—N2—C4	-94.97 (18)	Co1 <sup>iii</sup> —O1—C7—O2	1.72 (18)
O2 <sup>ii</sup> —Co1—N2—C4	179.43 (17)	Co1 <sup>iii</sup> —O1—C7—C1	-178.23 (15)
O1 <sup>ii</sup> —Co1—N2—C4	174.76 (16)	Co1 <sup>iii</sup> —O2—C7—O1	-1.74 (18)
C7 <sup>ii</sup> —Co1—N2—C4	178.90 (16)	Co1 <sup>iii</sup> —O2—C7—C1	178.21 (15)
C6—C1—C2—C3	1.4 (3)	C6—C1—C7—O1	-30.8 (3)
C7—C1—C2—C3	179.21 (18)	C2—C1—C7—O1	151.34 (18)
C6—C1—C2—C8	176.81 (17)	C6—C1—C7—O2	149.27 (18)
C7—C1—C2—C8	-5.4 (3)	C2—C1—C7—O2	-28.6 (3)
C1—C2—C3—C4	0.6 (3)	Co1 <sup>i</sup> —O4—C8—O3	0.8 (3)
C8—C2—C3—C4	-175.08 (17)	Co1 <sup>i</sup> —O4—C8—C2	-175.08 (12)
C2—C3—C4—N2	177.8 (2)	C3—C2—C8—O3	-62.2 (2)
C2—C3—C4—C5	-3.1 (3)	C1—C2—C8—O3	122.2 (2)
C9—N2—C4—C3	177.6 (2)	C3—C2—C8—O4	114.0 (2)
Co1—N2—C4—C3	-17.9 (3)	C1—C2—C8—O4	-61.7 (2)
C9—N2—C4—C5	-1.6 (2)	C4—N2—C9—N1	0.7 (2)
Co1—N2—C4—C5	162.99 (14)	Co1—N2—C9—N1	-165.38 (14)
C9—N1—C5—C6	177.3 (2)	C5—N1—C9—N2	0.4 (2)
C9—N1—C5—C4	-1.4 (2)		

Symmetry codes: (i)  $-x, -y, -z$ ; (ii)  $x+1/2, -y+1/2, z+1/2$ ; (iii)  $x-1/2, -y+1/2, z-1/2$ .

### Hydrogen-bond geometry ( $\text{\AA}, ^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3W—H3WA $\cdots$ O1 <sup>iv</sup>	0.84	1.97	2.798 (2)	170
N1—H1A $\cdots$ O1W <sup>iv</sup>	0.86	2.45	3.130 (2)	136
O2W—H2WA $\cdots$ O4 <sup>iv</sup>	0.84	1.99	2.78546	157
O3W—H3WB $\cdots$ O3 <sup>i</sup>	0.84	1.85	2.641 (2)	158
O4W—H4WB $\cdots$ O3 <sup>v</sup>	0.84	2.60	3.095 (2)	119
O4W—H4WA $\cdots$ O2 <sup>v</sup>	0.84	1.86	2.679 (2)	165
O4W—H4WB $\cdots$ O2W <sup>vi</sup>	0.84	2.02	2.76864	148
N1—H1A $\cdots$ O2W	0.86	2.31	2.89857	125
O1W—H1W $\cdots$ O3 <sup>vii</sup>	0.84	1.98	2.8218 (18)	180
O2W—H2WB $\cdots$ O3W <sup>viii</sup>	0.84	2.16	2.94922	157

Symmetry codes: (iv)  $-x, -y+1, -z$ ; (i)  $-x, -y, -z$ ; (v)  $-x, y, -z-1/2$ ; (vi)  $x, y-1, z$ ; (vii)  $x, -y, z+1/2$ ; (viii)  $-x+1/2, y+1/2, -z+1/2$ .

Fig. 1




Fig. 2

