

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Carbonyl[4-(2,6-dimethylphenylamino)-pent-3-en-2-onato- κ^2N,O](triphenylphosphine- κP)rhodium(I) acetone hemisolvate

Gertruida J. S. Venter,* Gideon Steyl and Andreas Roodt

Department of Chemistry, University of the Free State, P.O. Box 339, Bloemfontein, 9300, South Africa

Correspondence e-mail: truidie@hotmail.com


Received 23 October 2009; accepted 13 November 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.004$ Å; disorder in solvent or counterion; R factor = 0.038; wR factor = 0.112; data-to-parameter ratio = 21.0.

In the title compound, $[Rh(C_{13}H_{16}NO)(C_{18}H_{15}P)(CO)] \cdot 0.5C_3H_6O$, the Rh atom exhibits a square-planar coordination geometry, being coordinated by the N and O atoms of the bidentate β -diketonato ligand, a P atom from the triphenylphosphine unit and a C atom from the carbonyl group. The asymmetric unit also contains a disordered half-molecule, lying about an inversion center, of the acetone solvate. Intermolecular C—H \cdots O hydrogen bonds are observed between a C—H group of the triphenylphosphine unit and a carbonyl O atom and between the methyl group of the enaminketonato backbone and the solvent O atom. In addition, an intramolecular interaction is observed between a C—H group of the triphenylphosphine unit and the O atom of the enaminketonato ligand.

Related literature

For synthetic background, see: Shaheen *et al.* (2006); Cornils & Hermann (1996); Bonati & Wilkinson (1964). For applications of rhodium(I) dicarbonyl complexes, see: Cornils & Herrmann (1996); Trzeciak & Ziolkowski (1994); Van Rooy *et al.* (1995). For related structures, see: Damoense *et al.* (1994); Varshavsky *et al.* (2001); Venter *et al.* (2009).



Experimental

Crystal data

$[Rh(C_{13}H_{16}NO)(C_{18}H_{15}P)(CO)] \cdot 0.5C_3H_6O$
 $M_r = 624.5$
 Monoclinic, $P2_1/c$
 $a = 16.8558$ (8) Å
 $b = 11.4028$ (5) Å
 $c = 16.4059$ (8) Å
 $\beta = 108.733$ (1)°
 $V = 2986.2$ (2) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.66$ mm⁻¹
 $T = 100$ K
 $0.31 \times 0.15 \times 0.11$ mm

Data collection

Bruker X8 APEXII 4K Kappa CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2004)
 $T_{min} = 0.822$, $T_{max} = 0.931$
 23158 measured reflections
 7425 independent reflections
 6081 reflections with $> 2\sigma(I)$
 $R_{int} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.112$
 $S = 1.09$
 7425 reflections
 354 parameters
 1 restraint
 H-atom parameters constrained
 $\Delta\rho_{max} = 1.54$ e Å⁻³
 $\Delta\rho_{min} = -1.32$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C332—H332 \cdots O12	0.95	2.38	3.201 (3)	144
C334—H334 \cdots O14 ⁱ	0.95	2.51	3.201 (4)	130
C1—H1B \cdots O01 ⁱⁱ	0.98	2.54	3.372 (9)	142

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Table 2

Comparative geometrical parameters (Å, °) for similar $[Rh(N,O\text{-bid})(CO)(PPh_3)]$ complexes.

Parameters	(I) ^a	(II) ^b	(III) ^c	(IV) ^d
Rh1—N11	2.077 (2)	2.069 (2)	2.045 (4)	2.045 (3)
Rh1—O12	2.027 (2)	2.028 (2)	2.044 (3)	2.045 (2)
Rh1—P13	2.2704 (7)	2.2635 (6)	2.275 (1)	2.281 (2)
Rh1—C14	1.812 (3)	1.807 (2)	1.784 (5)	1.804 (3)
C14—O14	1.147 (3)	1.152 (3)	1.142 (7)	1.148 (4)
N11 \cdots O12	2.885 (3)	2.885 (3)	2.826 (6)	2.841 (3)
N11—Rh1—O12	89.31 (9)	89.54 (8)	87.4 (1)	87.95 (8)
O12—Rh1—P13	85.95 (6)	84.97 (5)	89.7 (1)	89.91 (5)
P13—Rh1—C14	91.57 (9)	91.87 (7)	90.3 (2)	89.48 (9)
N11—Rh1—C14	93.1 (1)	93.6 (1)	92.6 (2)	92.6 (1)
N11—C2—C4—O12	-2.6 (2)	4.1 (2)	1.2 (4)	1.5 (2)
θ_E^e	155.77 (2)	156.39 (3)	156.0 (2)	156.23 (4)

Notes: (a) This work; (b) N,O -bid = 4-(2,3-dimethyl phenylamino)pent-3-en-2-onato (Venter *et al.*, 2009); (c) N,O -bid = 4-amino-pent-3-en-2-onato (Damoense *et al.*, 1994); (d) N,O -bid = 4-amino-1,1,1-trifluoro-pent-3-en-2-onato (Varshavsky *et al.*, 2001); (e) cone angle (Tolman, 1977).

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT-Plus (Bruker, 2004); data reduction: SAINT-Plus; program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg & Putz, 2005); software used to prepare material for publication: WinGX (Farrugia, 1999).

Financial assistance from the University of the Free State is gratefully acknowledged, while Mr Leo Kirsten is thanked for

the data collection. We also express our gratitude to SASOL and the South African National Research Foundation (SANRF/THRIP) for financial support of this project. Part of this material is based on work supported by the SA-NRF/THRIP under grant No. GUN 2068915. Opinions, findings, conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the SANRF.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2227).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Bonati, F. & Wilkinson, G. (1964). *J. Chem. Soc.* pp. 3156–3160.
- Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2004). *SAINTE-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2005). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cornils, B. & Herrmann, W. A. (1996). *Applied Homogeneous Catalysis with Organometallic Compounds. A Comprehensive Handbook*, pp. 412–413. Weinheim: VCH.
- Damoense, L. J., Purcell, W., Roodt, A. & Leipoldt, J. G. (1994). *Rhodium Express*, **5**, 10–13.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Shaheen, F., Marchio, L., Badshah, A. & Khosa, M. K. (2006). *Acta Cryst.* **E62**, o873–o874.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tolman, C. A. (1977). *Chem. Rev.* **77**, 313–348.
- Trzeciak, A. M. & Ziolkowski, J. J. (1994). *J. Organomet. Chem.* **464**, 107–111.
- Van Rooy, A., Orji, E. N., Kramer, P. G. J. & Van Leeuwen, P. W. M. N. (1995). *Organometallics* **14**, 34–43.
- Varshavsky, Y. S., Galding, M. R., Cherkasova, T. G., Podkorytov, I. S., Nikol'skii, A. B., Trzeciak, A. M., Olejnik, Z., Lis, T. & Ziolkowski, J. J. (2001). *J. Organomet. Chem.* **628**, 195–210.
- Venter, G. J. S., Steyl, G. & Roodt, A. (2009). *Acta Cryst.* **E65**, m1321–m1322.

supplementary materials

Acta Cryst. (2009). E65, m1606-m1607 [doi:10.1107/S160053680904817X]

Carbonyl[4-(2,6-dimethylphenylamino)pent-3-en-2-onato- κ^2N,O](triphenylphosphine- κP)rhodium(I) acetone hemisolvate

G. J. S. Venter, G. Steyl and A. Roodt

Comment

Rhodium(I) dicarbonyl complexes of the type $[\text{Rh}(L,L')(\text{CO})_2]$ containing chelating mono-anionic bidentate (L,L') ligands coordinated to rhodium *via* (O,O) donor atoms have been studied as catalyst precursors (Cornils & Herrmann, 1996; Trzeciak *et al.*, 1994; Van Rooy *et al.*, 1995). The investigation of these complexes is followed by complexes containing bidentate β -enaminoketonato ligands such as 4-(phenylamino)pent-3-en-2-onato (Phony) (Shaheen *et al.*, 2006) coordinated to rhodium *via* (N,O) donor atoms. It was suggested that only one CO-group in a $[\text{Rh}(N,O\text{-bid})(\text{CO})_2]$ -type complex will be substituted by triphenyl phosphine, with the product being one of two possible isomers (Bonati & Wilkinson, 1964). Since the N-donor atom has a larger *trans*-influence than the O-atom, the CO-group *trans* to the N-atom will be substituted. This is evident in the title compound, (I), where $[\text{Rh}(2,6\text{-diMe-Phony})(\text{CO})(\text{PPh}_3)]$ is formed by the substitution of the carbonyl ligand in the dicarbonyl rhodium(I) complex $[\text{Rh}(2,6\text{-diMe-Phony})(\text{CO})_2]$ by PPh_3 .

In the title complex (Fig. 1), the bond distances involving rhodium differ significantly from the distances in related complexes, with the exception of $[\text{Rh}(2,3\text{-diMe-Phony})(\text{CO})(\text{PPh}_3)]$ (Venter *et al.*, 2009), wherein all angles and distances are comparable to (I) (Table 2). The distance, Rh—N, in (I) is longer than in similar complexes while the Rh—O bond distance is shorter. This is due to the steric influence of the phenyl group connected to nitrogen in the title compound, as opposed to the hydrogen in the related complexes. The Rh—C and the carbonyl C—O distances are comparable with those distances in the related complexes (Table 2). The N—Rh—O bite angle is slightly larger than that observed in similar complexes found in the literature. The effective cone angle, θ_E , (Tolman, 1977) of $155.77(2)^\circ$ is similar to the angles in the related compounds. The acetone solvate was disordered and was located about an inversion center. Intermolecular and intramolecular hydrogen bonds of the type C—H \cdots O are observed in the structure.

Experimental

To a 5 ml acetone solution of $[\text{Rh}(2,6\text{-diMe-Phony})(\text{CO})_2]$ (0.0302 g, 83.61 μmol) was added PPh_3 (0.0219 g, 83.50 μmol) resulting in an immediate evolution of gas with the formation of the title compound. Crystallization from acetone produced yellow crystals of (I) in quantitative (0.0516 g, 100%) yield. IR (KBr): ν_{CO} 1971.1 cm^{-1} .

Refinement

The methyl and aromatic H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H = 0.95 and 0.98 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ and $1.2U_{\text{eq}}(\text{C})$, respectively. The methyl groups were generated to fit the difference electron density and the groups were then refined as rigid rotors. The acetone solvate was disordered about inversion center. The highest residual electron-density peak in the final difference map was located 0.65 Å from H316 and was essentially meaningless.

Figures




Fig. 1. Molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability displacement level. H atoms and disordered solvate molecule have been omitted for clarity.




Fig. 2. A view of the unit cell of (I) illustrating the C—H...O interactions; hydrogen atoms have been omitted for clarity.

Carbonyl[4-(2,6-dimethylphenylamino)pent-3-en-2-onato- κ^2N,O](triphenylphosphine- κP)rhodium(I) acetone hemisolvate

Crystal data

$[\text{Rh}(\text{C}_{13}\text{H}_{16}\text{NO})(\text{C}_{18}\text{H}_{15}\text{P})(\text{CO})] \cdot 0.5\text{C}_3\text{H}_6\text{O}$

$M_r = 624.5$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 16.8558\ (8)\ \text{\AA}$

$b = 11.4028\ (5)\ \text{\AA}$

$c = 16.4059\ (8)\ \text{\AA}$

$\beta = 108.7330\ (10)^\circ$

$V = 2986.2\ (2)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 1288$

$D_x = 1.389\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 6670 reflections

$\theta = 2.5\text{--}28.2^\circ$

$\mu = 0.66\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Cuboid, yellow

$0.31 \times 0.15 \times 0.11\ \text{mm}$

Data collection

Bruker X8 APEXII 4K Kappa CCD diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100\ \text{K}$

ω and φ scans

Absorption correction: multi-scan (SADABS; Bruker, 2004)

$T_{\min} = 0.822$, $T_{\max} = 0.931$

23158 measured reflections

7425 independent reflections

6081 reflections with $> 2\sigma(I)$

$R_{\text{int}} = 0.042$

$\theta_{\text{max}} = 28.3^\circ$

$\theta_{\text{min}} = 1.3^\circ$

$h = -21 \rightarrow 22$

$k = -15 \rightarrow 11$

$l = -21 \rightarrow 21$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.038$	H-atom parameters constrained
$wR(F^2) = 0.112$	$w = 1/[\sigma^2(F_o^2) + (0.0577P)^2 + 1.4787P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
7425 reflections	$(\Delta/\sigma)_{\max} = 0.002$
354 parameters	$\Delta\rho_{\max} = 1.54 \text{ e } \text{\AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -1.32 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Experimental. The intensity data was collected on a Bruker X8 ApexII 4 K Kappa CCD diffractometer using an exposure time of 60 s/frame. A total of 688 frames were collected with a frame width of 0.5° covering up to $\theta = 28.24^\circ$ with 99.8% completeness accomplished.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.7144 (2)	0.7625 (3)	0.4744 (2)	0.0257 (7)	
H1A	0.7697	0.7807	0.5152	0.039*	
H1B	0.6712	0.7804	0.5007	0.039*	
H1C	0.7048	0.8097	0.4222	0.039*	
C2	0.71032 (17)	0.6337 (2)	0.45128 (18)	0.0168 (5)	
C3	0.65757 (17)	0.5637 (2)	0.48382 (18)	0.0183 (6)	
H3	0.6324	0.6017	0.5209	0.022*	
C4	0.63880 (17)	0.4461 (2)	0.46740 (19)	0.0180 (6)	
C5	0.57863 (19)	0.3856 (3)	0.5045 (2)	0.0265 (7)	
H5A	0.5288	0.3609	0.4577	0.04*	
H5B	0.5621	0.4399	0.5425	0.04*	
H5C	0.6058	0.3167	0.5375	0.04*	
C14	0.84522 (17)	0.4446 (2)	0.32802 (17)	0.0166 (5)	
C111	0.80357 (17)	0.6734 (2)	0.37352 (18)	0.0164 (5)	
C112	0.76869 (19)	0.7250 (2)	0.2919 (2)	0.0223 (6)	
C113	0.8183 (2)	0.8000 (3)	0.2624 (2)	0.0271 (7)	
H113	0.7952	0.8365	0.2078	0.033*	
C114	0.9008 (2)	0.8226 (2)	0.3109 (2)	0.0292 (7)	

supplementary materials

H114	0.9338	0.8746	0.2897	0.035*
C115	0.93503 (19)	0.7690 (2)	0.3902 (2)	0.0242 (6)
H115	0.9919	0.7836	0.4228	0.029*
C116	0.88743 (18)	0.6938 (2)	0.42310 (19)	0.0194 (6)
C117	0.6798 (2)	0.6980 (3)	0.2380 (2)	0.0318 (7)
H11A	0.6667	0.7385	0.1826	0.048*
H11B	0.6414	0.7247	0.2682	0.048*
H11C	0.6734	0.6132	0.2283	0.048*
C118	0.92484 (19)	0.6332 (3)	0.5082 (2)	0.0257 (7)
H11D	0.9859	0.6413	0.5269	0.039*
H11E	0.9099	0.5499	0.5021	0.039*
H11F	0.9031	0.6689	0.551	0.039*
C311	0.82259 (17)	0.1691 (2)	0.27715 (17)	0.0149 (5)
C312	0.90839 (17)	0.1910 (2)	0.30147 (18)	0.0168 (5)
H312	0.9352	0.2308	0.354	0.02*
C313	0.95533 (17)	0.1553 (2)	0.24967 (19)	0.0193 (6)
H313	1.0138	0.1707	0.267	0.023*
C314	0.91660 (18)	0.0976 (2)	0.17315 (19)	0.0212 (6)
H314	0.9483	0.0736	0.1375	0.025*
C315	0.83113 (18)	0.0749 (2)	0.14845 (19)	0.0190 (6)
H315	0.8045	0.035	0.0959	0.023*
C316	0.78449 (17)	0.1103 (2)	0.20019 (18)	0.0168 (5)
H316	0.7261	0.0943	0.1829	0.02*
C321	0.66208 (17)	0.1553 (2)	0.30122 (17)	0.0153 (5)
C322	0.59944 (17)	0.2176 (2)	0.24164 (19)	0.0202 (6)
H322	0.6093	0.296	0.2279	0.024*
C323	0.52197 (18)	0.1657 (3)	0.2017 (2)	0.0243 (6)
H323	0.4799	0.208	0.1594	0.029*
C324	0.50573 (19)	0.0534 (3)	0.2231 (2)	0.0260 (7)
H324	0.4523	0.019	0.1966	0.031*
C325	0.5676 (2)	-0.0085 (3)	0.2829 (2)	0.0386 (9)
H325	0.557	-0.086	0.2979	0.046*
C326	0.6459 (2)	0.0424 (3)	0.3217 (2)	0.0335 (8)
H326	0.6884	-0.001	0.3627	0.04*
C331	0.81170 (17)	0.1399 (2)	0.44688 (17)	0.0156 (5)
C332	0.7927 (2)	0.1752 (3)	0.51929 (19)	0.0266 (7)
H332	0.7579	0.2416	0.5163	0.032*
C333	0.8240 (2)	0.1144 (3)	0.5959 (2)	0.0334 (8)
H333	0.8091	0.1378	0.6447	0.04*
C334	0.8770 (2)	0.0196 (3)	0.60188 (19)	0.0263 (7)
H334	0.8999	-0.0205	0.655	0.032*
C335	0.89635 (18)	-0.0163 (2)	0.53013 (19)	0.0222 (6)
H335	0.932	-0.082	0.5339	0.027*
C336	0.86407 (18)	0.0429 (2)	0.45242 (19)	0.0196 (6)
H336	0.8775	0.0176	0.4032	0.024*
N11	0.75420 (14)	0.59212 (18)	0.40401 (15)	0.0148 (5)
O12	0.66821 (11)	0.37910 (16)	0.42069 (13)	0.0174 (4)
O14	0.90058 (13)	0.46375 (18)	0.30306 (13)	0.0225 (4)
P13	0.76530 (4)	0.22182 (6)	0.34722 (4)	0.01315 (15)

Rh1	0.761010 (12)	0.417429 (17)	0.371227 (13)	0.01263 (8)	
O01	0.4759 (6)	0.3631 (7)	0.0331 (6)	0.0963 (16)	0.5
C01	0.4182 (4)	0.5252 (5)	-0.0389 (4)	0.0963 (16)	
H01A	0.366	0.4813	-0.0482	0.144*	
H01B	0.4236	0.5472	-0.0946	0.144*	
H01C	0.4173	0.5961	-0.0055	0.144*	
C02	0.4899 (9)	0.4513 (10)	0.0086 (9)	0.0963 (16)	0.5

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0290 (16)	0.0158 (14)	0.0378 (18)	0.0010 (12)	0.0183 (14)	-0.0024 (13)
C2	0.0145 (13)	0.0149 (13)	0.0204 (14)	0.0019 (10)	0.0049 (11)	0.0014 (10)
C3	0.0165 (13)	0.0185 (13)	0.0215 (14)	0.0041 (11)	0.0082 (11)	0.0009 (11)
C4	0.0127 (13)	0.0192 (13)	0.0220 (14)	0.0039 (10)	0.0056 (11)	0.0037 (11)
C5	0.0233 (15)	0.0239 (15)	0.0378 (18)	-0.0002 (12)	0.0176 (14)	0.0045 (13)
C14	0.0190 (14)	0.0134 (12)	0.0157 (13)	-0.0017 (10)	0.0032 (11)	-0.0036 (10)
C111	0.0206 (14)	0.0097 (12)	0.0224 (14)	-0.0009 (10)	0.0118 (12)	-0.0016 (10)
C112	0.0273 (15)	0.0152 (13)	0.0279 (16)	0.0027 (11)	0.0137 (13)	0.0022 (11)
C113	0.0408 (19)	0.0149 (13)	0.0312 (17)	0.0020 (13)	0.0193 (15)	0.0041 (12)
C114	0.0410 (19)	0.0138 (14)	0.043 (2)	-0.0063 (13)	0.0286 (16)	-0.0043 (13)
C115	0.0243 (15)	0.0173 (14)	0.0375 (18)	-0.0055 (11)	0.0189 (14)	-0.0117 (12)
C116	0.0220 (14)	0.0136 (12)	0.0242 (15)	-0.0014 (11)	0.0099 (12)	-0.0038 (11)
C117	0.0293 (17)	0.0335 (17)	0.0289 (17)	0.0046 (14)	0.0043 (14)	0.0104 (14)
C118	0.0217 (15)	0.0267 (16)	0.0243 (16)	-0.0035 (12)	0.0013 (13)	-0.0047 (12)
C311	0.0186 (13)	0.0098 (12)	0.0171 (13)	0.0012 (10)	0.0066 (11)	0.0014 (10)
C312	0.0165 (13)	0.0147 (12)	0.0180 (13)	-0.0017 (10)	0.0037 (11)	-0.0003 (10)
C313	0.0138 (13)	0.0189 (13)	0.0249 (15)	0.0030 (11)	0.0059 (11)	0.0037 (11)
C314	0.0225 (15)	0.0196 (14)	0.0242 (15)	0.0043 (11)	0.0112 (13)	0.0025 (11)
C315	0.0211 (14)	0.0176 (13)	0.0180 (14)	-0.0002 (11)	0.0060 (12)	-0.0023 (10)
C316	0.0165 (13)	0.0143 (12)	0.0187 (14)	-0.0012 (10)	0.0041 (11)	-0.0001 (10)
C321	0.0159 (13)	0.0149 (12)	0.0163 (13)	-0.0011 (10)	0.0067 (11)	-0.0025 (10)
C322	0.0173 (14)	0.0168 (13)	0.0251 (15)	-0.0012 (11)	0.0049 (12)	-0.0008 (11)
C323	0.0155 (14)	0.0265 (16)	0.0261 (16)	0.0030 (12)	0.0001 (12)	0.0001 (12)
C324	0.0166 (14)	0.0312 (16)	0.0283 (16)	-0.0067 (12)	0.0045 (13)	-0.0052 (13)
C325	0.0317 (18)	0.0261 (17)	0.048 (2)	-0.0167 (14)	-0.0006 (16)	0.0098 (15)
C326	0.0249 (16)	0.0237 (16)	0.041 (2)	-0.0047 (13)	-0.0047 (15)	0.0129 (14)
C331	0.0162 (13)	0.0134 (12)	0.0147 (13)	-0.0016 (10)	0.0014 (11)	-0.0017 (10)
C332	0.0368 (18)	0.0224 (15)	0.0195 (15)	0.0132 (13)	0.0077 (13)	0.0011 (12)
C333	0.054 (2)	0.0313 (17)	0.0153 (15)	0.0234 (16)	0.0113 (15)	0.0034 (13)
C334	0.0340 (17)	0.0214 (15)	0.0188 (15)	0.0074 (13)	0.0022 (13)	0.0031 (12)
C335	0.0246 (15)	0.0168 (14)	0.0233 (15)	0.0049 (11)	0.0052 (12)	0.0005 (11)
C336	0.0220 (14)	0.0160 (13)	0.0213 (14)	0.0013 (11)	0.0075 (12)	-0.0017 (11)
N11	0.0142 (11)	0.0107 (10)	0.0191 (12)	-0.0009 (8)	0.0049 (9)	0.0012 (8)
O12	0.0152 (9)	0.0148 (9)	0.0242 (10)	-0.0008 (8)	0.0091 (8)	0.0014 (8)
O14	0.0261 (11)	0.0180 (10)	0.0280 (11)	-0.0059 (8)	0.0152 (9)	-0.0034 (8)
P13	0.0123 (3)	0.0117 (3)	0.0142 (3)	-0.0008 (2)	0.0027 (3)	-0.0002 (2)
Rh1	0.01250 (12)	0.01060 (11)	0.01490 (12)	-0.00047 (7)	0.00457 (8)	0.00021 (7)

supplementary materials

O01	0.139 (5)	0.057 (3)	0.104 (4)	-0.010 (3)	0.054 (4)	0.007 (2)
C01	0.139 (5)	0.057 (3)	0.104 (4)	-0.010 (3)	0.054 (4)	0.007 (2)
C02	0.139 (5)	0.057 (3)	0.104 (4)	-0.010 (3)	0.054 (4)	0.007 (2)

Geometric parameters (Å, °)

C1—C2	1.513 (4)	C313—H313	0.95
C1—H1A	0.98	C314—C315	1.390 (4)
C1—H1B	0.98	C314—H314	0.95
C1—H1C	0.98	C315—C316	1.389 (4)
C2—N11	1.320 (4)	C315—H315	0.95
C2—C3	1.420 (4)	C316—H316	0.95
C3—C4	1.384 (4)	C321—C326	1.380 (4)
C3—H3	0.95	C321—C322	1.383 (4)
C4—O12	1.289 (3)	C321—P13	1.824 (3)
C4—C5	1.507 (4)	C322—C323	1.391 (4)
C5—H5A	0.98	C322—H322	0.95
C5—H5B	0.98	C323—C324	1.378 (4)
C5—H5C	0.98	C323—H323	0.95
C14—O14	1.154 (3)	C324—C325	1.375 (4)
C14—Rh1	1.805 (3)	C324—H324	0.95
C111—C116	1.405 (4)	C325—C326	1.394 (4)
C111—C112	1.406 (4)	C325—H325	0.95
C111—N11	1.439 (3)	C326—H326	0.95
C112—C113	1.387 (4)	C331—C332	1.385 (4)
C112—C117	1.507 (4)	C331—C336	1.400 (4)
C113—C114	1.388 (5)	C331—P13	1.827 (3)
C113—H113	0.95	C332—C333	1.383 (4)
C114—C115	1.385 (5)	C332—H332	0.95
C114—H114	0.95	C333—C334	1.386 (4)
C115—C116	1.396 (4)	C333—H333	0.95
C115—H115	0.95	C334—C335	1.380 (4)
C116—C118	1.503 (4)	C334—H334	0.95
C117—H11A	0.98	C335—C336	1.390 (4)
C117—H11B	0.98	C335—H335	0.95
C117—H11C	0.98	C336—H336	0.95
C118—H11D	0.98	N11—Rh1	2.076 (2)
C118—H11E	0.98	O12—Rh1	2.0277 (19)
C118—H11F	0.98	P13—Rh1	2.2701 (7)
C311—C316	1.391 (4)	O01—C02	1.135 (12)
C311—C312	1.394 (4)	C01—C02	1.474 (13)
C311—P13	1.825 (3)	C01—H01A	0.98
C312—C313	1.395 (4)	C01—H01B	0.98
C312—H312	0.95	C01—H01C	0.98
C313—C314	1.382 (4)	C02—C01 ⁱ	1.492 (16)
C2—C1—H1A	109.5	C314—C315—H315	119.8
C2—C1—H1B	109.5	C315—C316—C311	120.5 (3)
H1A—C1—H1B	109.5	C315—C316—H316	119.7
C2—C1—H1C	109.5	C311—C316—H316	119.7

H1A—C1—H1C	109.5	C326—C321—C322	119.0 (3)
H1B—C1—H1C	109.5	C326—C321—P13	121.6 (2)
N11—C2—C3	123.8 (2)	C322—C321—P13	119.3 (2)
N11—C2—C1	120.5 (2)	C321—C322—C323	120.2 (3)
C3—C2—C1	115.8 (2)	C321—C322—H322	119.9
C4—C3—C2	127.1 (3)	C323—C322—H322	119.9
C4—C3—H3	116.5	C324—C323—C322	120.6 (3)
C2—C3—H3	116.5	C324—C323—H323	119.7
O12—C4—C3	125.6 (3)	C322—C323—H323	119.7
O12—C4—C5	113.9 (2)	C325—C324—C323	119.4 (3)
C3—C4—C5	120.4 (3)	C325—C324—H324	120.3
C4—C5—H5A	109.5	C323—C324—H324	120.3
C4—C5—H5B	109.5	C324—C325—C326	120.1 (3)
H5A—C5—H5B	109.5	C324—C325—H325	119.9
C4—C5—H5C	109.5	C326—C325—H325	119.9
H5A—C5—H5C	109.5	C321—C326—C325	120.6 (3)
H5B—C5—H5C	109.5	C321—C326—H326	119.7
O14—C14—Rh1	177.7 (2)	C325—C326—H326	119.7
C116—C111—C112	121.1 (3)	C332—C331—C336	119.2 (3)
C116—C111—N11	119.5 (2)	C332—C331—P13	117.9 (2)
C112—C111—N11	119.3 (2)	C336—C331—P13	122.8 (2)
C113—C112—C111	118.4 (3)	C333—C332—C331	120.5 (3)
C113—C112—C117	121.2 (3)	C333—C332—H332	119.7
C111—C112—C117	120.4 (3)	C331—C332—H332	119.7
C112—C113—C114	121.3 (3)	C332—C333—C334	120.3 (3)
C112—C113—H113	119.3	C332—C333—H333	119.9
C114—C113—H113	119.3	C334—C333—H333	119.9
C115—C114—C113	119.6 (3)	C335—C334—C333	119.6 (3)
C115—C114—H114	120.2	C335—C334—H334	120.2
C113—C114—H114	120.2	C333—C334—H334	120.2
C114—C115—C116	121.2 (3)	C334—C335—C336	120.5 (3)
C114—C115—H115	119.4	C334—C335—H335	119.7
C116—C115—H115	119.4	C336—C335—H335	119.7
C115—C116—C111	118.3 (3)	C335—C336—C331	119.8 (3)
C115—C116—C118	121.3 (3)	C335—C336—H336	120.1
C111—C116—C118	120.4 (3)	C331—C336—H336	120.1
C112—C117—H11A	109.5	C2—N11—C111	118.0 (2)
C112—C117—H11B	109.5	C2—N11—Rh1	125.79 (18)
H11A—C117—H11B	109.5	C111—N11—Rh1	116.24 (17)
C112—C117—H11C	109.5	C4—O12—Rh1	127.13 (18)
H11A—C117—H11C	109.5	C311—P13—C321	103.27 (12)
H11B—C117—H11C	109.5	C311—P13—C331	103.55 (12)
C116—C118—H11D	109.5	C321—P13—C331	103.59 (12)
C116—C118—H11E	109.5	C311—P13—Rh1	119.08 (9)
H11D—C118—H11E	109.5	C321—P13—Rh1	113.64 (9)
C116—C118—H11F	109.5	C331—P13—Rh1	112.06 (9)
H11D—C118—H11F	109.5	C14—Rh1—O12	177.36 (10)
H11E—C118—H11F	109.5	C14—Rh1—N11	93.02 (11)
C316—C311—C312	118.7 (3)	O12—Rh1—N11	89.38 (8)

supplementary materials

C316—C311—P13	123.2 (2)	C14—Rh1—P13	91.59 (9)
C312—C311—P13	118.1 (2)	O12—Rh1—P13	85.95 (6)
C311—C312—C313	120.8 (3)	N11—Rh1—P13	174.36 (7)
C311—C312—H312	119.6	C02—C01—H01A	109.5
C313—C312—H312	119.6	C02—C01—H01B	109.5
C314—C313—C312	119.8 (3)	H01A—C01—H01B	109.5
C314—C313—H313	120.1	C02—C01—H01C	109.5
C312—C313—H313	120.1	H01A—C01—H01C	109.5
C313—C314—C315	119.8 (3)	H01B—C01—H01C	109.5
C313—C314—H314	120.1	O01—C02—C01	117.7 (13)
C315—C314—H314	120.1	O01—C02—C01 ⁱ	110.6 (12)
C316—C315—C314	120.3 (3)	C01—C02—C01 ⁱ	131.4 (9)
C316—C315—H315	119.8		
N11—C2—C3—C4	-4.8 (5)	P13—C331—C336—C335	-179.2 (2)
C1—C2—C3—C4	175.4 (3)	C3—C2—N11—C111	178.5 (3)
C2—C3—C4—O12	1.8 (5)	C1—C2—N11—C111	-1.7 (4)
C2—C3—C4—C5	-177.6 (3)	C3—C2—N11—Rh1	-3.0 (4)
C116—C111—C112—C113	-2.1 (4)	C1—C2—N11—Rh1	176.8 (2)
N11—C111—C112—C113	-177.7 (3)	C116—C111—N11—C2	91.5 (3)
C116—C111—C112—C117	177.3 (3)	C112—C111—N11—C2	-92.9 (3)
N11—C111—C112—C117	1.7 (4)	C116—C111—N11—Rh1	-87.2 (3)
C111—C112—C113—C114	1.2 (4)	C112—C111—N11—Rh1	88.5 (3)
C117—C112—C113—C114	-178.3 (3)	C3—C4—O12—Rh1	9.0 (4)
C112—C113—C114—C115	0.4 (5)	C5—C4—O12—Rh1	-171.64 (18)
C113—C114—C115—C116	-1.1 (4)	C316—C311—P13—C321	9.9 (3)
C114—C115—C116—C111	0.1 (4)	C312—C311—P13—C321	-171.4 (2)
C114—C115—C116—C118	178.5 (3)	C316—C311—P13—C331	117.7 (2)
C112—C111—C116—C115	1.5 (4)	C312—C311—P13—C331	-63.6 (2)
N11—C111—C116—C115	177.1 (2)	C316—C311—P13—Rh1	-117.1 (2)
C112—C111—C116—C118	-176.9 (3)	C312—C311—P13—Rh1	61.6 (2)
N11—C111—C116—C118	-1.3 (4)	C326—C321—P13—C311	83.9 (3)
C316—C311—C312—C313	0.3 (4)	C322—C321—P13—C311	-93.4 (2)
P13—C311—C312—C313	-178.5 (2)	C326—C321—P13—C331	-23.8 (3)
C311—C312—C313—C314	0.1 (4)	C322—C321—P13—C331	158.9 (2)
C312—C313—C314—C315	-0.3 (4)	C326—C321—P13—Rh1	-145.7 (2)
C313—C314—C315—C316	0.2 (4)	C322—C321—P13—Rh1	37.0 (2)
C314—C315—C316—C311	0.1 (4)	C332—C331—P13—C311	168.3 (2)
C312—C311—C316—C315	-0.4 (4)	C336—C331—P13—C311	-12.7 (3)
P13—C311—C316—C315	178.3 (2)	C332—C331—P13—C321	-84.2 (2)
C326—C321—C322—C323	-1.5 (4)	C336—C331—P13—C321	94.8 (3)
P13—C321—C322—C323	175.9 (2)	C332—C331—P13—Rh1	38.7 (3)
C321—C322—C323—C324	2.2 (5)	C336—C331—P13—Rh1	-142.3 (2)
C322—C323—C324—C325	-1.5 (5)	C4—O12—Rh1—N11	-11.8 (2)
C323—C324—C325—C326	0.2 (6)	C4—O12—Rh1—P13	165.0 (2)
C322—C321—C326—C325	0.2 (5)	C2—N11—Rh1—C14	-170.1 (2)
P13—C321—C326—C325	-177.2 (3)	C111—N11—Rh1—C14	8.5 (2)
C324—C325—C326—C321	0.5 (6)	C2—N11—Rh1—O12	8.8 (2)
C336—C331—C332—C333	-0.9 (5)	C111—N11—Rh1—O12	-172.62 (19)

P13—C331—C332—C333	178.1 (3)	C311—P13—Rh1—C14	-14.40 (13)
C331—C332—C333—C334	2.0 (5)	C321—P13—Rh1—C14	-136.42 (13)
C332—C333—C334—C335	-2.0 (5)	C331—P13—Rh1—C14	106.57 (13)
C333—C334—C335—C336	0.9 (5)	C311—P13—Rh1—O12	166.55 (12)
C334—C335—C336—C331	0.2 (4)	C321—P13—Rh1—O12	44.53 (11)
C332—C331—C336—C335	-0.2 (4)	C331—P13—Rh1—O12	-72.48 (11)

Symmetry codes: (i) $-x+1, -y+1, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C332—H332 \cdots O12	0.95	2.38	3.201 (3)	144
C334—H334 \cdots O14 ⁱⁱ	0.95	2.51	3.201 (4)	130
C1—H1B \cdots O01 ⁱⁱⁱ	0.98	2.54	3.372 (9)	142

Symmetry codes: (ii) $x, -y+1/2, z+1/2$; (iii) $-x+1, y+1/2, -z+1/2$.

Table 2

Comparative geometrical parameters (\AA , $^\circ$) *for similar* $[Rh(N,O\text{-bid})(CO)(PPh_3)]$ *complexes*

Parameters	(I) ^a	(II) ^b	(III) ^c	(IV) ^d
Rh1—N11	2.077 (2)	2.069 (2)	2.045 (4)	2.045 (3)
Rh1—O12	2.027 (2)	2.028 (2)	2.044 (3)	2.045 (2)
Rh1—P13	2.2704 (7)	2.2635 (6)	2.275 (1)	2.281 (2)
Rh1—C14	1.812 (3)	1.807 (2)	1.784 (5)	1.804 (3)
C14—O14	1.147 (3)	1.152 (3)	1.142 (7)	1.148 (4)
N11 \cdots O12	2.885 (3)	2.885 (3)	2.826 (6)	2.841 (3)
N11—Rh1—O12	89.31 (9)	89.54 (8)	87.4 (1)	87.95 (8)
O12—Rh1—P13	85.95 (6)	84.97 (5)	89.7 (1)	89.91 (5)
P13—Rh1—C14	91.57 (9)	91.87 (7)	90.3 (2)	89.48 (9)
N11—Rh1—C14	93.1 (1)	93.6 (1)	92.6 (2)	92.6 (1)
N11—C2—C4—O12	-2.6 (2)	4.1 (2)	1.2 (4)	1.5 (2)
θ_E^e	155.77 (2)	156.39 (3)	156.0 (2)	156.23 (4)

Notes: (a) This work; (b) *N,O*-bid = 4-(2,3-dimethyl phenylamino)pent-3-en-2-onato (Venter *et al.*, 2009); (c) *N,O*-bid = 4-amino-pent-3-en-2-onato (Damoense *et al.*, 1994); (d) *N,O*-bid = 4-amino-1,1,1-trifluoro-pent-3-en-2-onato (Varshavsky *et al.*, 2001); (e) Tolman (1977).

Fig. 1




Fig. 2

