

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(3-Chlorobenzoyloxy)urea

 Xi Mai,^{a,b} Hong-Ying Xia,^c Yu-Sheng Cao,^{a,d*} Wei Tong^b
and Guo-Gang Tu^b

^aState Key Laboratory of Food Science and Technology, Nanchang University, Nanchang 330047, People's Republic of China, ^bDepartment of Pharmacy, Medical College of Nanchang University, Nanchang 330006, People's Republic of China, ^cDepartment of Pharmacy, Shangrao Branch of Jiangxi Medical College, Shangrao 334000, People's Republic of China, and ^dSino-German Joint Research Institute of Nanchang University, Nanchang 330006, People's Republic of China
Correspondence e-mail: cmxlf2008@163.com


Received 29 October 2009; accepted 29 October 2009

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.047; wR factor = 0.094; data-to-parameter ratio = 13.9.

The asymmetric unit of the crystal structure of the title compound, $\text{C}_8\text{H}_9\text{ClN}_2\text{O}_2$, contains four independent molecules. The dihedral angles between the urea $\text{N}-(\text{C}=\text{O})-\text{N}$ planes and the benzene rings are 83.3 (3), 87.8 (1), 89.1 (1) and 17.5 (2)° in the four molecules. Extensive $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonding is present in the crystal structure.

Related literature

For general background to the design and synthesis of hydroxyurea derivatives and their *in vitro* antitumor activity, see: Mai *et al.* (2009). For related structures, see: Armagan *et al.* (1976); Nielsen *et al.* (1993); Berman & Kim (1967); Howard *et al.* (1967); Larsen & Jerslev (1966); Thiessen *et al.* (1978); Yoshitaka *et al.* (1993).



Experimental

Crystal data

$\text{C}_8\text{H}_9\text{ClN}_2\text{O}_2$
 $M_r = 200.62$
 Triclinic, $P\bar{1}$
 $a = 10.830$ (1) Å
 $b = 13.9410$ (14) Å
 $c = 14.2750$ (15) Å
 $\alpha = 69.672$ (1)°
 $\beta = 75.828$ (2)°
 $\gamma = 70.388$ (1)°
 $V = 1883.6$ (3) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.37$ mm⁻¹
 $T = 298$ K
 $0.43 \times 0.40 \times 0.05$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.856$, $T_{\max} = 0.982$
 9908 measured reflections
 6533 independent reflections
 3124 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.094$
 $S = 1.01$
 6533 reflections
 469 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O5}^i$	0.90	2.20	3.096 (3)	173
$\text{N2}-\text{H2A}\cdots\text{O1}^i$	0.86	2.16	3.023 (3)	177
$\text{N2}-\text{H2B}\cdots\text{O3}^{ii}$	0.86	2.29	2.971 (3)	136
$\text{N4}-\text{H4A}\cdots\text{O7}^{iii}$	0.86	2.11	2.971 (3)	176
$\text{N4}-\text{H4B}\cdots\text{O5}$	0.86	2.39	3.017 (3)	130
$\text{N5}-\text{H5}\cdots\text{O1}^i$	0.90	2.19	3.090 (3)	176
$\text{N6}-\text{H6A}\cdots\text{O5}^{iv}$	0.86	2.07	2.925 (3)	177
$\text{N7}-\text{H7}\cdots\text{O7}^v$	0.90	2.04	2.937 (3)	171
$\text{N8}-\text{H8A}\cdots\text{O3}^{ii}$	0.86	2.09	2.947 (3)	177
$\text{N8}-\text{H8B}\cdots\text{O1}^i$	0.86	2.25	2.976 (3)	142

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x+1, y, z$; (iii) $x-1, y, z$; (iv) $-x, -y+2, -z+1$; (v) $-x+1, -y+2, -z+1$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: SHELXL97.

The authors gratefully acknowledge the financial support of this study by the National Key S&T Special Project of China: Grand New Drug R&D (NO. 2009ZX09103-087) and the program of the Nanchang Department of Science and Technology, China (No. 2008368).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2661).

References

- Armagan, N., Richards, J. P. G. & Uraz, A. A. (1976). *Acta Cryst.* **B32**, 1042–1047.
 Berman, H. & Kim, S. H. (1967). *Acta Cryst.* **23**, 180–181.
 Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Howard, W., Shields, P. J., Hamrick, J. & Welby, R. (1967). *J. Chem. Phys.* **46**, 2510–2514.
 Larsen, I. K. & Jerslev, B. (1966). *Acta Chem. Scand.* **20**, 983–991.
 Mai, X., Xia, H.-Y., Cao, Y.-S., Lu, X.-S. & Fang, X.-N. (2009). *Acta Cryst.* **E65**, o442.
 Nielsen, B. B., Frydenvang, K. & Larsen, I. K. (1993). *Acta Cryst.* **C49**, 1018–1022.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Thiessen, W. E., Levy, H. A. & Flaig, B. D. (1978). *Acta Cryst.* **B34**, 2495–2502.
 Yoshitaka, S., James, L. S., Alan, J. H., Adam, H. L., Timothy, J. K., Warren, H. L. D., Hope, W. & Earl, F. K. (1993). *J. Med. Chem.* **36**, 3580–3594.

supplementary materials

Acta Cryst. (2009). E65, o2983 [doi:10.1107/S160053680904553X]

1-(3-Chlorobenzoyloxy)urea

X. Mai, H.-Y. Xia, Y.-S. Cao, W. Tong and G.-G. Tu

Comment

Hydroxyurea (HU) is a substance used in cancer chemotherapy for many years, but it has several disadvantages, such as short half-life, extremely polar nature, the rapid development of resistance and so on. To obtain more potent compound, we have designed and synthesized HU derivatives, and evaluated their *in vitro* antitumor activities in our previous work (Mai *et al.*, 2009). Here we report the crystal structure of the title compound, 3-chlorobenzoyloxyurea.

The structure of 3-chlorobenzoyloxyurea is shown in Fig. 1. The conformations of the N–O and C=O bonds are opposite to each other, similar to that observed in N-hydroxyurea (Howard *et al.*, 1967; Thiessen *et al.*, 1978; Armagan *et al.*, 1976; Ber-man *et al.*, 1967; Larsen & Jerslev, 1966), 1-hydroxy-1-methylurea, 1-hydroxy-3-methylurea (Nielsen *et al.*, 1993), N-(6-phenoxy-2H-chromen-3-ylmethyl)-N-hydroxyurea (Yoshitaka *et al.*, 1993) and 1-(2-fluorobenzyl)-1-(2-fluorobenzoyloxy)urea (Mai *et al.*, 2009). The bond parameters are similar to 1-(2-fluorobenzyl)-1-(2-fluorobenzoyloxy)urea (Mai *et al.*, 2009). The asymmetric unit of the title compound contains four independent molecules. The dihedral angles between the urea N–(C=O)–N planes and benzene ring are 83.3 (3)°, 87.8 (1)°, 89.1 (1)° and 17.5 (2)° for the four molecules. The N–O bonds are twisted out of the urea N–(C=O)–N planes by 18.4 (3)°, 17.9 (3)°, 19.2 (4)° and -17.8 (3)°, respectively in the four molecules. In the crystal structure, molecules are linked through intermolecular N–H...O hydrogen bonds, forming the zigzag chain.

Experimental

The title compound was synthesized by hydroxyurea (0.026 mol) with 3-chlorobenzyl chloride (0.034 mol) in methanol (80 ml) in the presence of potassium hydroxide (0.034 mol). After refluxing for 13 h, solvent was removed under reduced pressure at 308 K. The resulting crude solid was filtered and washed in trichloromethane, then recrystallized in acetone and trichloromethane solution (5:2), filtered and dried. Colorless platelet single crystals of the title compound were recrystallized from the mixed solvent acetone and n-hexane (5:10).

Refinement

H atoms were placed in calculated positions with N–H = 0.90 (imino), 0.86 Å (amino), C–H = 0.93 (aromatic) and 0.97 Å (methylene), and refined in riding mode with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

Figures




Fig. 1. Molecular structure of the title compound showing the atom labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. The unit cell diagram showing intermolecular hydrogen bonding as dashed lines

1-(3-Chlorobenzoyloxy)urea

Crystal data

$C_8H_9ClN_2O_2$

$M_r = 200.62$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 10.8300$ (10) Å

$b = 13.9410$ (14) Å

$c = 14.2750$ (15) Å

$\alpha = 69.6720$ (10)°

$\beta = 75.828$ (2)°

$\gamma = 70.3880$ (10)°

$V = 1883.6$ (3) Å³

$Z = 8$

$F_{000} = 832$

$D_x = 1.415$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1978 reflections

$\theta = 2.2$ – 22.6 °

$\mu = 0.37$ mm⁻¹

$T = 298$ K

Platelet, colourless

$0.43 \times 0.40 \times 0.05$ mm

Data collection

Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298$ K

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.856$, $T_{\max} = 0.982$

9908 measured reflections

6533 independent reflections

3124 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 25.0$ °

$\theta_{\min} = 1.5$ °

$h = -12 \rightarrow 11$

$k = -16 \rightarrow 16$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.047$	H-atom parameters constrained
$wR(F^2) = 0.094$	$w = 1/[\sigma^2(F_o^2) + (0.0205P)^2]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
6533 reflections	$(\Delta/\sigma)_{\max} = 0.001$
469 parameters	$\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	1.11832 (10)	0.31531 (8)	-0.02808 (6)	0.0839 (3)
C12	0.25832 (11)	0.58558 (9)	-0.04798 (7)	0.0998 (4)
C13	0.39222 (11)	0.83726 (9)	-0.04386 (7)	0.1020 (4)
C14	0.16163 (12)	1.13164 (10)	0.84040 (10)	0.1271 (5)
N1	0.8146 (2)	0.36994 (18)	0.39732 (16)	0.0401 (6)
H1	0.8418	0.3029	0.4370	0.048*
N2	0.6676 (2)	0.51825 (17)	0.44071 (16)	0.0419 (6)
H2A	0.5900	0.5522	0.4631	0.050*
H2B	0.7301	0.5491	0.4177	0.050*
N3	0.0506 (2)	0.61462 (18)	0.39981 (16)	0.0380 (6)
H3	0.0757	0.5489	0.4424	0.046*
N4	-0.0889 (2)	0.76897 (17)	0.43587 (16)	0.0453 (7)
H4A	-0.1647	0.8054	0.4583	0.054*
H4B	-0.0251	0.7980	0.4089	0.054*
N5	0.2948 (2)	0.86095 (18)	0.39097 (17)	0.0438 (6)
H5	0.3203	0.7948	0.4327	0.053*
N6	0.1619 (2)	1.01583 (18)	0.42966 (17)	0.0529 (7)
H6A	0.0882	1.0524	0.4549	0.063*
H6B	0.2252	1.0448	0.3991	0.063*
N7	0.4342 (2)	0.89042 (18)	0.59283 (16)	0.0422 (6)
H7	0.4119	0.9574	0.5520	0.051*
N8	0.5745 (2)	0.73892 (17)	0.55203 (16)	0.0447 (6)
H8A	0.6513	0.7025	0.5309	0.054*
H8B	0.5093	0.7111	0.5735	0.054*

supplementary materials

O1	0.60499 (19)	0.36862 (15)	0.47386 (15)	0.0501 (6)
O2	0.91190 (18)	0.42284 (15)	0.38500 (13)	0.0421 (5)
O3	-0.15777 (19)	0.62023 (14)	0.48081 (14)	0.0462 (5)
O4	0.15349 (18)	0.66423 (15)	0.38002 (14)	0.0440 (5)
O5	0.09142 (19)	0.86693 (15)	0.48084 (15)	0.0524 (6)
O6	0.4015 (2)	0.90763 (15)	0.36655 (15)	0.0510 (6)
O7	0.64654 (19)	0.88364 (14)	0.52150 (14)	0.0469 (5)
O8	0.33178 (18)	0.84381 (15)	0.60348 (14)	0.0458 (5)
C1	0.6909 (3)	0.4182 (2)	0.4417 (2)	0.0379 (7)
C2	0.9543 (3)	0.4670 (2)	0.2794 (2)	0.0463 (8)
H2C	0.9940	0.5227	0.2714	0.056*
H2D	0.8774	0.4992	0.2444	0.056*
C3	1.0527 (3)	0.3856 (2)	0.2309 (2)	0.0400 (8)
C4	1.0418 (3)	0.3863 (2)	0.1359 (2)	0.0473 (8)
H4	0.9715	0.4348	0.1032	0.057*
C5	1.1353 (3)	0.3151 (3)	0.0903 (2)	0.0489 (8)
C6	1.2408 (3)	0.2433 (3)	0.1353 (2)	0.0573 (9)
H6	1.3038	0.1964	0.1028	0.069*
C7	1.2522 (3)	0.2418 (3)	0.2306 (3)	0.0619 (10)
H7A	1.3230	0.1931	0.2627	0.074*
C8	1.1592 (3)	0.3119 (3)	0.2776 (2)	0.0551 (9)
H8	1.1675	0.3100	0.3416	0.066*
C9	-0.0704 (3)	0.6681 (2)	0.4428 (2)	0.0373 (7)
C10	0.1852 (3)	0.7051 (2)	0.2732 (2)	0.0506 (9)
H10A	0.2443	0.7490	0.2591	0.061*
H10B	0.1045	0.7503	0.2456	0.061*
C11	0.2494 (3)	0.6202 (3)	0.2204 (2)	0.0462 (8)
C12	0.2222 (3)	0.6366 (3)	0.1247 (2)	0.0552 (9)
H12	0.1590	0.6975	0.0957	0.066*
C13	0.2892 (4)	0.5626 (3)	0.0729 (2)	0.0578 (9)
C14	0.3784 (4)	0.4728 (3)	0.1150 (3)	0.0694 (11)
H14	0.4233	0.4234	0.0795	0.083*
C15	0.4031 (4)	0.4544 (3)	0.2107 (3)	0.0804 (12)
H15	0.4632	0.3915	0.2403	0.096*
C16	0.3399 (3)	0.5278 (3)	0.2632 (2)	0.0659 (10)
H16	0.3582	0.5151	0.3275	0.079*
C17	0.1778 (3)	0.9153 (3)	0.4379 (2)	0.0424 (8)
C18	0.4308 (3)	0.9459 (3)	0.2589 (2)	0.0649 (10)
H18A	0.5039	0.9773	0.2420	0.078*
H18B	0.3544	1.0016	0.2333	0.078*
C19	0.4666 (3)	0.8607 (3)	0.2068 (2)	0.0498 (9)
C20	0.4215 (3)	0.8854 (3)	0.1150 (3)	0.0608 (10)
H20	0.3710	0.9537	0.0858	0.073*
C21	0.4533 (3)	0.8067 (3)	0.0688 (2)	0.0625 (10)
C22	0.5279 (3)	0.7067 (3)	0.1085 (3)	0.0665 (10)
H22	0.5489	0.6552	0.0754	0.080*
C23	0.5723 (3)	0.6827 (3)	0.1992 (3)	0.0687 (10)
H23	0.6225	0.6141	0.2278	0.082*
C24	0.5427 (3)	0.7592 (3)	0.2471 (2)	0.0575 (9)

H24	0.5744	0.7424	0.3074	0.069*
C25	0.5569 (3)	0.8378 (2)	0.5515 (2)	0.0364 (7)
C26	0.2837 (3)	0.8033 (2)	0.7081 (2)	0.0504 (9)
H26A	0.3582	0.7727	0.7456	0.061*
H26B	0.2461	0.7465	0.7157	0.061*
C27	0.1814 (3)	0.8842 (2)	0.7541 (2)	0.0443 (8)
C28	0.2123 (3)	0.9650 (3)	0.7696 (2)	0.0560 (9)
H28	0.2968	0.9742	0.7459	0.067*
C29	0.1197 (4)	1.0319 (3)	0.8198 (3)	0.0605 (10)
C30	-0.0046 (4)	1.0205 (3)	0.8561 (3)	0.0815 (12)
H30	-0.0658	1.0643	0.8926	0.098*
C31	-0.0373 (4)	0.9426 (3)	0.8373 (4)	0.1146 (18)
H31	-0.1230	0.9356	0.8587	0.137*
C32	0.0543 (4)	0.8753 (3)	0.7876 (3)	0.0877 (13)
H32	0.0304	0.8226	0.7763	0.105*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.1008 (8)	0.0997 (8)	0.0604 (6)	-0.0235 (7)	-0.0021 (6)	-0.0446 (6)
C12	0.0901 (8)	0.1568 (11)	0.0720 (7)	-0.0397 (8)	-0.0086 (6)	-0.0533 (7)
C13	0.1091 (9)	0.1371 (11)	0.0636 (6)	-0.0405 (8)	-0.0252 (6)	-0.0189 (6)
C14	0.1133 (10)	0.1276 (11)	0.1915 (13)	-0.0182 (8)	-0.0241 (10)	-0.1208 (10)
N1	0.0363 (16)	0.0381 (16)	0.0505 (15)	-0.0135 (14)	0.0008 (13)	-0.0199 (13)
N2	0.0341 (15)	0.0312 (16)	0.0625 (16)	-0.0105 (13)	0.0005 (13)	-0.0198 (13)
N3	0.0380 (16)	0.0337 (15)	0.0433 (15)	-0.0127 (13)	-0.0003 (13)	-0.0134 (12)
N4	0.0397 (16)	0.0333 (17)	0.0641 (17)	-0.0126 (13)	0.0039 (13)	-0.0209 (13)
N5	0.0432 (17)	0.0370 (17)	0.0543 (16)	-0.0154 (14)	-0.0014 (14)	-0.0168 (13)
N6	0.0454 (17)	0.0355 (17)	0.0766 (19)	-0.0135 (14)	0.0034 (15)	-0.0211 (14)
N7	0.0389 (16)	0.0341 (16)	0.0532 (16)	-0.0117 (14)	-0.0013 (14)	-0.0144 (13)
N8	0.0382 (16)	0.0316 (16)	0.0665 (17)	-0.0097 (13)	-0.0005 (13)	-0.0218 (13)
O1	0.0376 (13)	0.0378 (13)	0.0820 (16)	-0.0174 (11)	0.0014 (12)	-0.0260 (11)
O2	0.0374 (12)	0.0513 (14)	0.0452 (12)	-0.0174 (11)	0.0006 (10)	-0.0226 (10)
O3	0.0372 (13)	0.0399 (13)	0.0690 (14)	-0.0186 (11)	0.0037 (11)	-0.0250 (11)
O4	0.0387 (13)	0.0486 (14)	0.0505 (13)	-0.0196 (11)	0.0025 (11)	-0.0201 (11)
O5	0.0414 (14)	0.0408 (14)	0.0827 (16)	-0.0187 (12)	0.0033 (12)	-0.0283 (12)
O6	0.0466 (14)	0.0544 (15)	0.0578 (14)	-0.0230 (12)	0.0024 (11)	-0.0215 (11)
O7	0.0359 (13)	0.0333 (13)	0.0743 (14)	-0.0120 (11)	-0.0013 (11)	-0.0211 (11)
O8	0.0391 (13)	0.0521 (14)	0.0522 (13)	-0.0186 (11)	0.0001 (11)	-0.0212 (11)
C1	0.040 (2)	0.034 (2)	0.0448 (19)	-0.0074 (17)	-0.0077 (17)	-0.0187 (16)
C2	0.051 (2)	0.046 (2)	0.0417 (19)	-0.0195 (18)	0.0019 (17)	-0.0133 (16)
C3	0.042 (2)	0.039 (2)	0.0374 (18)	-0.0149 (17)	0.0032 (16)	-0.0115 (15)
C4	0.047 (2)	0.043 (2)	0.050 (2)	-0.0110 (17)	-0.0046 (17)	-0.0145 (17)
C5	0.053 (2)	0.051 (2)	0.045 (2)	-0.0189 (19)	0.0003 (18)	-0.0179 (17)
C6	0.061 (3)	0.047 (2)	0.057 (2)	-0.012 (2)	0.011 (2)	-0.0221 (18)
C7	0.056 (2)	0.055 (2)	0.058 (2)	0.0005 (19)	-0.004 (2)	-0.0146 (19)
C8	0.054 (2)	0.060 (3)	0.047 (2)	-0.012 (2)	-0.0039 (19)	-0.0160 (18)
C9	0.037 (2)	0.037 (2)	0.0447 (19)	-0.0107 (17)	-0.0061 (16)	-0.0193 (16)

supplementary materials

C10	0.051 (2)	0.042 (2)	0.050 (2)	-0.0157 (17)	0.0065 (18)	-0.0099 (17)
C11	0.045 (2)	0.044 (2)	0.043 (2)	-0.0166 (18)	0.0088 (17)	-0.0103 (17)
C12	0.046 (2)	0.053 (2)	0.059 (2)	-0.0114 (18)	0.0005 (19)	-0.0160 (19)
C13	0.055 (2)	0.064 (3)	0.050 (2)	-0.022 (2)	0.0023 (19)	-0.014 (2)
C14	0.086 (3)	0.058 (3)	0.056 (2)	-0.022 (2)	0.021 (2)	-0.025 (2)
C15	0.096 (3)	0.047 (3)	0.056 (3)	0.014 (2)	0.005 (2)	-0.008 (2)
C16	0.076 (3)	0.053 (3)	0.045 (2)	0.004 (2)	-0.002 (2)	-0.0105 (19)
C17	0.041 (2)	0.038 (2)	0.053 (2)	-0.0084 (18)	-0.0080 (18)	-0.0211 (17)
C18	0.075 (3)	0.052 (2)	0.062 (2)	-0.027 (2)	0.011 (2)	-0.013 (2)
C19	0.051 (2)	0.046 (2)	0.047 (2)	-0.0181 (19)	0.0096 (18)	-0.0128 (18)
C20	0.054 (2)	0.053 (2)	0.057 (2)	-0.0084 (19)	0.001 (2)	-0.006 (2)
C21	0.053 (2)	0.080 (3)	0.048 (2)	-0.021 (2)	0.0020 (19)	-0.015 (2)
C22	0.062 (3)	0.076 (3)	0.060 (2)	-0.014 (2)	0.006 (2)	-0.034 (2)
C23	0.066 (3)	0.060 (3)	0.063 (2)	0.004 (2)	-0.005 (2)	-0.020 (2)
C24	0.057 (2)	0.057 (3)	0.052 (2)	-0.009 (2)	-0.0074 (19)	-0.015 (2)
C25	0.036 (2)	0.030 (2)	0.0425 (18)	-0.0054 (17)	-0.0067 (16)	-0.0129 (15)
C26	0.054 (2)	0.041 (2)	0.054 (2)	-0.0180 (18)	0.0034 (18)	-0.0124 (17)
C27	0.039 (2)	0.043 (2)	0.0460 (19)	-0.0100 (17)	-0.0013 (16)	-0.0103 (16)
C28	0.043 (2)	0.069 (3)	0.063 (2)	-0.015 (2)	-0.0005 (18)	-0.033 (2)
C29	0.067 (3)	0.056 (3)	0.060 (2)	-0.009 (2)	-0.011 (2)	-0.0252 (19)
C30	0.078 (3)	0.057 (3)	0.083 (3)	-0.002 (2)	0.022 (2)	-0.026 (2)
C31	0.061 (3)	0.078 (3)	0.195 (5)	-0.027 (3)	0.048 (3)	-0.067 (3)
C32	0.060 (3)	0.062 (3)	0.142 (4)	-0.025 (2)	0.024 (3)	-0.049 (3)

Geometric parameters (Å, °)

C11—C5	1.742 (3)	C6—C7	1.388 (4)
C12—C13	1.737 (3)	C6—H6	0.9300
C13—C21	1.750 (3)	C7—C8	1.372 (4)
C14—C29	1.732 (3)	C7—H7A	0.9300
N1—C1	1.387 (3)	C8—H8	0.9300
N1—O2	1.424 (2)	C10—C11	1.504 (4)
N1—H1	0.9000	C10—H10A	0.9700
N2—C1	1.327 (3)	C10—H10B	0.9700
N2—H2A	0.8600	C11—C16	1.376 (4)
N2—H2B	0.8600	C11—C12	1.392 (4)
N3—C9	1.385 (3)	C12—C13	1.380 (4)
N3—O4	1.424 (2)	C12—H12	0.9300
N3—H3	0.9000	C13—C14	1.346 (4)
N4—C9	1.323 (3)	C14—C15	1.376 (4)
N4—H4A	0.8600	C14—H14	0.9300
N4—H4B	0.8600	C15—C16	1.375 (4)
N5—C17	1.386 (3)	C15—H15	0.9300
N5—O6	1.426 (3)	C16—H16	0.9300
N5—H5	0.9000	C18—C19	1.510 (4)
N6—C17	1.320 (3)	C18—H18A	0.9700
N6—H6A	0.8600	C18—H18B	0.9700
N6—H6B	0.8600	C19—C24	1.377 (4)
N7—C25	1.384 (3)	C19—C20	1.400 (4)

N7—O8	1.417 (2)	C20—C21	1.378 (4)
N7—H7	0.9000	C20—H20	0.9300
N8—C25	1.324 (3)	C21—C22	1.355 (4)
N8—H8A	0.8600	C22—C23	1.383 (4)
N8—H8B	0.8600	C22—H22	0.9300
O1—C1	1.247 (3)	C23—C24	1.371 (4)
O2—C2	1.439 (3)	C23—H23	0.9300
O3—C9	1.247 (3)	C24—H24	0.9300
O4—C10	1.428 (3)	C26—C27	1.500 (4)
O5—C17	1.249 (3)	C26—H26A	0.9700
O6—C18	1.432 (3)	C26—H26B	0.9700
O7—C25	1.248 (3)	C27—C32	1.374 (4)
O8—C26	1.432 (3)	C27—C28	1.377 (4)
C2—C3	1.507 (4)	C28—C29	1.369 (4)
C2—H2C	0.9700	C28—H28	0.9300
C2—H2D	0.9700	C29—C30	1.360 (4)
C3—C4	1.387 (4)	C30—C31	1.372 (5)
C3—C8	1.391 (4)	C30—H30	0.9300
C4—C5	1.374 (4)	C31—C32	1.364 (5)
C4—H4	0.9300	C31—H31	0.9300
C5—C6	1.366 (4)	C32—H32	0.9300
C1—N1—O2	113.2 (2)	C13—C12—H12	120.0
C1—N1—H1	108.3	C11—C12—H12	120.0
O2—N1—H1	108.2	C14—C13—C12	120.6 (3)
C1—N2—H2A	120.0	C14—C13—C12	119.7 (3)
C1—N2—H2B	120.0	C12—C13—C12	119.7 (3)
H2A—N2—H2B	120.0	C13—C14—C15	119.8 (3)
C9—N3—O4	114.3 (2)	C13—C14—H14	120.1
C9—N3—H3	108.1	C15—C14—H14	120.1
O4—N3—H3	108.1	C16—C15—C14	120.8 (3)
C9—N4—H4A	120.0	C16—C15—H15	119.6
C9—N4—H4B	120.0	C14—C15—H15	119.6
H4A—N4—H4B	120.0	C15—C16—C11	119.8 (3)
C17—N5—O6	114.9 (2)	C15—C16—H16	120.1
C17—N5—H5	108.0	C11—C16—H16	120.1
O6—N5—H5	107.9	O5—C17—N6	124.3 (3)
C17—N6—H6A	120.0	O5—C17—N5	117.3 (3)
C17—N6—H6B	120.0	N6—C17—N5	118.2 (3)
H6A—N6—H6B	120.0	O6—C18—C19	113.6 (2)
C25—N7—O8	114.0 (2)	O6—C18—H18A	108.8
C25—N7—H7	107.9	C19—C18—H18A	108.8
O8—N7—H7	107.9	O6—C18—H18B	108.8
C25—N8—H8A	120.0	C19—C18—H18B	108.8
C25—N8—H8B	120.0	H18A—C18—H18B	107.7
H8A—N8—H8B	120.0	C24—C19—C20	119.1 (3)
N1—O2—C2	110.04 (18)	C24—C19—C18	121.7 (3)
N3—O4—C10	108.3 (2)	C20—C19—C18	119.2 (3)
N5—O6—C18	108.6 (2)	C21—C20—C19	118.7 (3)
N7—O8—C26	110.3 (2)	C21—C20—H20	120.7

supplementary materials

O1—C1—N2	123.4 (3)	C19—C20—H20	120.7
O1—C1—N1	118.8 (3)	C22—C21—C20	122.2 (3)
N2—C1—N1	117.6 (3)	C22—C21—Cl3	119.4 (3)
O2—C2—C3	113.2 (2)	C20—C21—Cl3	118.4 (3)
O2—C2—H2C	108.9	C21—C22—C23	118.9 (3)
C3—C2—H2C	108.9	C21—C22—H22	120.6
O2—C2—H2D	108.9	C23—C22—H22	120.6
C3—C2—H2D	108.9	C24—C23—C22	120.5 (3)
H2C—C2—H2D	107.7	C24—C23—H23	119.8
C4—C3—C8	118.5 (3)	C22—C23—H23	119.8
C4—C3—C2	120.1 (3)	C23—C24—C19	120.6 (3)
C8—C3—C2	121.4 (3)	C23—C24—H24	119.7
C5—C4—C3	119.8 (3)	C19—C24—H24	119.7
C5—C4—H4	120.1	O7—C25—N8	123.8 (3)
C3—C4—H4	120.1	O7—C25—N7	118.3 (3)
C6—C5—C4	121.9 (3)	N8—C25—N7	117.8 (3)
C6—C5—Cl1	118.9 (3)	O8—C26—C27	114.8 (2)
C4—C5—Cl1	119.2 (3)	O8—C26—H26A	108.6
C5—C6—C7	118.7 (3)	C27—C26—H26A	108.6
C5—C6—H6	120.7	O8—C26—H26B	108.6
C7—C6—H6	120.7	C27—C26—H26B	108.6
C8—C7—C6	120.2 (3)	H26A—C26—H26B	107.5
C8—C7—H7A	119.9	C32—C27—C28	117.9 (3)
C6—C7—H7A	119.9	C32—C27—C26	120.1 (3)
C7—C8—C3	121.0 (3)	C28—C27—C26	121.9 (3)
C7—C8—H8	119.5	C29—C28—C27	120.5 (3)
C3—C8—H8	119.5	C29—C28—H28	119.8
O3—C9—N4	123.9 (3)	C27—C28—H28	119.8
O3—C9—N3	118.1 (3)	C30—C29—C28	121.4 (3)
N4—C9—N3	117.9 (3)	C30—C29—Cl4	118.9 (3)
O4—C10—C11	113.7 (2)	C28—C29—Cl4	119.7 (3)
O4—C10—H10A	108.8	C29—C30—C31	118.2 (4)
C11—C10—H10A	108.8	C29—C30—H30	120.9
O4—C10—H10B	108.8	C31—C30—H30	120.9
C11—C10—H10B	108.8	C32—C31—C30	120.8 (4)
H10A—C10—H10B	107.7	C32—C31—H31	119.6
C16—C11—C12	119.0 (3)	C30—C31—H31	119.6
C16—C11—C10	120.9 (3)	C31—C32—C27	121.1 (4)
C12—C11—C10	120.0 (3)	C31—C32—H32	119.5
C13—C12—C11	120.0 (3)	C27—C32—H32	119.5
C1—N1—O2—C2	-114.4 (2)	C12—C11—C16—C15	0.9 (5)
C9—N3—O4—C10	-110.6 (3)	C10—C11—C16—C15	-176.0 (3)
C17—N5—O6—C18	-112.7 (3)	O6—N5—C17—O5	-164.4 (2)
C25—N7—O8—C26	114.6 (3)	O6—N5—C17—N6	19.2 (4)
O2—N1—C1—O1	-166.4 (2)	N5—O6—C18—C19	-58.1 (3)
O2—N1—C1—N2	18.4 (3)	O6—C18—C19—C24	-38.7 (4)
N1—O2—C2—C3	-79.1 (3)	O6—C18—C19—C20	141.4 (3)
O2—C2—C3—C4	137.8 (3)	C24—C19—C20—C21	0.9 (5)
O2—C2—C3—C8	-45.1 (4)	C18—C19—C20—C21	-179.2 (3)

C8—C3—C4—C5	-0.1 (4)	C19—C20—C21—C22	-0.8 (5)
C2—C3—C4—C5	177.0 (3)	C19—C20—C21—C13	178.1 (2)
C3—C4—C5—C6	-0.7 (5)	C20—C21—C22—C23	0.8 (5)
C3—C4—C5—C11	178.8 (2)	C13—C21—C22—C23	-178.0 (3)
C4—C5—C6—C7	0.9 (5)	C21—C22—C23—C24	-0.9 (5)
C11—C5—C6—C7	-178.5 (2)	C22—C23—C24—C19	1.1 (5)
C5—C6—C7—C8	-0.5 (5)	C20—C19—C24—C23	-1.1 (5)
C6—C7—C8—C3	-0.3 (5)	C18—C19—C24—C23	179.0 (3)
C4—C3—C8—C7	0.5 (5)	O8—N7—C25—O7	166.5 (2)
C2—C3—C8—C7	-176.5 (3)	O8—N7—C25—N8	-17.8 (3)
O4—N3—C9—O3	-165.9 (2)	N7—O8—C26—C27	83.7 (3)
O4—N3—C9—N4	17.9 (3)	O8—C26—C27—C32	114.8 (3)
N3—O4—C10—C11	-67.9 (3)	O8—C26—C27—C28	-69.1 (4)
O4—C10—C11—C16	-38.4 (4)	C32—C27—C28—C29	2.0 (5)
O4—C10—C11—C12	144.7 (3)	C26—C27—C28—C29	-174.2 (3)
C16—C11—C12—C13	-2.2 (5)	C27—C28—C29—C30	0.3 (5)
C10—C11—C12—C13	174.7 (3)	C27—C28—C29—C14	178.9 (2)
C11—C12—C13—C14	1.7 (5)	C28—C29—C30—C31	-2.8 (6)
C11—C12—C13—C12	-178.1 (2)	C14—C29—C30—C31	178.6 (3)
C12—C13—C14—C15	0.2 (5)	C29—C30—C31—C32	2.9 (7)
C12—C13—C14—C15	180.0 (3)	C30—C31—C32—C27	-0.7 (7)
C13—C14—C15—C16	-1.6 (6)	C28—C27—C32—C31	-1.8 (6)
C14—C15—C16—C11	1.0 (6)	C26—C27—C32—C31	174.4 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots O5 ⁱ	0.90	2.20	3.096 (3)	173
N2—H2A \cdots O1 ⁱ	0.86	2.16	3.023 (3)	177
N2—H2B \cdots O3 ⁱⁱ	0.86	2.29	2.971 (3)	136
N4—H4A \cdots O7 ⁱⁱⁱ	0.86	2.11	2.971 (3)	176
N4—H4B \cdots O5	0.86	2.39	3.017 (3)	130
N5—H5 \cdots O1 ⁱ	0.90	2.19	3.090 (3)	176
N6—H6A \cdots O5 ^{iv}	0.86	2.07	2.925 (3)	177
N7—H7 \cdots O7 ^v	0.90	2.04	2.937 (3)	171
N8—H8A \cdots O3 ⁱⁱ	0.86	2.09	2.947 (3)	177
N8—H8B \cdots O1 ⁱ	0.86	2.25	2.976 (3)	142

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x+1, y, z$; (iii) $x-1, y, z$; (iv) $-x, -y+2, -z+1$; (v) $-x+1, -y+2, -z+1$.

Fig. 1




Fig. 2

