

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-(2-Bromophenyl)-N-phenyloxirane-2-carboxamide

 Long He,^{a*} Hong-Mei Qin^b and Lian-Mei Chen^a
^aCollege of Chemistry and Chemical Engineering, China West Normal University, Nanchong 637002, People's Republic of China, and ^bCollege of Life Science, China West Normal University, Nanchong 637002, People's Republic of China

Correspondence e-mail: helongcwnu@yahoo.com.cn


Received 1 November 2009; accepted 2 November 2009

 Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.028; wR factor = 0.069; data-to-parameter ratio = 15.3.

In the molecule of the title compound, $\text{C}_{15}\text{H}_{12}\text{BrNO}_2$, the two benzene rings adopt a *syn* configuration with respect to the epoxy ring; the dihedral angles between the epoxy ring and the two benzene rings are $59.90(13)$ and $68.01(12)^\circ$. Intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonding is present in the crystal structure.

Related literature

For epoxide-containing compounds used as building blocks in synthesis, see: Flisak *et al.* (1993); Watanabe *et al.* (1998); Zhu & Espenson (1995). For related structures, see: He (2009); He & Chen (2009).



Experimental

Crystal data

 $\text{C}_{15}\text{H}_{12}\text{BrNO}_2$
 $M_r = 318.17$

 Orthorhombic, $P2_12_12_1$
 $a = 6.71700(10)$ Å

 $b = 10.0370(2)$ Å

 $c = 20.4287(3)$ Å

 $V = 1377.27(4)$ Å³
 $Z = 4$

 Cu $K\alpha$ radiation

 $\mu = 4.05$ mm⁻¹
 $T = 295$ K

 $0.40 \times 0.40 \times 0.36$ mm

Data collection

Oxford Diffraction Gemini S Ultra diffractometer

Absorption correction: multi-scan

 (*CrysAlis Pro*; Oxford Diffraction, 2009)

 $T_{\min} = 0.294$, $T_{\max} = 0.324$

17721 measured reflections

2701 independent reflections

 2675 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.069$
 $S = 1.01$

2701 reflections

177 parameters

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.42$ e Å⁻³

Absolute structure: Flack (1983),

1104 Friedel pairs

 Flack parameter: $-0.008(18)$
Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O2}^i$	0.79 (3)	2.22 (3)	2.971 (2)	161 (2)
$\text{C15}-\text{H15}\cdots\text{O1}^{ii}$	0.93	2.59	3.442 (3)	153

 Symmetry codes: (i) $-x, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CrysAlis Pro* (Oxford Diffraction, 2009); cell refinement: *CrysAlis Pro*; data reduction: *CrysAlis Pro*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The diffraction data were collected at The Centre for Testing and Analysis, Sichuan University. We acknowledge financial support from China West Normal University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2664).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Flisak, J. R., Gombatz, K. J., Holmes, M. M., Jarmas, A. A., Lantos, I., Mendelson, W. L., Novack, V. J., Remich, J. J. & Snyder, L. (1993). *J. Org. Chem.* **58**, 6247–6254.
 He, L. (2009). *Acta Cryst.* **E65**, o2052.
 He, L. & Chen, L. M. (2009). *Acta Cryst.* **E65**, o2976.
 Oxford Diffraction (2009). *CrysAlis Pro*. Oxford Diffraction Ltd, Yarnton, England.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Watanabe, S., Arai, T., Sasai, H., Bougauchi, M. & Shibasaki, M. (1998). *J. Org. Chem.* **63**, 8090–8091.
 Zhu, Z. L. & Espenson, J. H. (1995). *J. Org. Chem.* **60**, 7090–7091.

supplementary materials

Acta Cryst. (2009). E65, o2999 [doi:10.1107/S1600536809045929]

3-(2-Bromophenyl)-*N*-phenyloxirane-2-carboxamide

L. He, H.-M. Qin and L.-M. Chen

Comment

Epoxides are important intermediates in organic synthesis. Glycidic esters and amides are particularly useful as they can be further transformed to key intermediates of several pharmaceutical products (Flisak *et al.* 1993; Watanabe *et al.* 1998). The Darzens reaction, is one of the most powerful methodologies for the synthesis of α , β -epoxy carbonyl and related compounds (Zhu & Espenson, 1995). We report herein the crystal structure of the title compound.

The molecular structure of (I) is shown in Fig. 1. Bond lengths and angles in (I) are normal. In the molecule, the two phenyl ring adopts a *cis* configuration about the epoxides ring. The dihedral angle between the C1—C6 and C10—C15 is $77.05(7)^\circ$, O1/C7/C8 epoxide ring makes dihedral angles of $59.90(13)^\circ$ and $68.01(12)^\circ$ with C6 and C15 phenyl ring, respectively, which is similar to that found in a related structure (He & Chen, 2009). The crystal packing is stabilized by N—H \cdots O and C—H \cdots O hydrogen bonding (Table 1).

Experimental

2-Chloro-*N*-phenylacetamide (0.17 g, 1.0 mmol) and potassium hydroxide (0.112 g, 2.0 mmol) were dissolved in acetonitrile (2 ml). To the solution was added 2-bromophenylaldehyde (0.15 g, 1.0 mmol) at 298 K, the solution was stirred for 60 min and removal of solvent under reduced pressure, the residue was purified through column chromatography. Single crystals suitable for X-ray diffraction were obtained by slow evaporation of an ethyl acetate solution at room temperature for 1 d.

Refinement

Imine H atom was located in a difference Fourier map and refined isotropically. The carbon-bound hydrogen atoms were placed in calculated positions, with C—H = 0.93–0.98 Å, and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of (I) with 30% probability displacement ellipsoids (arbitrary spheres for H atoms).

3-(2-Bromophenyl)-*N*-phenyloxirane-2-carboxamide

Crystal data

$C_{15}H_{12}BrNO_2$	$F_{000} = 640$
$M_r = 318.17$	$D_x = 1.534 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Cu $K\alpha$ radiation, $\lambda = 1.54184 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 15647 reflections
$a = 6.71700 (10) \text{ \AA}$	$\theta = 2.2\text{--}72.1^\circ$
$b = 10.0370 (2) \text{ \AA}$	$\mu = 4.05 \text{ mm}^{-1}$
$c = 20.4287 (3) \text{ \AA}$	$T = 295 \text{ K}$
$V = 1377.27 (4) \text{ \AA}^3$	Block, colorless
$Z = 4$	$0.40 \times 0.40 \times 0.36 \text{ mm}$

Data collection

Oxford Diffraction Gemini S Ultra diffractometer	2701 independent reflections
Radiation source: Enhance Ultra (Cu) X-ray Source	2675 reflections with $I > 2\sigma(I)$
Monochromator: mirror	$R_{\text{int}} = 0.028$
Detector resolution: $15.9149 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 72.3^\circ$
$T = 295 \text{ K}$	$\theta_{\text{min}} = 4.3^\circ$
ω scans	$h = -8 \rightarrow 8$
Absorption correction: multi-scan (CrysAlis Pro; Oxford Diffraction, 2009)	$k = -11 \rightarrow 12$
$T_{\text{min}} = 0.294$, $T_{\text{max}} = 0.324$	$l = -24 \rightarrow 25$
17721 measured reflections	

Refinement

Refinement on F^2	H atoms treated by a mixture of independent and constrained refinement
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.035P)^2 + 0.5235P]$
$R[F^2 > 2\sigma(F^2)] = 0.028$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.069$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.01$	$\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
2701 reflections	$\Delta\rho_{\text{min}} = -0.42 \text{ e \AA}^{-3}$
177 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008),
Primary atom site location: structure-invariant direct methods	$F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.0074 (5)
Hydrogen site location: inferred from neighbouring sites	Absolute structure: Flack (1983), 1104 Friedel pairs
	Flack parameter: $-0.008 (18)$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	-0.14003 (5)	0.63054 (3)	0.557209 (16)	0.07563 (15)
O1	-0.1933 (3)	0.23595 (16)	0.66346 (9)	0.0567 (4)
O2	-0.0059 (3)	0.50889 (16)	0.76057 (10)	0.0637 (5)
N1	0.1077 (3)	0.29560 (18)	0.75313 (9)	0.0437 (4)
C5	0.1656 (4)	0.2794 (3)	0.59008 (13)	0.0582 (6)
H5	0.1480	0.2015	0.6140	0.070*
C6	0.0186 (3)	0.3772 (2)	0.59138 (10)	0.0470 (5)
C15	0.3410 (4)	0.4080 (2)	0.82806 (11)	0.0520 (5)
H15	0.2588	0.4823	0.8314	0.062*
C9	-0.0201 (3)	0.3947 (2)	0.74074 (11)	0.0440 (4)
C8	-0.1951 (3)	0.3600 (2)	0.69813 (11)	0.0467 (5)
H8	-0.3253	0.3893	0.7143	0.056*
C10	0.2882 (3)	0.3016 (2)	0.78857 (10)	0.0407 (4)
C7	-0.1724 (3)	0.3564 (2)	0.62606 (11)	0.0499 (5)
H7	-0.2912	0.3828	0.6014	0.060*
C11	0.4155 (3)	0.1931 (2)	0.78350 (11)	0.0494 (5)
H11	0.3828	0.1227	0.7559	0.059*
C1	0.0535 (4)	0.4931 (2)	0.55591 (11)	0.0502 (5)
C12	0.5905 (4)	0.1889 (3)	0.81914 (14)	0.0607 (6)
H12	0.6734	0.1149	0.8162	0.073*
C2	0.2261 (5)	0.5119 (3)	0.52052 (13)	0.0654 (7)
H2	0.2477	0.5908	0.4977	0.078*
C3	0.3657 (5)	0.4116 (3)	0.51958 (14)	0.0723 (7)
H3	0.4815	0.4227	0.4952	0.087*
C13	0.6420 (4)	0.2934 (3)	0.85879 (13)	0.0655 (7)
H13	0.7594	0.2905	0.8829	0.079*
C14	0.5199 (4)	0.4020 (3)	0.86281 (13)	0.0641 (7)
H14	0.5567	0.4734	0.8892	0.077*
C4	0.3375 (4)	0.2965 (3)	0.55369 (14)	0.0680 (7)
H4	0.4334	0.2297	0.5525	0.082*
H1	0.074 (3)	0.225 (3)	0.7406 (12)	0.037 (6)*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0896 (2)	0.05464 (17)	0.0827 (2)	0.01486 (15)	0.01432 (17)	0.01807 (14)
O1	0.0611 (10)	0.0400 (8)	0.0689 (10)	-0.0133 (7)	-0.0136 (8)	0.0087 (7)
O2	0.0677 (10)	0.0342 (8)	0.0891 (12)	0.0036 (8)	-0.0200 (10)	-0.0011 (8)
N1	0.0487 (10)	0.0311 (8)	0.0515 (9)	-0.0032 (8)	-0.0056 (8)	-0.0002 (7)
C5	0.0618 (15)	0.0547 (13)	0.0581 (13)	0.0042 (12)	-0.0126 (12)	-0.0037 (11)
C6	0.0500 (11)	0.0458 (11)	0.0453 (10)	-0.0031 (11)	-0.0122 (8)	-0.0024 (9)
C15	0.0584 (13)	0.0455 (11)	0.0520 (11)	0.0046 (10)	-0.0047 (11)	-0.0052 (9)
C9	0.0473 (10)	0.0332 (10)	0.0515 (11)	-0.0039 (8)	-0.0011 (9)	0.0071 (8)
C8	0.0418 (10)	0.0375 (10)	0.0609 (12)	-0.0063 (9)	-0.0036 (8)	0.0089 (10)
C10	0.0442 (10)	0.0386 (10)	0.0392 (9)	-0.0030 (8)	0.0020 (8)	0.0079 (8)
C7	0.0459 (11)	0.0448 (11)	0.0589 (12)	-0.0055 (10)	-0.0150 (9)	0.0106 (10)
C11	0.0517 (12)	0.0425 (11)	0.0541 (12)	0.0011 (9)	0.0042 (9)	0.0029 (9)
C1	0.0604 (12)	0.0462 (11)	0.0441 (10)	-0.0012 (9)	-0.0009 (10)	-0.0020 (10)
C12	0.0477 (12)	0.0615 (14)	0.0730 (16)	0.0100 (11)	0.0029 (11)	0.0138 (12)
C2	0.0782 (17)	0.0652 (16)	0.0528 (13)	-0.0086 (14)	0.0095 (12)	-0.0001 (12)
C3	0.0631 (16)	0.091 (2)	0.0629 (15)	-0.0026 (16)	0.0099 (14)	-0.0143 (14)
C13	0.0525 (13)	0.0838 (19)	0.0602 (13)	0.0019 (15)	-0.0100 (12)	0.0111 (13)
C14	0.0696 (16)	0.0679 (16)	0.0548 (13)	-0.0056 (14)	-0.0149 (12)	-0.0083 (12)
C4	0.0624 (15)	0.0748 (17)	0.0669 (15)	0.0141 (13)	-0.0095 (15)	-0.0156 (14)

Geometric parameters (\AA , $^\circ$)

Br1—C1	1.896 (2)	C8—H8	0.9800
O1—C8	1.433 (3)	C10—C11	1.388 (3)
O1—C7	1.437 (3)	C7—H7	0.9800
O2—C9	1.219 (3)	C11—C12	1.384 (4)
N1—C9	1.338 (3)	C11—H11	0.9300
N1—C10	1.414 (3)	C1—C2	1.380 (4)
N1—H1	0.78 (3)	C12—C13	1.370 (4)
C5—C4	1.384 (4)	C12—H12	0.9300
C5—C6	1.392 (3)	C2—C3	1.375 (5)
C5—H5	0.9300	C2—H2	0.9300
C6—C1	1.390 (3)	C3—C4	1.363 (5)
C6—C7	1.481 (3)	C3—H3	0.9300
C15—C10	1.385 (3)	C13—C14	1.366 (4)
C15—C14	1.397 (4)	C13—H13	0.9300
C15—H15	0.9300	C14—H14	0.9300
C9—C8	1.503 (3)	C4—H4	0.9300
C8—C7	1.481 (3)		
C8—O1—C7	62.11 (14)	O1—C7—H7	114.9
C9—N1—C10	127.97 (19)	C6—C7—H7	114.9
C9—N1—H1	115.1 (18)	C8—C7—H7	114.9
C10—N1—H1	116.8 (18)	C12—C11—C10	120.6 (2)
C4—C5—C6	121.0 (3)	C12—C11—H11	119.7

C4—C5—H5	119.5	C10—C11—H11	119.7
C6—C5—H5	119.5	C2—C1—C6	121.9 (2)
C1—C6—C5	117.4 (2)	C2—C1—Br1	118.96 (19)
C1—C6—C7	120.9 (2)	C6—C1—Br1	119.09 (17)
C5—C6—C7	121.7 (2)	C13—C12—C11	120.1 (2)
C10—C15—C14	118.9 (2)	C13—C12—H12	119.9
C10—C15—H15	120.6	C11—C12—H12	119.9
C14—C15—H15	120.6	C3—C2—C1	118.7 (3)
O2—C9—N1	125.9 (2)	C3—C2—H2	120.6
O2—C9—C8	118.1 (2)	C1—C2—H2	120.6
N1—C9—C8	116.05 (19)	C4—C3—C2	121.2 (3)
O1—C8—C7	59.09 (14)	C4—C3—H3	119.4
O1—C8—C9	118.74 (18)	C2—C3—H3	119.4
C7—C8—C9	120.07 (19)	C14—C13—C12	119.7 (2)
O1—C8—H8	115.7	C14—C13—H13	120.2
C7—C8—H8	115.7	C12—C13—H13	120.2
C9—C8—H8	115.7	C13—C14—C15	121.3 (2)
C15—C10—C11	119.4 (2)	C13—C14—H14	119.3
C15—C10—N1	123.5 (2)	C15—C14—H14	119.3
C11—C10—N1	117.1 (2)	C3—C4—C5	119.7 (3)
O1—C7—C6	117.2 (2)	C3—C4—H4	120.1
O1—C7—C8	58.80 (14)	C5—C4—H4	120.1
C6—C7—C8	124.15 (18)		
C4—C5—C6—C1	-1.3 (3)	C9—C8—C7—O1	107.5 (2)
C4—C5—C6—C7	175.4 (2)	O1—C8—C7—C6	-103.6 (2)
C10—N1—C9—O2	-3.1 (4)	C9—C8—C7—C6	3.9 (4)
C10—N1—C9—C8	176.5 (2)	C15—C10—C11—C12	2.1 (3)
C7—O1—C8—C9	-109.7 (2)	N1—C10—C11—C12	-176.9 (2)
O2—C9—C8—O1	167.2 (2)	C5—C6—C1—C2	0.1 (3)
N1—C9—C8—O1	-12.5 (3)	C7—C6—C1—C2	-176.7 (2)
O2—C9—C8—C7	98.3 (3)	C5—C6—C1—Br1	-178.98 (16)
N1—C9—C8—C7	-81.4 (3)	C7—C6—C1—Br1	4.2 (3)
C14—C15—C10—C11	-1.2 (3)	C10—C11—C12—C13	-1.4 (4)
C14—C15—C10—N1	177.8 (2)	C6—C1—C2—C3	1.1 (4)
C9—N1—C10—C15	14.1 (3)	Br1—C1—C2—C3	-179.8 (2)
C9—N1—C10—C11	-166.9 (2)	C1—C2—C3—C4	-1.2 (4)
C8—O1—C7—C6	115.3 (2)	C11—C12—C13—C14	-0.2 (4)
C1—C6—C7—O1	-177.83 (19)	C12—C13—C14—C15	1.1 (4)
C5—C6—C7—O1	5.5 (3)	C10—C15—C14—C13	-0.4 (4)
C1—C6—C7—C8	-108.6 (2)	C2—C3—C4—C5	0.0 (4)
C5—C6—C7—C8	74.7 (3)	C6—C5—C4—C3	1.3 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O2 ⁱ	0.79 (3)	2.22 (3)	2.971 (2)	161 (2)
C15—H15 \cdots O1 ⁱⁱ	0.93	2.59	3.442 (3)	153

Symmetry codes: (i) $-x, y-1/2, -z+3/2$; (ii) $-x, y+1/2, -z+3/2$.

Fig. 1

