

Bis(trimethylphenylammonium) tetra-bromidobis(4-chlorophenyl)stannate(IV)

Kong Mun Lo and Seik Weng Ng*

 Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
 Correspondence e-mail: seikweng@um.edu.my


Received 18 November 2009; accepted 19 November 2009

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å;
 R factor = 0.031; wR factor = 0.098; data-to-parameter ratio = 21.6.

The Sn^{IV} atom in the title salt, $[\text{N}(\text{CH}_3)_3(\text{C}_6\text{H}_5)]_2\text{[SnBr}_4(\text{C}_6\text{H}_4\text{Cl})_2]$, exists in a distorted all-*trans* SnC_2Br_4 octahedral geometry. The Sn^{IV} atom lies on a center of inversion. Weak intermolecular $\text{C}-\text{H}\cdots\text{Br}$ hydrogen bonding is observed between trimethylphenylammonium cations and the Sn complex anion in the crystal structure.

Related literature

For bis(4-dimethylaminopyridinium) tetrabromidodiphenylstannate, see: Yap *et al.* (2008).



Experimental

Crystal data

 $(\text{C}_9\text{H}_{14}\text{N})_2[\text{SnBr}_4(\text{C}_6\text{H}_4\text{Cl})_2]$
 $M_r = 933.84$

 Monoclinic, $C2/c$
 $a = 25.7930$ (3) Å

 $b = 9.0937$ (1) Å
 $c = 15.8303$ (2) Å
 $\beta = 113.4146$ (6)°
 $V = 3407.30$ (7) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 5.62$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.25 \times 0.20$ mm

Data collection

 Bruker SMART APEX
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.482$, $T_{\text{max}} = 0.756$

 15816 measured reflections
 3916 independent reflections
 3427 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.098$
 $S = 1.24$
 3916 reflections

 181 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.28$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.03$ e Å⁻³
Table 1

Selected bond lengths (Å).

Sn1—C1	2.156 (3)	Sn1—Br2	2.7386 (4)
Sn1—Br1	2.7368 (4)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C15—H15C \cdots Br1	0.96	2.91	3.839 (5)	163

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

We thank the University of Malaya (grant No. RG020/09AFR) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2683).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2009). publCIF. In preparation.
 Yap, Q. L., Lo, K. M. & Ng, S. W. (2008). *Acta Cryst.* **E64**, m696.

supplementary materials

Acta Cryst. (2009). E65, m1677 [doi:10.1107/S160053680904940X]

Bis(trimethylphenylammonium) tetrabromidobis(4-chlorophenyl)stannate(IV)

K. M. Lo and S. W. Ng

Experimental

In an attempt to cleave a tin-carbon bond in a tetraorganotin compound, tetrakis(4-chlorophenyl)tin (0.57 g, 1 mmol) and trimethylphenylammonium tribromide (0.38 g, 1 mmol) were heated in ethanol for six hours. Colorless crystals of the stannate separated from the solution after a few days.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93–0.96 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to 1.2–1.5 $U(C)$.

The final difference Fourier map had peak near Sn1 and a hole near Br2.

The refinement program suggested a rather larger second parameter for the weighting scheme. The scheme was arbitrarily selected as (0.05 5.00), which led to a satisfactory, albeit somewhat large, goodness-of-fit.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of the two independent formula units of $2[N(CH_3)_3(C_6H_5)] [SnBr_4(C_6H_4Cl)_2]$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

Bis(trimethylphenylammonium) tetrabromidobis(4-chlorophenyl)stannate(IV)

Crystal data

$(C_9H_{14}N)_2[SnBr_4(C_6H_4Cl)_2]$

$M_r = 933.84$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 25.7930$ (3) Å

$b = 9.0937$ (1) Å

$c = 15.8303$ (2) Å

$\beta = 113.4146$ (6)°

$V = 3407.30$ (7) Å³

$F(000) = 1816$

$D_x = 1.820$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 946 reflections

$\theta = 2.4$ – 28.2 °

$\mu = 5.62$ mm⁻¹

$T = 293$ K

Block, colorless

$0.30 \times 0.25 \times 0.20$ mm

supplementary materials

Z = 4

Data collection

Bruker SMART APEX diffractometer	3916 independent reflections
Radiation source: fine-focus sealed tube	3427 reflections with $I > 2\sigma(I)$
graphite	$R_{\text{int}} = 0.025$
ω scans	$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 1.7^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -33 \rightarrow 33$
$T_{\text{min}} = 0.482$, $T_{\text{max}} = 0.756$	$k = -11 \rightarrow 11$
15816 measured reflections	$l = -20 \rightarrow 20$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.031$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.098$	H-atom parameters constrained
$S = 1.24$	$w = 1/[\sigma^2(F_o^2) + (0.05P)^2 + 5.0P]$
3916 reflections	where $P = (F_o^2 + 2F_c^2)/3$
181 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 1.28 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\text{min}} = -1.03 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.2500	0.7500	0.5000	0.02955 (10)
Br1	0.183541 (17)	0.69102 (4)	0.59433 (3)	0.04572 (12)
Br2	0.292712 (17)	0.99167 (4)	0.60875 (3)	0.04584 (12)
Cl1	0.45173 (6)	0.34442 (18)	0.80689 (10)	0.0818 (4)
N1	0.13915 (14)	0.2207 (3)	0.5931 (2)	0.0400 (7)
C1	0.31591 (13)	0.6149 (3)	0.5957 (2)	0.0295 (6)
C2	0.30386 (17)	0.4780 (4)	0.6223 (3)	0.0422 (8)
H2	0.2670	0.4426	0.5964	0.051*
C3	0.34514 (18)	0.3939 (4)	0.6858 (3)	0.0466 (9)
H3	0.3366	0.3028	0.7037	0.056*
C4	0.39961 (17)	0.4473 (5)	0.7226 (3)	0.0473 (9)
C5	0.41310 (16)	0.5804 (5)	0.6971 (3)	0.0518 (10)
H5	0.4502	0.6141	0.7222	0.062*
C6	0.37092 (15)	0.6646 (4)	0.6334 (3)	0.0420 (8)
H6	0.3798	0.7556	0.6158	0.050*
C7	0.08627 (16)	0.2803 (4)	0.5981 (2)	0.0383 (8)
C8	0.0346 (2)	0.2255 (6)	0.5397 (4)	0.0658 (13)

H8	0.0325	0.1501	0.4988	0.079*
C9	-0.0136 (2)	0.2830 (8)	0.5424 (5)	0.0817 (17)
H9	-0.0485	0.2469	0.5022	0.098*
C10	-0.0121 (2)	0.3909 (6)	0.6018 (4)	0.0702 (14)
H10	-0.0454	0.4290	0.6026	0.084*
C11	0.0390 (3)	0.4433 (6)	0.6607 (4)	0.0732 (15)
H11	0.0403	0.5161	0.7027	0.088*
C12	0.0888 (2)	0.3908 (5)	0.6594 (3)	0.0564 (11)
H12	0.1233	0.4291	0.6990	0.068*
C13	0.1911 (2)	0.2531 (7)	0.6789 (3)	0.0683 (14)
H13A	0.1960	0.3575	0.6867	0.102*
H13B	0.2236	0.2107	0.6731	0.102*
H13C	0.1867	0.2115	0.7313	0.102*
C14	0.1361 (2)	0.0572 (5)	0.5795 (5)	0.0728 (15)
H14A	0.1059	0.0337	0.5218	0.109*
H14B	0.1292	0.0111	0.6286	0.109*
H14C	0.1712	0.0222	0.5795	0.109*
C15	0.1479 (2)	0.2913 (6)	0.5144 (3)	0.0573 (11)
H15A	0.1171	0.2660	0.4577	0.086*
H15B	0.1827	0.2570	0.5129	0.086*
H15C	0.1494	0.3961	0.5221	0.086*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.02499 (16)	0.03104 (17)	0.03048 (17)	0.00219 (11)	0.00875 (12)	0.00523 (11)
Br1	0.0450 (2)	0.0460 (2)	0.0533 (2)	0.00187 (16)	0.02717 (18)	0.00690 (16)
Br2	0.0481 (2)	0.0417 (2)	0.0424 (2)	-0.00224 (16)	0.01229 (17)	-0.00120 (15)
Cl1	0.0651 (8)	0.0921 (9)	0.0723 (8)	0.0379 (7)	0.0105 (6)	0.0363 (7)
N1	0.0452 (18)	0.0404 (15)	0.0392 (16)	-0.0008 (13)	0.0219 (14)	0.0015 (12)
C1	0.0289 (15)	0.0301 (14)	0.0286 (14)	0.0038 (12)	0.0105 (12)	0.0032 (11)
C2	0.040 (2)	0.0401 (18)	0.0415 (19)	-0.0019 (16)	0.0111 (16)	0.0051 (15)
C3	0.055 (2)	0.0371 (18)	0.043 (2)	0.0073 (17)	0.0149 (18)	0.0098 (15)
C4	0.044 (2)	0.053 (2)	0.0394 (19)	0.0212 (18)	0.0114 (16)	0.0108 (17)
C5	0.0280 (18)	0.062 (3)	0.057 (2)	0.0037 (18)	0.0080 (17)	0.010 (2)
C6	0.0301 (17)	0.0438 (19)	0.048 (2)	0.0015 (15)	0.0110 (15)	0.0100 (16)
C7	0.040 (2)	0.0405 (18)	0.0363 (18)	-0.0006 (15)	0.0168 (16)	0.0021 (14)
C8	0.049 (3)	0.081 (3)	0.058 (3)	-0.008 (2)	0.012 (2)	-0.021 (2)
C9	0.039 (3)	0.102 (4)	0.089 (4)	-0.008 (3)	0.010 (3)	-0.007 (3)
C10	0.053 (3)	0.069 (3)	0.099 (4)	0.012 (2)	0.041 (3)	0.016 (3)
C11	0.083 (4)	0.055 (3)	0.098 (4)	0.009 (3)	0.053 (3)	-0.009 (3)
C12	0.051 (2)	0.051 (2)	0.068 (3)	-0.005 (2)	0.024 (2)	-0.014 (2)
C13	0.047 (3)	0.105 (4)	0.047 (2)	0.016 (3)	0.013 (2)	-0.003 (2)
C14	0.082 (4)	0.040 (2)	0.120 (5)	0.001 (2)	0.065 (4)	-0.001 (3)
C15	0.070 (3)	0.064 (3)	0.051 (2)	-0.006 (2)	0.038 (2)	0.004 (2)

supplementary materials

Geometric parameters (Å, °)

Sn1—C1 ⁱ	2.156 (3)	C7—C12	1.380 (6)
Sn1—C1	2.156 (3)	C7—C8	1.380 (6)
Sn1—Br1	2.7368 (4)	C8—C9	1.365 (8)
Sn1—Br1 ⁱ	2.7368 (4)	C8—H8	0.9300
Sn1—Br2 ⁱ	2.7386 (4)	C9—C10	1.349 (8)
Sn1—Br2	2.7386 (4)	C9—H9	0.9300
C11—C4	1.743 (4)	C10—C11	1.362 (8)
N1—C14	1.499 (5)	C10—H10	0.9300
N1—C15	1.495 (5)	C11—C12	1.377 (7)
N1—C7	1.499 (5)	C11—H11	0.9300
N1—C13	1.510 (6)	C12—H12	0.9300
C1—C6	1.379 (5)	C13—H13A	0.9600
C1—C2	1.389 (5)	C13—H13B	0.9600
C2—C3	1.371 (5)	C13—H13C	0.9600
C2—H2	0.9300	C14—H14A	0.9600
C3—C4	1.378 (6)	C14—H14B	0.9600
C3—H3	0.9300	C14—H14C	0.9600
C4—C5	1.364 (6)	C15—H15A	0.9600
C5—C6	1.384 (5)	C15—H15B	0.9600
C5—H5	0.9300	C15—H15C	0.9600
C6—H6	0.9300		
C1 ⁱ —Sn1—C1	180.000 (1)	C1—C6—H6	119.7
C1 ⁱ —Sn1—Br1	90.29 (8)	C12—C7—C8	119.8 (4)
C1—Sn1—Br1	89.71 (8)	C12—C7—N1	120.8 (4)
C1 ⁱ —Sn1—Br1 ⁱ	89.71 (8)	C8—C7—N1	119.4 (4)
C1—Sn1—Br1 ⁱ	90.29 (8)	C9—C8—C7	119.4 (5)
Br1—Sn1—Br1 ⁱ	180.0	C9—C8—H8	120.3
C1 ⁱ —Sn1—Br2 ⁱ	90.43 (8)	C7—C8—H8	120.3
C1—Sn1—Br2 ⁱ	89.57 (8)	C8—C9—C10	121.7 (5)
Br1—Sn1—Br2 ⁱ	90.173 (13)	C8—C9—H9	119.1
Br1 ⁱ —Sn1—Br2 ⁱ	89.827 (12)	C10—C9—H9	119.1
C1 ⁱ —Sn1—Br2	89.57 (8)	C11—C10—C9	118.9 (5)
C1—Sn1—Br2	90.43 (8)	C11—C10—H10	120.5
Br1—Sn1—Br2	89.827 (12)	C9—C10—H10	120.5
Br1 ⁱ —Sn1—Br2	90.173 (13)	C10—C11—C12	121.5 (5)
Br2 ⁱ —Sn1—Br2	180.000 (10)	C10—C11—H11	119.2
C14—N1—C15	108.9 (4)	C12—C11—H11	119.2
C14—N1—C7	111.7 (3)	C11—C12—C7	118.6 (5)
C15—N1—C7	109.3 (3)	C11—C12—H12	120.7
C14—N1—C13	107.4 (4)	C7—C12—H12	120.7
C15—N1—C13	107.0 (4)	N1—C13—H13A	109.5
C7—N1—C13	112.5 (3)	N1—C13—H13B	109.5
C6—C1—C2	118.6 (3)	H13A—C13—H13B	109.5

C6—C1—Sn1	120.4 (2)	N1—C13—H13C	109.5
C2—C1—Sn1	121.0 (3)	H13A—C13—H13C	109.5
C3—C2—C1	121.4 (4)	H13B—C13—H13C	109.5
C3—C2—H2	119.3	N1—C14—H14A	109.5
C1—C2—H2	119.3	N1—C14—H14B	109.5
C2—C3—C4	118.6 (4)	H14A—C14—H14B	109.5
C2—C3—H3	120.7	N1—C14—H14C	109.5
C4—C3—H3	120.7	H14A—C14—H14C	109.5
C5—C4—C3	121.6 (3)	H14B—C14—H14C	109.5
C5—C4—C11	119.5 (3)	N1—C15—H15A	109.5
C3—C4—C11	118.8 (3)	N1—C15—H15B	109.5
C4—C5—C6	119.2 (4)	H15A—C15—H15B	109.5
C4—C5—H5	120.4	N1—C15—H15C	109.5
C6—C5—H5	120.4	H15A—C15—H15C	109.5
C5—C6—C1	120.6 (4)	H15B—C15—H15C	109.5
C5—C6—H6	119.7		
Br1—Sn1—C1—C6	-133.2 (3)	C2—C1—C6—C5	-0.7 (6)
Br1 ⁱ —Sn1—C1—C6	46.8 (3)	Sn1—C1—C6—C5	177.9 (3)
Br2 ⁱ —Sn1—C1—C6	136.6 (3)	C14—N1—C7—C12	140.8 (5)
Br2—Sn1—C1—C6	-43.4 (3)	C15—N1—C7—C12	-98.7 (5)
Br1—Sn1—C1—C2	45.3 (3)	C13—N1—C7—C12	20.0 (5)
Br1 ⁱ —Sn1—C1—C2	-134.7 (3)	C14—N1—C7—C8	-40.4 (6)
Br2 ⁱ —Sn1—C1—C2	-44.8 (3)	C15—N1—C7—C8	80.2 (5)
Br2—Sn1—C1—C2	135.2 (3)	C13—N1—C7—C8	-161.2 (4)
C6—C1—C2—C3	1.2 (6)	C12—C7—C8—C9	0.9 (8)
Sn1—C1—C2—C3	-177.4 (3)	N1—C7—C8—C9	-178.0 (5)
C1—C2—C3—C4	-0.8 (6)	C7—C8—C9—C10	-1.0 (10)
C2—C3—C4—C5	-0.2 (6)	C8—C9—C10—C11	-0.1 (10)
C2—C3—C4—C11	177.4 (3)	C9—C10—C11—C12	1.4 (9)
C3—C4—C5—C6	0.7 (7)	C10—C11—C12—C7	-1.5 (8)
C11—C4—C5—C6	-176.9 (3)	C8—C7—C12—C11	0.3 (7)
C4—C5—C6—C1	-0.2 (7)	N1—C7—C12—C11	179.2 (4)

Symmetry codes: (i) $-x+1/2, -y+3/2, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C15—H15C \cdots Br1	0.96	2.91	3.839 (5)	163

Fig. 1

