

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

Dibromidodi- $\mu$ -hydroxido-di- $\mu_3$ -oxido-octaphenyltetratin(IV)

Quai Ling Yap, Kong Mun Lo and Seik Weng Ng\*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 16 November 2009; accepted 1 December 2009

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.015$  Å;  
 $R$  factor = 0.042;  $wR$  factor = 0.160; data-to-parameter ratio = 20.3.

In the centrosymmetric title compound,  $[\text{Sn}_4\text{Br}_2(\text{C}_6\text{H}_5)_8\text{O}_2(\text{OH})_2]$ , the four tin(IV) atoms are bridged by the hydroxo and oxo ligands, forming a ladder-like array of three edge-connected  $\text{Sn}_2\text{O}_2$  squares. The two independent tin atoms show distorted trigonal-bipyramidal  $\text{SnC}_2\text{O}_3$  and  $\text{SnC}_2\text{O}_2\text{Br}$  coordination geometries.

## Related literature

For other  $[\text{Sn}_4\text{X}_2(\text{O})_2(\text{OH})_2\text{R}_8]$  ( $X = \text{halogen}$ ,  $R = \text{organic group}$ ) structures, see: Baumeister *et al.* (2002); Beckmann *et al.* (2001); Cox & Tiekink (1994); Kresinski *et al.* (1994); Lo & Ng (2009); Mohamed *et al.* (2004); Puff *et al.* (1983); Tiekink (1991); Vollano *et al.* (1984).



## Experimental

## Crystal data

 $[\text{Sn}_4\text{Br}_2(\text{C}_6\text{H}_5)_8\text{O}_2(\text{OH})_2]$   
 $M_r = 1317.40$   
 Triclinic,  $P\bar{1}$ 
 $a = 10.3759$  (1) Å  
 $b = 10.8247$  (1) Å  
 $c = 12.0341$  (2) Å

 $\alpha = 77.1809$  (7)°  
 $\beta = 65.4426$  (7)°  
 $\gamma = 75.2040$  (7)°  
 $V = 1178.26$  (3) Å<sup>3</sup>  
 $Z = 1$ 

 Mo  $K\alpha$  radiation  
 $\mu = 3.83$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.25 \times 0.25 \times 0.25$  mm

## Data collection

 Bruker SMART APEX  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.539$ ,  $T_{\max} = 0.746$ 

 10751 measured reflections  
 5309 independent reflections  
 4449 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.160$   
 $S = 1.06$   
 5309 reflections

 262 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 2.39$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -1.38$  e Å<sup>-3</sup>

Table 1

Selected bond lengths (Å).

Sn1—O1 <sup>i</sup>	2.068 (4)	Sn2—O1 <sup>i</sup>	2.029 (4)
Sn1—O1	2.095 (4)	Sn2—C13	2.118 (7)
Sn1—C1	2.117 (6)	Sn2—C19	2.126 (7)
Sn1—C7	2.120 (6)	Sn2—O2	2.167 (4)
Sn1—O2	2.171 (4)	Sn2—Br1	2.6444 (9)

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

Data collection: APEX2 software (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

We thank the University of Malaya (grant No. RG020/09AFR) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5234).

## References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
 Baumeister, U., Dakternieks, D., Jurkschat, K. & Schürmann, M. (2002). *Main Group Met. Chem.* **26**, 521–522.  
 Beckmann, J., Jurkschat, K., Rabe, S., Schürmann, M. & Dakternieks, D. (2001). *Z. Anorg. Allg. Chem.* **627**, 458–464.  
 Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Cox, M. J. & Tiekink, E. R. T. (1994). *Z. Kristallogr.* **209**, 622–623.  
 Kresinski, R. A., Staples, R. J. & Fackler, J. P. (1994). *Acta Cryst.* **C50**, 40–41.  
 Lo, K. M. & Ng, S. W. (2009). *Acta Cryst.* **E65**, m593.  
 Mohamed, E. M., Panchanatheswaran, K., Low, J. N. & Glidewell, C. (2004). *Acta Cryst.* **E60**, m489–m491.  
 Puff, H., Bung, I., Friedrichs, E. & Jansen, A. (1983). *J. Organomet. Chem.* **254**, 23–32.  
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Tiekink, E. R. T. (1991). *Acta Cryst.* **C47**, 661–662.  
 Vollano, J. F., Day, R. O. & Holmes, R. R. (1984). *Organometallics*, **3**, 745–750.  
 Westrip, S. P. (2009). publCIF. In preparation.

---

## supporting information

*Acta Cryst.* (2010). E66, m8 [doi:10.1107/S1600536809051691]

### **Dibromidodi- $\mu$ -hydroxido-di- $\mu_3$ -oxido-octaphenyltin(IV)**

**Quai Ling Yap, Kong Mun Lo and Seik Weng Ng**


#### **S1. Experimental**

In an attempt to cleave a tin-carbon bond in a tetraorganotin compound, bis(3-methyl-2-thienyl)diphenyltin (0.47 g, 1 mmol) and trimethylphenylammonium tribromide (0.38 g, 1 mmol) were heated in ethanol for three hours. Colourless blocks of (I) separated from solution after a few days.

#### **S2. Refinement**

Carbon- and oxygen-bound H-atoms were placed in calculated positions (C–H 0.93 Å) and were included in the refinement in the riding model approximation, with  $U(\text{H})$  set to  $1.2U(\text{C}, \text{O})$ .

The final difference Fourier map had a peak near Sn1 and a hole near Sn2.

**Figure 1**

View of (I) at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. Unlabelled atoms are generated by the symmetry operation  $(1-x, 1-y, 1-z)$ .

### Dibromidodi- $\mu$ -hydroxido-di- $\mu_3$ -oxido-octaphenyltetra tin(IV)

#### Crystal data

$[\text{Sn}_4\text{Br}_2(\text{C}_6\text{H}_5)_8\text{O}_2(\text{OH})_2]$

$M_r = 1317.40$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 10.3759$  (1) Å

$b = 10.8247$  (1) Å

$c = 12.0341$  (2) Å

$\alpha = 77.1809$  (7)°

$\beta = 65.4426$  (7)°

$\gamma = 75.2040$  (7)°

$V = 1178.26$  (3) Å<sup>3</sup>

$Z = 1$

$F(000) = 632$

$D_x = 1.857$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 7354 reflections

$\theta = 2.2\text{--}28.5^\circ$

$\mu = 3.83$  mm<sup>-1</sup>

$T = 293$  K

Block, colourless

$0.25 \times 0.25 \times 0.25$  mm

#### Data collection

Bruker SMART APEX  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.539$ ,  $T_{\max} = 0.746$

10751 measured reflections  
 5309 independent reflections  
 4449 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

$\theta_{\text{max}} = 27.5^\circ$ ,  $\theta_{\text{min}} = 1.9^\circ$   
 $h = -13 \rightarrow 13$   
 $k = -14 \rightarrow 13$   
 $l = -15 \rightarrow 15$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.160$   
 $S = 1.06$   
 5309 reflections  
 262 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0952P)^2 + 4.7584P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 2.39 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -1.38 \text{ e } \text{\AA}^{-3}$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.38584 (4)	0.45011 (4)	0.46172 (4)	0.03147 (14)
Sn2	0.52171 (5)	0.20525 (4)	0.63535 (4)	0.03495 (15)
Br1	0.70321 (9)	0.23768 (8)	0.72424 (7)	0.0508 (2)
O1	0.4796 (5)	0.6078 (4)	0.4416 (4)	0.0315 (8)
O2	0.3774 (5)	0.2494 (4)	0.5377 (4)	0.0364 (9)
H2O	0.3312	0.2022	0.5311	0.044*
C1	0.4869 (8)	0.4083 (7)	0.2777 (6)	0.0437 (15)
C2	0.6000 (10)	0.4680 (9)	0.1951 (7)	0.062 (2)
H2	0.6262	0.5303	0.2198	0.074*
C3	0.6748 (14)	0.4366 (14)	0.0765 (9)	0.092 (4)
H3	0.7537	0.4739	0.0230	0.111*
C4	0.6308 (17)	0.3499 (15)	0.0393 (11)	0.104 (5)
H4	0.6786	0.3303	-0.0410	0.125*
C5	0.5169 (15)	0.2913 (12)	0.1190 (11)	0.088 (4)
H5	0.4865	0.2338	0.0921	0.106*
C6	0.4471 (11)	0.3183 (9)	0.2404 (8)	0.060 (2)
H6	0.3736	0.2753	0.2959	0.072*
C7	0.1737 (7)	0.5207 (7)	0.5813 (6)	0.0379 (13)
C8	0.0633 (8)	0.4546 (8)	0.6167 (8)	0.0524 (18)
H8	0.0782	0.3810	0.5819	0.063*
C9	-0.0724 (10)	0.4978 (12)	0.7057 (10)	0.075 (3)
H9	-0.1485	0.4550	0.7284	0.090*
C10	-0.0900 (10)	0.6034 (13)	0.7578 (9)	0.082 (3)
H10	-0.1780	0.6296	0.8196	0.098*
C11	0.0158 (11)	0.6711 (12)	0.7228 (11)	0.084 (3)
H11	-0.0008	0.7451	0.7576	0.100*
C12	0.1506 (9)	0.6298 (9)	0.6340 (8)	0.0554 (19)
H12	0.2247	0.6754	0.6103	0.067*
C13	0.3559 (8)	0.1680 (7)	0.8092 (6)	0.0453 (15)
C14	0.3843 (12)	0.0721 (11)	0.8954 (9)	0.077 (3)

H14	0.4784	0.0286	0.8804	0.093*
C15	0.2745 (16)	0.0389 (14)	1.0052 (10)	0.103 (4)
H15	0.2949	-0.0282	1.0623	0.123*
C16	0.1375 (15)	0.1039 (13)	1.0295 (10)	0.091 (4)
H16	0.0637	0.0796	1.1022	0.109*
C17	0.1077 (13)	0.2043 (14)	0.9484 (10)	0.094 (4)
H17	0.0144	0.2512	0.9670	0.113*
C18	0.2175 (10)	0.2370 (10)	0.8367 (8)	0.065 (2)
H18	0.1971	0.3056	0.7809	0.078*
C19	0.6770 (7)	0.0737 (6)	0.5153 (6)	0.0388 (13)
C20	0.7810 (9)	-0.0145 (8)	0.5494 (8)	0.0527 (18)
H20	0.7787	-0.0227	0.6288	0.063*
C21	0.8893 (9)	-0.0909 (9)	0.4642 (9)	0.062 (2)
H21	0.9578	-0.1515	0.4879	0.074*
C22	0.8961 (9)	-0.0784 (8)	0.3475 (9)	0.059 (2)
H22	0.9705	-0.1285	0.2911	0.071*
C23	0.7926 (9)	0.0089 (8)	0.3123 (8)	0.0541 (19)
H23	0.7975	0.0177	0.2321	0.065*
C24	0.6816 (9)	0.0834 (7)	0.3962 (7)	0.0495 (17)
H24	0.6101	0.1399	0.3732	0.059*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Sn1	0.0317 (2)	0.0315 (2)	0.0317 (2)	-0.00616 (16)	-0.01214 (17)	-0.00462 (15)
Sn2	0.0381 (2)	0.0287 (2)	0.0345 (2)	-0.00560 (17)	-0.01235 (18)	-0.00091 (16)
Br1	0.0555 (4)	0.0517 (4)	0.0531 (4)	-0.0067 (3)	-0.0305 (3)	-0.0063 (3)
O1	0.033 (2)	0.026 (2)	0.036 (2)	-0.0070 (16)	-0.0157 (17)	0.0008 (16)
O2	0.040 (2)	0.030 (2)	0.045 (2)	-0.0123 (18)	-0.0197 (19)	-0.0005 (17)
C1	0.055 (4)	0.039 (3)	0.033 (3)	0.000 (3)	-0.017 (3)	-0.008 (3)
C2	0.068 (5)	0.070 (6)	0.040 (4)	-0.024 (5)	-0.006 (4)	-0.010 (4)
C3	0.104 (9)	0.114 (10)	0.041 (5)	-0.033 (8)	-0.002 (5)	-0.014 (5)
C4	0.127 (11)	0.120 (11)	0.050 (6)	-0.001 (9)	-0.019 (7)	-0.036 (6)
C5	0.123 (10)	0.082 (7)	0.080 (7)	0.005 (7)	-0.055 (7)	-0.046 (6)
C6	0.069 (5)	0.061 (5)	0.061 (5)	-0.005 (4)	-0.032 (4)	-0.024 (4)
C7	0.031 (3)	0.041 (3)	0.038 (3)	-0.005 (3)	-0.012 (2)	-0.004 (3)
C8	0.039 (4)	0.049 (4)	0.066 (5)	-0.010 (3)	-0.017 (3)	-0.005 (3)
C9	0.044 (5)	0.096 (8)	0.077 (6)	-0.024 (5)	-0.016 (4)	0.002 (6)
C10	0.044 (5)	0.114 (9)	0.061 (6)	-0.008 (5)	0.006 (4)	-0.018 (6)
C11	0.058 (6)	0.086 (8)	0.090 (7)	-0.003 (5)	-0.001 (5)	-0.046 (6)
C12	0.042 (4)	0.059 (5)	0.060 (5)	-0.011 (4)	-0.007 (3)	-0.021 (4)
C13	0.051 (4)	0.043 (4)	0.038 (3)	-0.016 (3)	-0.010 (3)	-0.003 (3)
C14	0.075 (6)	0.073 (6)	0.059 (5)	-0.011 (5)	-0.019 (5)	0.023 (5)
C15	0.110 (10)	0.107 (10)	0.058 (6)	-0.033 (8)	-0.015 (6)	0.033 (6)
C16	0.101 (9)	0.094 (8)	0.049 (5)	-0.038 (7)	0.010 (5)	-0.009 (5)
C17	0.069 (7)	0.115 (10)	0.065 (6)	-0.025 (7)	0.016 (5)	-0.024 (7)
C18	0.058 (5)	0.068 (6)	0.053 (5)	-0.009 (4)	-0.009 (4)	-0.005 (4)
C19	0.036 (3)	0.030 (3)	0.050 (4)	-0.007 (3)	-0.016 (3)	-0.008 (3)

C20	0.054 (4)	0.046 (4)	0.060 (5)	-0.002 (3)	-0.025 (4)	-0.012 (3)
C21	0.051 (5)	0.048 (5)	0.087 (6)	0.011 (4)	-0.031 (4)	-0.024 (4)
C22	0.040 (4)	0.051 (5)	0.081 (6)	-0.003 (3)	-0.013 (4)	-0.023 (4)
C23	0.054 (4)	0.058 (5)	0.048 (4)	-0.018 (4)	-0.009 (3)	-0.014 (3)
C24	0.052 (4)	0.038 (4)	0.055 (4)	-0.002 (3)	-0.019 (3)	-0.010 (3)

*Geometric parameters (Å, °)*

Sn1—O1 <sup>i</sup>	2.068 (4)	C9—H9	0.9300
Sn1—O1	2.095 (4)	C10—C11	1.345 (16)
Sn1—C1	2.117 (6)	C10—H10	0.9300
Sn1—C7	2.120 (6)	C11—C12	1.396 (12)
Sn1—O2	2.171 (4)	C11—H11	0.9300
Sn2—O1 <sup>i</sup>	2.029 (4)	C12—H12	0.9300
Sn2—C13	2.118 (7)	C13—C14	1.365 (11)
Sn2—C19	2.126 (7)	C13—C18	1.377 (12)
Sn2—O2	2.167 (4)	C14—C15	1.385 (14)
Sn2—Br1	2.6444 (9)	C14—H14	0.9300
O1—Sn2 <sup>i</sup>	2.029 (4)	C15—C16	1.355 (18)
O1—Sn1 <sup>i</sup>	2.068 (4)	C15—H15	0.9300
O2—H2O	0.8200	C16—C17	1.356 (19)
C1—C6	1.369 (11)	C16—H16	0.9300
C1—C2	1.384 (11)	C17—C18	1.397 (13)
C2—C3	1.384 (12)	C17—H17	0.9300
C2—H2	0.9300	C18—H18	0.9300
C3—C4	1.365 (19)	C19—C20	1.383 (10)
C3—H3	0.9300	C19—C24	1.394 (11)
C4—C5	1.37 (2)	C20—C21	1.393 (11)
C4—H4	0.9300	C20—H20	0.9300
C5—C6	1.395 (13)	C21—C22	1.355 (13)
C5—H5	0.9300	C21—H21	0.9300
C6—H6	0.9300	C22—C23	1.381 (12)
C7—C12	1.384 (10)	C22—H22	0.9300
C7—C8	1.373 (10)	C23—C24	1.385 (11)
C8—C9	1.407 (12)	C23—H23	0.9300
C8—H8	0.9300	C24—H24	0.9300
C9—C10	1.360 (16)		
O1 <sup>i</sup> —Sn1—O1	73.05 (18)	C9—C8—H8	119.9
O1 <sup>i</sup> —Sn1—C1	113.5 (2)	C10—C9—C8	118.7 (9)
O1—Sn1—C1	98.9 (2)	C10—C9—H9	120.6
O1 <sup>i</sup> —Sn1—C7	109.8 (2)	C8—C9—H9	120.6
O1—Sn1—C7	98.4 (2)	C11—C10—C9	122.1 (9)
C1—Sn1—C7	136.4 (3)	C11—C10—H10	118.9
O1 <sup>i</sup> —Sn1—O2	72.57 (17)	C9—C10—H10	118.9
O1—Sn1—O2	145.58 (17)	C10—C11—C12	119.7 (10)
C1—Sn1—O2	93.1 (2)	C10—C11—H11	120.2
C7—Sn1—O2	94.7 (2)	C12—C11—H11	120.2

O1 <sup>i</sup> —Sn2—C13	116.9 (2)	C7—C12—C11	119.8 (8)
O1 <sup>i</sup> —Sn2—C19	113.2 (2)	C7—C12—H12	120.1
C13—Sn2—C19	129.4 (3)	C11—C12—H12	120.1
O1 <sup>i</sup> —Sn2—O2	73.38 (17)	C14—C13—C18	118.6 (8)
C13—Sn2—O2	93.3 (2)	C14—C13—Sn2	120.1 (7)
C19—Sn2—O2	94.0 (2)	C18—C13—Sn2	121.3 (6)
O1 <sup>i</sup> —Sn2—Br1	86.33 (12)	C13—C14—C15	120.8 (10)
C13—Sn2—Br1	95.6 (2)	C13—C14—H14	119.6
C19—Sn2—Br1	94.25 (18)	C15—C14—H14	119.6
O2—Sn2—Br1	159.71 (12)	C16—C15—C14	120.1 (11)
Sn2 <sup>i</sup> —O1—Sn1 <sup>i</sup>	111.45 (19)	C16—C15—H15	120.0
Sn2 <sup>i</sup> —O1—Sn1	141.6 (2)	C14—C15—H15	120.0
Sn1 <sup>i</sup> —O1—Sn1	106.95 (18)	C17—C16—C15	120.3 (10)
Sn2—O2—Sn1	102.60 (18)	C17—C16—H16	119.9
Sn2—O2—H2O	128.7	C15—C16—H16	119.9
Sn1—O2—H2O	128.7	C16—C17—C18	119.8 (12)
C6—C1—C2	119.3 (7)	C16—C17—H17	120.1
C6—C1—Sn1	121.1 (6)	C18—C17—H17	120.1
C2—C1—Sn1	119.5 (6)	C13—C18—C17	120.2 (10)
C3—C2—C1	121.1 (9)	C13—C18—H18	119.9
C3—C2—H2	119.5	C17—C18—H18	119.9
C1—C2—H2	119.5	C20—C19—C24	119.3 (7)
C4—C3—C2	119.0 (11)	C20—C19—Sn2	121.2 (5)
C4—C3—H3	120.5	C24—C19—Sn2	119.4 (5)
C2—C3—H3	120.5	C19—C20—C21	119.6 (8)
C3—C4—C5	120.9 (10)	C19—C20—H20	120.2
C3—C4—H4	119.6	C21—C20—H20	120.2
C5—C4—H4	119.6	C22—C21—C20	121.0 (8)
C4—C5—C6	119.7 (11)	C22—C21—H21	119.5
C4—C5—H5	120.1	C20—C21—H21	119.5
C6—C5—H5	120.1	C21—C22—C23	119.9 (8)
C1—C6—C5	119.9 (10)	C21—C22—H22	120.0
C1—C6—H6	120.0	C23—C22—H22	120.0
C5—C6—H6	120.0	C24—C23—C22	120.1 (8)
C12—C7—C8	119.5 (7)	C24—C23—H23	119.9
C12—C7—Sn1	118.4 (5)	C22—C23—H23	119.9
C8—C7—Sn1	121.8 (5)	C23—C24—C19	120.0 (7)
C7—C8—C9	120.1 (9)	C23—C24—H24	120.0
C7—C8—H8	119.9	C19—C24—H24	120.0
O1 <sup>i</sup> —Sn1—O1—Sn2 <sup>i</sup>	-177.8 (5)	C12—C7—C8—C9	0.2 (12)
C1—Sn1—O1—Sn2 <sup>i</sup>	70.2 (4)	Sn1—C7—C8—C9	173.6 (7)
C7—Sn1—O1—Sn2 <sup>i</sup>	-69.5 (4)	C7—C8—C9—C10	-1.9 (15)
O2—Sn1—O1—Sn2 <sup>i</sup>	179.3 (3)	C8—C9—C10—C11	3.2 (19)
O1 <sup>i</sup> —Sn1—O1—Sn1 <sup>i</sup>	0.0	C9—C10—C11—C12	-3 (2)
C1—Sn1—O1—Sn1 <sup>i</sup>	-112.0 (2)	C8—C7—C12—C11	0.2 (13)
C7—Sn1—O1—Sn1 <sup>i</sup>	108.2 (2)	Sn1—C7—C12—C11	-173.4 (8)
O2—Sn1—O1—Sn1 <sup>i</sup>	-2.9 (4)	C10—C11—C12—C7	1.0 (17)

O1 <sup>i</sup> —Sn2—O2—Sn1	-0.23 (16)	O1 <sup>i</sup> —Sn2—C13—C14	-143.7 (8)
C13—Sn2—O2—Sn1	116.9 (3)	C19—Sn2—C13—C14	45.3 (9)
C19—Sn2—O2—Sn1	-113.2 (2)	O2—Sn2—C13—C14	143.2 (8)
Br1—Sn2—O2—Sn1	0.7 (5)	Br1—Sn2—C13—C14	-55.0 (8)
O1 <sup>i</sup> —Sn1—O2—Sn2	0.23 (16)	O1 <sup>i</sup> —Sn2—C13—C18	37.9 (8)
O1—Sn1—O2—Sn2	3.2 (4)	C19—Sn2—C13—C18	-133.2 (7)
C1—Sn1—O2—Sn2	113.9 (3)	O2—Sn2—C13—C18	-35.2 (7)
C7—Sn1—O2—Sn2	-109.1 (2)	Br1—Sn2—C13—C18	126.6 (7)
O1 <sup>i</sup> —Sn1—C1—C6	112.1 (6)	C18—C13—C14—C15	4.3 (17)
O1—Sn1—C1—C6	-172.6 (6)	Sn2—C13—C14—C15	-174.2 (11)
C7—Sn1—C1—C6	-60.4 (8)	C13—C14—C15—C16	-2 (2)
O2—Sn1—C1—C6	39.7 (7)	C14—C15—C16—C17	-2 (2)
O1 <sup>i</sup> —Sn1—C1—C2	-64.6 (7)	C15—C16—C17—C18	3 (2)
O1—Sn1—C1—C2	10.7 (7)	C14—C13—C18—C17	-3.3 (16)
C7—Sn1—C1—C2	122.9 (7)	Sn2—C13—C18—C17	175.2 (9)
O2—Sn1—C1—C2	-137.0 (7)	C16—C17—C18—C13	-0.3 (19)
C6—C1—C2—C3	-1.1 (15)	O1 <sup>i</sup> —Sn2—C19—C20	123.9 (6)
Sn1—C1—C2—C3	175.6 (9)	C13—Sn2—C19—C20	-64.8 (7)
C1—C2—C3—C4	3.3 (19)	O2—Sn2—C19—C20	-162.4 (6)
C2—C3—C4—C5	-2 (2)	Br1—Sn2—C19—C20	36.1 (6)
C3—C4—C5—C6	-1 (2)	O1 <sup>i</sup> —Sn2—C19—C24	-50.8 (6)
C2—C1—C6—C5	-2.4 (13)	C13—Sn2—C19—C24	120.5 (6)
Sn1—C1—C6—C5	-179.1 (8)	O2—Sn2—C19—C24	22.9 (6)
C4—C5—C6—C1	3.7 (17)	Br1—Sn2—C19—C24	-138.6 (6)
O1 <sup>i</sup> —Sn1—C7—C12	63.6 (6)	C24—C19—C20—C21	0.7 (12)
O1—Sn1—C7—C12	-11.3 (6)	Sn2—C19—C20—C21	-174.0 (6)
C1—Sn1—C7—C12	-123.7 (6)	C19—C20—C21—C22	1.4 (14)
O2—Sn1—C7—C12	136.8 (6)	C20—C21—C22—C23	-1.7 (14)
O1 <sup>i</sup> —Sn1—C7—C8	-109.8 (6)	C21—C22—C23—C24	-0.2 (13)
O1—Sn1—C7—C8	175.3 (6)	C22—C23—C24—C19	2.3 (12)
C1—Sn1—C7—C8	62.9 (7)	C20—C19—C24—C23	-2.5 (12)
O2—Sn1—C7—C8	-36.7 (6)	Sn2—C19—C24—C23	172.3 (6)

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .