

Dichlorido(1,10-phenanthroline- $\kappa^2 N,N'$)-palladium(II)

Kwang Ha

School of Applied Chemical Engineering, The Research Institute of Catalysis, Chonnam National University, Gwangju 500-757, Republic of Korea
Correspondence e-mail: hakwang@chonnam.ac.kr


Received 1 December 2009; accepted 5 December 2009

Key indicators: single-crystal X-ray study; $T = 200\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.011\text{ \AA}$; R factor = 0.056; wR factor = 0.169; data-to-parameter ratio = 18.0.

In the title complex, $[\text{PdCl}_2(\text{C}_{12}\text{H}_8\text{N}_2)]$, the Pd^{2+} ion is four-coordinated in a slightly distorted square-planar environment by two N atoms of the chelating 1,10-phenanthroline ligand and two chloride ions. The nearly planar molecules, with a maximum deviation of $0.120(3)\text{ \AA}$ from the least-squares plane, are stacked in columns along the c axis with a $\text{Pd}\cdots\text{Pd}$ distance of $4.8340(9)\text{ \AA}$. In the column, $\pi\cdots\pi$ interactions between adjacent six-membered rings are present, the shortest centroid–centroid distance being $3.680(4)\text{ \AA}$. A weak $\text{C}\cdots\text{H}\cdots\text{Cl}$ interaction is observed between the columns.

Related literature

For the syntheses of $[\text{Pd}X_2(\text{phen})]$ ($\text{phen} = 1,10\text{-phenanthroline}; X = \text{Cl}, \text{Br}, \text{I} \text{ or } \text{SCN}$), see: Cheng *et al.* (1977). For the crystal structure of yellow $[\text{PtCl}_2(\text{phen})]$ which is isotypic to the title complex, see: Grzesiak & Matzger (2007). For the crystal structures of related Pd-bipy complexes, $[\text{Pd}X_2(\text{bipy})]$ ($\text{bipy} = 2,2'\text{-bipyridine}; X = \text{Cl}, \text{Br} \text{ or } \text{I}$), see: Maekawa *et al.* (1991); Smeets *et al.* (1997); Ha (2009).



Experimental

Crystal data

$[\text{PdCl}_2(\text{C}_{12}\text{H}_8\text{N}_2)]$	$V = 1128.26(16)\text{ \AA}^3$
$M_r = 357.50$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 9.6170(8)\text{ \AA}$	$\mu = 2.09\text{ mm}^{-1}$
$b = 17.1402(14)\text{ \AA}$	$T = 200\text{ K}$
$c = 7.2529(6)\text{ \AA}$	$0.36 \times 0.06 \times 0.04\text{ mm}$
$\beta = 109.314(2)^\circ$	

Data collection

Bruker SMART 1000 CCD diffractometer	8118 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2000)	2767 independent reflections
$T_{\min} = 0.769$, $T_{\max} = 0.920$	1852 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.060$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$	154 parameters
$wR(F^2) = 0.169$	H-atom parameters constrained
$S = 1.10$	$\Delta\rho_{\max} = 2.90\text{ e \AA}^{-3}$
2767 reflections	$\Delta\rho_{\min} = -1.29\text{ e \AA}^{-3}$

Table 1
Selected geometric parameters (\AA , $^\circ$).

Pd1—N2	2.035 (6)	Pd1—Cl2	2.283 (2)
Pd1—N1	2.036 (6)	Pd1—Cl1	2.2914 (19)
N2—Pd1—N1	81.5 (2)		

Table 2

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C2—H2 \cdots Cl1 ⁱ	0.95	2.80	3.733 (11)	169

Symmetry code: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Priority Research Centers Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2009-0094056).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2500).

References

- Bruker (2000). *SADABS, SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cheng, C. P., Plankey, B., Rund, J. V. & Brown, T. L. (1977). *J. Am. Chem. Soc.* **99**, 8413–8417.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Grzesiak, A. L. & Matzger, A. J. (2007). *Inorg. Chem.* **46**, 453–457.
- Ha, K. (2009). *Acta Cryst. E65*, m1588.
- Maekawa, M., Munakata, M., Kitagawa, S. & Nakamura, M. (1991). *Anal. Sci.* **7**, 521–522.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Smeets, W. J. J., Spek, A. L., Hoare, J. L., Carty, A. J., Hovestad, N. & van Koten, G. (1997). *Acta Cryst. C53*, 1045–1047.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supporting information

Acta Cryst. (2010). E66, m38 [doi:10.1107/S1600536809052313]

Dichlorido(1,10-phenanthroline- κ^2N,N')palladium(II)

Kwang Ha

S1. Comment

The title complex, $[PdCl_2(\text{phen})]$ (where phen is 1,10-phenanthroline, $C_{12}H_8N_2$), is isomorphous with the yellow form of $[PtCl_2(\text{phen})]$, whereas the orange form of $[PtCl_2(\text{phen})]$ crystallized in the orthorhombic space group $Pca2_1$ (Grzesiak & Matzger, 2007).


In the title complex, the Pd^{2+} ion is four-coordinated in a slightly distorted square-planar environment by two N atoms of the chelating 1,10-phenanthroline ligand and two chloride ions (Fig. 1). The main contribution to the distortion is the tight $N1—Pd1—N2$ chelate angle [$81.5(2)^\circ$], which results in non-linear *trans* arrangement [$<N1—Pd1—Cl1 = 174.64(17)^\circ$ and $<N2—Pd1—Cl2 = 175.11(17)^\circ$]. The $Pd1—N$ and $Pd1—Cl$ bond lengths are almost equal, respectively [$Pd1—N: 2.036(6)$ and $2.035(6)$ Å; $Pd1—Cl: 2.2914(19)$ and $2.283(2)$ Å]. The complex displays numerous intermolecular $\pi\cdots\pi$ interactions between adjacent six-membered rings, with a shortest centroid-centroid distance of 3.680(4) Å and the dihedral angle between the ring planes is $5.6(4)^\circ$. The nearly planar $[PdCl_2(\text{phen})]$ molecules with a largest deviation of 0.120(3) Å from the least-squares plane stack in columns along the c axis with a $Pd\cdots Pd$ distance of 4.8340(9) Å (Fig. 2).

S2. Experimental


To a solution of Na_2PdCl_4 (0.5095 g, 1.732 mmol) in H_2O (40 ml) was added 1,10-phenanthroline (0.3122 g, 1.732 mmol) and refluxed for 3 h. The precipitate obtained was separated by filtration, washed with water and acetone, and dried at 70 °C, to give a pale yellow powder (0.5259 g). Crystals suitable for X-ray analysis were obtained by slow evaporation from a dimethyl sulfoxide solution at 80 °C.

S3. Refinement

H atoms were positioned geometrically and allowed to ride on their respective parent atoms [$C—H = 0.95$ Å and $U_{iso}(H) = 1.2U_{eq}(C)$]. The highest peak and the deepest hole in the difference Fourier map are located 1.14 and 0.85 Å, respectively, from atom Pd1.

**Figure 1**

The structure of the title complex, with displacement ellipsoids drawn at the 50% probability level for non-H atoms.

**Figure 2**

Crystal packing diagram of the title complex.

Dichlorido(1,10-phenanthroline- κ^2N,N')palladium(II)

Crystal data

[PdCl₂(C₁₂H₈N₂)]

$M_r = 357.50$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 9.6170 (8) \text{ \AA}$

$b = 17.1402 (14) \text{ \AA}$

$c = 7.2529 (6) \text{ \AA}$

$\beta = 109.314 (2)^\circ$

$V = 1128.26 (16) \text{ \AA}^3$

$Z = 4$

$F(000) = 696$

$D_x = 2.105 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2653 reflections

$\theta = 2.4\text{--}28.2^\circ$

$\mu = 2.09 \text{ mm}^{-1}$

$T = 200 \text{ K}$

Needle, yellow

$0.36 \times 0.06 \times 0.04 \text{ mm}$

Data collection

Bruker SMART 1000 CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2000)
 $T_{\min} = 0.769$, $T_{\max} = 0.920$

8118 measured reflections
2767 independent reflections
1852 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.060$
 $\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -12 \rightarrow 12$
 $k = -22 \rightarrow 19$
 $l = -9 \rightarrow 9$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.169$
 $S = 1.10$
2767 reflections
154 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.084P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 2.90 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -1.29 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Pd1	0.70739 (6)	0.34324 (3)	0.39119 (8)	0.0214 (2)
Cl1	0.7453 (2)	0.47533 (11)	0.4189 (3)	0.0313 (5)
Cl2	0.4658 (2)	0.36052 (12)	0.2078 (3)	0.0366 (5)
N1	0.6941 (6)	0.2247 (3)	0.3805 (9)	0.0235 (13)
N2	0.9182 (6)	0.3184 (3)	0.5594 (9)	0.0211 (13)
C1	0.5819 (9)	0.1794 (5)	0.2838 (12)	0.0319 (18)
H1	0.4931	0.2030	0.2036	0.038*
C2	0.5898 (10)	0.0983 (5)	0.2958 (13)	0.039 (2)
H2	0.5073	0.0676	0.2244	0.046*
C3	0.7163 (9)	0.0629 (4)	0.4102 (12)	0.033 (2)
H3	0.7214	0.0076	0.4202	0.040*
C4	0.8382 (9)	0.1085 (4)	0.5127 (11)	0.0286 (17)
C5	0.9777 (10)	0.0785 (4)	0.6369 (12)	0.0330 (19)
H5	0.9904	0.0238	0.6561	0.040*
C6	1.0895 (9)	0.1263 (5)	0.7254 (12)	0.0318 (18)
H6	1.1805	0.1045	0.8044	0.038*

C7	1.0771 (8)	0.2089 (4)	0.7057 (11)	0.0263 (16)
C8	1.1907 (8)	0.2626 (5)	0.7913 (12)	0.0301 (17)
H8	1.2850	0.2445	0.8702	0.036*
C9	1.1660 (8)	0.3405 (4)	0.7613 (12)	0.0287 (17)
H9	1.2423	0.3768	0.8210	0.034*
C10	1.0283 (8)	0.3669 (4)	0.6426 (11)	0.0269 (16)
H10	1.0131	0.4213	0.6207	0.032*
C11	0.9420 (8)	0.2407 (4)	0.5896 (10)	0.0234 (15)
C12	0.8212 (8)	0.1902 (4)	0.4927 (11)	0.0224 (15)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pd1	0.0171 (3)	0.0198 (3)	0.0258 (3)	-0.0001 (2)	0.0050 (2)	0.0001 (2)
C11	0.0285 (10)	0.0198 (9)	0.0422 (12)	0.0034 (7)	0.0070 (9)	0.0016 (7)
Cl2	0.0178 (9)	0.0379 (11)	0.0457 (13)	0.0017 (8)	-0.0007 (8)	0.0059 (9)
N1	0.022 (3)	0.016 (3)	0.031 (3)	-0.003 (2)	0.006 (3)	-0.004 (2)
N2	0.018 (3)	0.014 (3)	0.030 (3)	0.000 (2)	0.006 (3)	0.000 (2)
C1	0.026 (4)	0.037 (4)	0.031 (4)	-0.005 (3)	0.007 (4)	-0.005 (3)
C2	0.040 (5)	0.029 (5)	0.048 (5)	-0.017 (4)	0.016 (4)	-0.015 (4)
C3	0.043 (5)	0.011 (4)	0.053 (6)	-0.005 (3)	0.025 (5)	-0.004 (3)
C4	0.038 (5)	0.022 (4)	0.030 (4)	-0.005 (3)	0.018 (4)	0.000 (3)
C5	0.054 (6)	0.021 (4)	0.029 (4)	0.005 (4)	0.021 (4)	0.006 (3)
C6	0.039 (5)	0.026 (4)	0.035 (5)	0.013 (4)	0.019 (4)	0.008 (3)
C7	0.023 (4)	0.026 (4)	0.034 (4)	0.011 (3)	0.015 (3)	0.007 (3)
C8	0.016 (4)	0.039 (5)	0.033 (5)	0.003 (3)	0.006 (3)	0.005 (3)
C9	0.020 (4)	0.032 (4)	0.029 (4)	-0.001 (3)	0.001 (3)	-0.003 (3)
C10	0.029 (4)	0.021 (4)	0.031 (4)	-0.003 (3)	0.009 (3)	-0.004 (3)
C11	0.022 (4)	0.026 (4)	0.023 (4)	0.000 (3)	0.009 (3)	-0.003 (3)
C12	0.032 (4)	0.013 (3)	0.027 (4)	-0.003 (3)	0.016 (3)	-0.005 (3)

Geometric parameters (\AA , $^\circ$)

Pd1—N2	2.035 (6)	C4—C12	1.412 (10)
Pd1—N1	2.036 (6)	C4—C5	1.441 (12)
Pd1—Cl2	2.283 (2)	C5—C6	1.335 (11)
Pd1—Cl1	2.2914 (19)	C5—H5	0.9500
N1—C1	1.327 (9)	C6—C7	1.423 (11)
N1—C12	1.359 (9)	C6—H6	0.9500
N2—C10	1.324 (9)	C7—C11	1.404 (10)
N2—C11	1.357 (9)	C7—C8	1.407 (10)
C1—C2	1.392 (12)	C8—C9	1.360 (10)
C1—H1	0.9500	C8—H8	0.9500
C2—C3	1.368 (12)	C9—C10	1.394 (11)
C2—H2	0.9500	C9—H9	0.9500
C3—C4	1.401 (11)	C10—H10	0.9500
C3—H3	0.9500	C11—C12	1.433 (10)

N2—Pd1—N1	81.5 (2)	C6—C5—C4	121.1 (7)
N2—Pd1—Cl2	175.11 (17)	C6—C5—H5	119.5
N1—Pd1—Cl2	93.93 (17)	C4—C5—H5	119.5
N2—Pd1—Cl1	93.20 (17)	C5—C6—C7	122.3 (7)
N1—Pd1—Cl1	174.64 (17)	C5—C6—H6	118.8
Cl2—Pd1—Cl1	91.42 (7)	C7—C6—H6	118.8
C1—N1—C12	118.4 (7)	C11—C7—C8	116.1 (7)
C1—N1—Pd1	129.4 (5)	C11—C7—C6	118.5 (7)
C12—N1—Pd1	112.2 (4)	C8—C7—C6	125.4 (7)
C10—N2—C11	118.4 (6)	C9—C8—C7	120.2 (7)
C10—N2—Pd1	129.1 (5)	C9—C8—H8	119.9
C11—N2—Pd1	112.6 (5)	C7—C8—H8	119.9
N1—C1—C2	122.2 (8)	C8—C9—C10	119.8 (7)
N1—C1—H1	118.9	C8—C9—H9	120.1
C2—C1—H1	118.9	C10—C9—H9	120.1
C3—C2—C1	120.0 (7)	N2—C10—C9	122.1 (7)
C3—C2—H2	120.0	N2—C10—H10	119.0
C1—C2—H2	120.0	C9—C10—H10	119.0
C2—C3—C4	119.6 (7)	N2—C11—C7	123.5 (7)
C2—C3—H3	120.2	N2—C11—C12	116.6 (6)
C4—C3—H3	120.2	C7—C11—C12	119.9 (7)
C3—C4—C12	116.8 (7)	N1—C12—C4	122.9 (6)
C3—C4—C5	125.1 (7)	N1—C12—C11	117.0 (6)
C12—C4—C5	118.1 (7)	C4—C12—C11	120.1 (7)
N2—Pd1—N1—C1	-177.2 (7)	Pd1—N2—C10—C9	177.3 (5)
Cl2—Pd1—N1—C1	4.4 (7)	C8—C9—C10—N2	1.3 (12)
N2—Pd1—N1—C12	3.4 (5)	C10—N2—C11—C7	0.5 (11)
Cl2—Pd1—N1—C12	-174.9 (5)	Pd1—N2—C11—C7	-178.0 (6)
N1—Pd1—N2—C10	178.1 (7)	C10—N2—C11—C12	-178.3 (6)
Cl1—Pd1—N2—C10	-1.4 (6)	Pd1—N2—C11—C12	3.1 (8)
N1—Pd1—N2—C11	-3.6 (5)	C8—C7—C11—N2	-0.4 (11)
Cl1—Pd1—N2—C11	176.9 (5)	C6—C7—C11—N2	-179.5 (7)
C12—N1—C1—C2	0.8 (11)	C8—C7—C11—C12	178.4 (7)
Pd1—N1—C1—C2	-178.5 (6)	C6—C7—C11—C12	-0.7 (10)
N1—C1—C2—C3	0.1 (13)	C1—N1—C12—C4	-0.9 (10)
C1—C2—C3—C4	-1.0 (12)	Pd1—N1—C12—C4	178.6 (5)
C2—C3—C4—C12	0.9 (11)	C1—N1—C12—C11	177.8 (6)
C2—C3—C4—C5	-179.6 (8)	Pd1—N1—C12—C11	-2.8 (8)
C3—C4—C5—C6	178.3 (7)	C3—C4—C12—N1	0.0 (11)
C12—C4—C5—C6	-2.2 (11)	C5—C4—C12—N1	-179.5 (7)
C4—C5—C6—C7	1.1 (12)	C3—C4—C12—C11	-178.6 (7)
C5—C6—C7—C11	0.3 (11)	C5—C4—C12—C11	1.8 (10)
C5—C6—C7—C8	-178.7 (8)	N2—C11—C12—N1	-0.2 (10)
C11—C7—C8—C9	0.7 (11)	C7—C11—C12—N1	-179.2 (6)
C6—C7—C8—C9	179.7 (7)	N2—C11—C12—C4	178.5 (6)
C7—C8—C9—C10	-1.1 (12)	C7—C11—C12—C4	-0.5 (10)
C11—N2—C10—C9	-1.0 (11)		

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
C2—H2···Cl1 ⁱ	0.95	2.80	3.733 (11)	169

Symmetry code: (i) $-x+1, y-1/2, -z+1/2$.