

N,N'-Dicyclohexylethylenediammonium dichloride

Bob-Dan Lechner and Kurt Merzweiler*

Institut für Chemie, Naturwissenschaftliche Fakultät II, Martin-Luther-Universität Halle-Wittenberg, Kurt-Mothes-Str. 2, 06120 Halle, Germany
Correspondence e-mail: kurt.merzweiler@chemie.uni-halle.de


Received 4 November 2009; accepted 4 December 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.035; wR factor = 0.089; data-to-parameter ratio = 11.8.

In the title compound, $\text{C}_{14}\text{H}_{30}\text{N}_2^{2+}\cdot 2\text{Cl}^-$, the N,N' -dicyclohexylethylenediammonium cation possesses crystallographic $\bar{1}$ symmetry, and thus the compound crystallizes with two formula units per unit cell. In the crystal, the cations and anions are linked by $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds, giving a two-dimensional network with {6,3} topology.

Related literature

For the crystal structures of cyclohexylammonium derivatives, see Smith *et al.* (1994); Martell & Zaworotko (1991). For the crystal structure of an iridium complex with the N,N' -dicyclohexylethylenediamine ligand, see: Greulich *et al.* (2002). For a review of hydrogen bonding, see Steiner (2002). N,N' -dicyclohexylethylenediamine was prepared according to Denk *et al.* (2003). For the topology of {6,3} ring systems and three-dimensional polyhedra and networks, see: Wells & Sharpe (1963).



Experimental

Crystal data



$M_r = 297.30$

Monoclinic, $P2_1/c$
 $a = 11.551(3)\text{ \AA}$
 $b = 6.785(2)\text{ \AA}$
 $c = 10.8434(17)\text{ \AA}$
 $\beta = 91.892(15)^\circ$
 $V = 849.3(4)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.37\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.65 \times 0.28 \times 0.12\text{ mm}$

Data collection

Stoe STADI4 diffractometer
3349 measured reflections
1675 independent reflections
1430 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$
2 standard reflections every 120 min
intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.089$
 $S = 1.10$
1675 reflections
142 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.23\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.28\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}-\text{H}4\cdots\text{Cl}$	0.91 (2)	2.20 (2)	3.1088 (16)	175.8 (18)
$\text{N}-\text{H}3\cdots\text{Cl}^1$	0.84 (2)	2.30 (2)	3.1250 (16)	168.8 (18)

Symmetry code: (i) $-x, y - \frac{1}{2}, -z + \frac{3}{2}$

Data collection: STADI4 (Stoe & Cie, 1996); cell refinement: STADI4; data reduction: X-RED (Stoe & Cie, 1996); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg, 2009); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SI2221).

References

- Brandenburg, K. (2009). DIAMOND. Crystal Impact GbR, Bonn, Germany.
- Denk, M. K., Krause, M. J., Niyogi, D. F. & Gill, N. K. (2003). *Tetrahedron*, **59**, 7565–7570.
- Greulich, S., Klein, A., Knödler, A. & Kaim, W. (2002). *Organometallics*, **21**, 765–769.
- Martell, J. M. & Zaworotko, M. (1991). *J. Chem. Soc. Dalton Trans.* pp. 1495–1498.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Smith, H. W., Mastropaoletti, D., Camerman, A. & Camerman, N. (1994). *J. Chem. Crystallogr.* **24**, 239–242.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Steiner, T. (2002). *Angew. Chem. Int. Ed.* **41**, 48–76.
- Stoe & Cie. (1996). STADI4 and X-RED. Stoe & Cie, Darmstadt, Germany.
- Wells, A. F. & Sharpe, R. R. (1963). *Acta Cryst.* **16**, 857–871.

supporting information

Acta Cryst. (2010). E66, o122 [doi:10.1107/S1600536809052222]

N,N'-Dicyclohexylethylenediammonium dichloride

Bob-Dan Lechner and Kurt Merzweiler

S1. Comment


Compared to other *N,N'*-disubstituted ethylenediamine compounds RNH-CH₂CH₂-NHR (*R* = Me, Ph, etc.) *N,N'*-Di-cyclohexylethylenediamine derivatives have been studied only rarely by X-ray diffraction. One of the few examples is the Iridium complex [Cp*(CyNH₂CH₂CH₂NHCy)HIr][H₃BCN] (Greulich *et al.*, 2002).

The crystal structure of the title compound (Fig. 1) consists of [CyNH₂CH₂CH₂NH₂Cy]²⁺ cations and Cl⁻ anions. The [CyNH₂CH₂CH₂NH₂Cy]²⁺ cations exhibit crystallographic $\bar{1}$ symmetry and thus an exactly staggered conformation with a N—C—C—N torsion angle of 180 ° is observed. The N atoms display a distorted tetrahedral coordination with H—N—C angles of 108.6 to 111.2 and a C—N—C angle of 114.50 (1)°. The cyclohexyl groups adopt a chair conformation (Fig. 2) as it was observed in [Cp*(CyNH₂CH₂CH₂NHCy)HIr][H₃BCN] (Greulich *et al.*, 2002). Both hydrogen atoms of the NH₂ groups are involved in hydrogen bridges to neighbouring Cl⁻ anions. The NH···Cl distances of 2.20 (2) and 2.30 (2) Å are comparable to those found in other cyclohexylammonium derivatives like [CyNH₃]Cl (2.187–2.354 Å) (Smith *et al.*, 1994) and [CyNH₃]₂(AlCl₄)Cl (2.305–2.478 Å) (Martell & Zaworotko, 1991) respectively. The N—H···Cl angles of 169 (2)° and 176 (2)° are in the expected range for hydrogen bridges of moderate strength (Steiner, 2002).


On balance each [CyNH₂CH₂CH₂NH₂Cy]²⁺ cation forms four N—H···Cl bridges to neighbouring Cl⁻ anions and each Cl⁻ anion acts as H-acceptor for two NH hydrogen atoms. As a result of the hydrogen bonding between [CyNH₂CH₂CH₂NH₂Cy]²⁺ cations and Cl⁻ anions a two-dimensional layer structure is formed. The layers consist of puckered C₄H₈N₆Cl₄ rings that are interconnected to give a honeycomb arrangement with {6,3} net topology (Wells & Sharpe, 1963).

S2. Experimental

An excess of hydrochloric acid was added dropwise to a solution of *N,N'*-Dicyclohexylethylenediamine monohydrate (1.11 g, 5 mmol) prepared by standard techniques (Denk *et al.*, 2003) in a ethanol/water mixture(10:1, 20 ml) and stirred for 6 h at 140 °C in an autoclave. The mixture was slowly cooled to ambient temperature and colourless plate-like crystals were obtained. Spectroscopic data: ¹H NMR (D₂O, 500 MHz, 298 K, p.p.m.): δ 1.07–1.99 (m, 20 H, CH₂, Cy), 3.09 (m, 2H, CH), 3.32 (s, 4H, CH₂); ¹³C NMR (D₂O, 125 MHz, 298 K, p.p.m.): δ 23.7 (s, CH₂, Cy: C4, C6), 24.3 (s, CH₂, Cy: C5), 26.7 (s, CH₂, Cy: C3, C7), 40.0 (s, CH₂), 57.934 (s, CH, Cy).

**Figure 1**


Molecular structure of the $[\text{CyNH}_2\text{CH}_2\text{CH}_2\text{NH}_2\text{Cy}]^{2+}$ cation with surrounding Cl^- anions. The asymmetric unit is shown by filled bonds. Thermal ellipsoids are drawn at the 50% probability level. Symmetry codes: (i) $-x, -y, -z$; (ii) $-x, -1/2 + y, 1.5 - z$; (iii) $x, 1.5 - y, -1/2 + z$

**Figure 2**

Part of the layer structure formed by hydrogen bonded $[\text{CyNH}_2\text{CH}_2\text{CH}_2\text{NH}_2\text{Cy}]^{2+}$ cations and Cl^- anions. Cyclohexyl groups are omitted for clarity.

N,N'-Dicyclohexylenediammonium dichloride

Crystal data



$M_r = 297.30$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 11.551(3)$ Å

$b = 6.785(2)$ Å

$c = 10.8434(17)$ Å

$\beta = 91.892(15)^\circ$

$V = 849.3(4)$ Å³

$Z = 2$

Data collection

Stoe STADI4

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$2\theta/\omega$ scans

3349 measured reflections

1675 independent reflections

1430 reflections with $I > 2\sigma(I)$

$F(000) = 324$

$D_x = 1.163 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 22 reflections

$\theta = 10.2\text{--}14.6^\circ$

$\mu = 0.37 \text{ mm}^{-1}$

$T = 293$ K

Plate, colourless

$0.65 \times 0.28 \times 0.12$ mm

$R_{\text{int}} = 0.036$

$\theta_{\text{max}} = 26.1^\circ, \theta_{\text{min}} = 1.8^\circ$

$h = -14 \rightarrow 14$

$k = -8 \rightarrow 0$

$l = -13 \rightarrow 13$

2 standard reflections every 120 min

intensity decay: none

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.035$$

$$wR(F^2) = 0.089$$

$$S = 1.10$$

1675 reflections

142 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0304P)^2 + 0.223P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.05608 (15)	0.4477 (3)	0.48641 (16)	0.0404 (4)
H2	0.1138 (18)	0.537 (3)	0.4631 (19)	0.054 (6)*
H1	0.0449 (18)	0.353 (3)	0.419 (2)	0.059 (6)*
C2	0.21086 (13)	0.2307 (2)	0.58171 (14)	0.0342 (3)
H5	0.2039 (14)	0.163 (3)	0.5053 (16)	0.038 (4)*
C3	0.22533 (16)	0.0777 (3)	0.6834 (2)	0.0478 (5)
H6	0.2248 (18)	0.147 (3)	0.763 (2)	0.059 (6)*
H7	0.1619 (18)	-0.009 (4)	0.6755 (18)	0.058 (6)*
C4	0.33978 (18)	-0.0313 (3)	0.6726 (3)	0.0574 (5)
H8	0.3397 (19)	-0.095 (3)	0.596 (2)	0.063 (7)*
H9	0.346 (2)	-0.123 (4)	0.738 (2)	0.069 (7)*
C5	0.44122 (17)	0.1104 (3)	0.6751 (2)	0.0513 (5)
H10	0.4456 (18)	0.173 (3)	0.757 (2)	0.064 (6)*
H11	0.516 (2)	0.045 (3)	0.668 (2)	0.068 (6)*
C6	0.42621 (16)	0.2660 (3)	0.5755 (2)	0.0522 (5)
H12	0.4254 (18)	0.198 (3)	0.497 (2)	0.064 (6)*
H13	0.492 (2)	0.354 (4)	0.581 (2)	0.075 (7)*
C7	0.31115 (15)	0.3757 (3)	0.58388 (18)	0.0419 (4)
H14	0.3073 (17)	0.447 (3)	0.6637 (19)	0.054 (6)*
H15	0.3005 (17)	0.466 (3)	0.5188 (18)	0.053 (6)*
N	0.09763 (12)	0.3358 (2)	0.59677 (13)	0.0335 (3)
H4	0.1015 (17)	0.416 (3)	0.6647 (19)	0.052 (6)*
H3	0.0454 (17)	0.257 (3)	0.6169 (17)	0.045 (5)*
Cl	0.12073 (4)	0.59342 (7)	0.83342 (4)	0.04938 (17)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0421 (9)	0.0418 (9)	0.0375 (8)	0.0074 (8)	0.0039 (7)	0.0097 (7)
C2	0.0376 (8)	0.0297 (8)	0.0351 (8)	0.0025 (6)	-0.0004 (6)	-0.0027 (6)
C3	0.0413 (10)	0.0349 (9)	0.0670 (13)	-0.0006 (8)	-0.0007 (8)	0.0165 (9)
C4	0.0521 (11)	0.0354 (9)	0.0838 (16)	0.0089 (9)	-0.0074 (10)	0.0073 (11)
C5	0.0405 (10)	0.0492 (11)	0.0638 (12)	0.0098 (8)	-0.0038 (8)	0.0007 (9)
C6	0.0391 (9)	0.0583 (12)	0.0596 (12)	0.0033 (9)	0.0088 (8)	0.0046 (10)
C7	0.0383 (9)	0.0361 (9)	0.0514 (10)	-0.0011 (7)	0.0039 (7)	0.0081 (8)
N	0.0338 (7)	0.0314 (7)	0.0352 (7)	-0.0018 (6)	0.0011 (5)	0.0043 (6)
C1	0.0569 (3)	0.0446 (3)	0.0467 (3)	0.0160 (2)	0.00215 (19)	-0.00631 (19)

Geometric parameters (\AA , $^\circ$)

C1—N	1.484 (2)	C4—H8	0.94 (2)
C1—C1 ⁱ	1.514 (3)	C4—H9	0.95 (2)
C1—H2	0.94 (2)	C5—C6	1.516 (3)
C1—H1	0.98 (2)	C5—H10	0.98 (2)
C2—N	1.503 (2)	C5—H11	0.98 (2)
C2—C7	1.519 (2)	C6—C7	1.529 (2)
C2—C3	1.519 (2)	C6—H12	0.97 (2)
C2—H5	0.948 (18)	C6—H13	0.97 (3)
C3—C4	1.523 (3)	C7—H14	0.99 (2)
C3—H6	0.98 (2)	C7—H15	0.94 (2)
C3—H7	0.94 (2)	N—H4	0.91 (2)
C4—C5	1.515 (3)	N—H3	0.84 (2)
N—C1—C1 ⁱ	109.80 (17)	C4—C5—C6	111.01 (17)
N—C1—H2	109.4 (13)	C4—C5—H10	108.0 (13)
C1 ⁱ —C1—H2	111.7 (12)	C6—C5—H10	109.9 (13)
N—C1—H1	107.3 (13)	C4—C5—H11	113.4 (14)
C1 ⁱ —C1—H1	111.1 (12)	C6—C5—H11	109.8 (13)
H2—C1—H1	107.4 (17)	H10—C5—H11	104.5 (18)
N—C2—C7	110.89 (13)	C5—C6—C7	112.07 (16)
N—C2—C3	108.69 (13)	C5—C6—H12	107.2 (13)
C7—C2—C3	111.44 (15)	C7—C6—H12	107.4 (13)
N—C2—H5	106.0 (10)	C5—C6—H13	108.4 (14)
C7—C2—H5	111.7 (10)	C7—C6—H13	112.4 (14)
C3—C2—H5	107.9 (11)	H12—C6—H13	109.2 (18)
C2—C3—C4	110.54 (17)	C2—C7—C6	110.34 (15)
C2—C3—H6	108.0 (12)	C2—C7—H14	105.8 (12)
C4—C3—H6	109.2 (12)	C6—C7—H14	110.7 (12)
C2—C3—H7	107.1 (13)	C2—C7—H15	109.3 (12)
C4—C3—H7	111.5 (13)	C6—C7—H15	111.5 (12)
H6—C3—H7	110.5 (17)	H14—C7—H15	109.1 (17)
C5—C4—C3	111.30 (17)	C1—N—C2	114.50 (13)
C5—C4—H8	106.8 (14)	C1—N—H4	110.6 (12)

C3—C4—H8	108.5 (14)	C2—N—H4	110.5 (13)
C5—C4—H9	111.3 (14)	C1—N—H3	108.6 (13)
C3—C4—H9	107.9 (14)	C2—N—H3	111.1 (13)
H8—C4—H9	111 (2)	H4—N—H3	100.6 (17)
N—C2—C3—C4	-179.40 (16)	C3—C2—C7—C6	55.7 (2)
C7—C2—C3—C4	-56.9 (2)	C5—C6—C7—C2	-54.8 (2)
C2—C3—C4—C5	56.5 (3)	C1 ⁱ —C1—N—C2	-178.21 (18)
C3—C4—C5—C6	-55.5 (3)	C7—C2—N—C1	73.57 (19)
C4—C5—C6—C7	54.9 (2)	C3—C2—N—C1	-163.61 (15)
N—C2—C7—C6	176.95 (14)		

Symmetry code: (i) $-x, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^{\circ}$)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N—H4 \cdots Cl	0.91 (2)	2.20 (2)	3.1088 (16)	175.8 (18)
N—H3 \cdots Cl ⁱⁱ	0.84 (2)	2.30 (2)	3.1250 (16)	168.8 (18)

Symmetry code: (ii) $-x, y-1/2, -z+3/2$.