

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(2-Naphthyl)-1,3-dioxane

 Damien Thevenet,^a Reinhard Neier^{a*} and Helen Stoeckli-Evans^b
^aInstitute of Chemistry, University of Neuchâtel, rue Emile-Argand 11, 2009 Neuchâtel, Switzerland, and ^bInstitute of Physics, University of Neuchâtel, rue Emile-Argand 11, 2009 Neuchâtel, Switzerland

Correspondence e-mail: rienhard.neier@unine.ch


Received 30 December 2009; accepted 6 January 2010

 Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.024; wR factor = 0.061; data-to-parameter ratio = 7.6.

The title compound, $\text{C}_{14}\text{H}_{14}\text{O}_2$, crystallizes in the chiral monoclinic space group $P2_1$. This acetal is composed of a planar naphthalene ring with a 1,3-dioxane ring substituent, which has a chair conformation. In the crystal structure, symmetry-related molecules are connected *via* a weak $\text{C}-\text{H}\cdots\text{O}$ interaction to form a helical chain propagating in $[010]$. While there are no $\pi-\pi$ stacking interactions present, there are weak $\text{C}-\text{H}\cdots\pi$ interactions involving the naphthalene aromatic rings, which link the helical chains to form a two-dimensional network in the (011) plane.

Related literature

For information on commonly used protecting groups for carbonyl compounds, see: Kocienski (1994); Showler & Darley (1967). For methods for their deprotection, see: Cordes & Bull (1974); Fujioka *et al.* (2004); Ates *et al.* (2003). For kinetic and thermodynamic studies of acetals and ketals in the naphthalene series and other physical data, see: Newman & Dickson (1970); Carmichael & Hug (1986). For the synthesis of 2-naphthaldehyde acetal, see Gopinath *et al.* (2002). For details of the new photochemical reaction to hydrolyse the acetal into an aldehyde, see Thevenet & Neier (2010). For information on 1,3-dioxane ring related compounds, see: Buys & Eliel (1970). For the synthesis and crystal structure of a related compound, see: Borbas *et al.* (2002). For normal geometric parameters for molecular compounds, see: Allen (2002).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{14}\text{O}_2$	$V = 556.20$ (9) Å ³
$M_r = 214.25$	$Z = 2$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation
$a = 7.5351$ (6) Å	$\mu = 0.08$ mm ⁻¹
$b = 7.8575$ (8) Å	$T = 173$ K
$c = 9.4057$ (9) Å	$0.38 \times 0.30 \times 0.08$ mm
$\beta = 92.839$ (11)°	

Data collection

Stoe IPDS diffractometer	951 reflections with $I > 2\sigma(I)$
4461 measured reflections	$R_{\text{int}} = 0.024$
1098 independent reflections	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.024$	1 restraint
$wR(F^2) = 0.061$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\text{max}} = 0.13$ e Å ⁻³
1098 reflections	$\Delta\rho_{\text{min}} = -0.11$ e Å ⁻³
145 parameters	

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 and Cg2 are the centroids of the $\text{C1}'-\text{C4}'/\text{C9}'/\text{C10}'$ and $\text{C5}'-\text{C10}'$ rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C1}'-\text{H1}'\cdots\text{O2}^{\text{i}}$	0.95	2.60	3.349 (2)	136
$\text{C5}'-\text{H5}'\cdots\text{Cg1}^{\text{ii}}$	0.95	2.70	3.555 (2)	151
$\text{C4}'-\text{H4}'\cdots\text{Cg2}^{\text{ii}}$	0.95	2.92	3.776 (2)	150
$\text{C3}-\text{H3A}\cdots\text{Cg1}^{\text{i}}$	0.99	2.99	3.927 (2)	159

Symmetry codes: (i) $-x + 2, y - \frac{1}{2}, -z + 1$; (ii) $-x + 2, y + \frac{1}{2}, -z + 2$.

Data collection: *EXPOSE* in *IPDS-I* (Stoe & Cie, 2000); cell refinement: *CELL* in *IPDS-I*; data reduction: *INTEGRATE* in *IPDS-I*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *SHELXL97* and *PLATON*.

HSE is grateful to the XRD Application Laboratory, Microsystems Technology Division, Swiss Center for Electronics and Microtechnology, Neuchâtel, for access to the X-ray diffraction equipment.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2685).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
 Ates, A., Gautier, A., Leroy, B., Plancher, J. M., Quesnel, Y., Vanherck, J. C. & Marko, I. E. (2003). *Tetrahedron*, **59**, 8989–8999.
 Borbas, A., Szoba, Z. B., Szilagy, L., Benyei, A. & Liptak, A. (2002). *Tetrahedron*, **58**, 5723–5732.
 Buys, H. R. & Eliel, E. L. (1970). *Tetrahedron Lett.* **32**, 2779–2782.
 Carmichael, I. & Hug, G. L. (1986). *J. Phys. Chem. Ref. Data*, **15**, 1–250.
 Cordes, E. H. & Bull, H. G. (1974). *Chem. Rev.* **74**, 581–603.
 Fujioka, H., Sawama, Y., Murata, N., Okitsu, T., Kubo, O., Matsuda, S. & Kita, Y. (2004). *J. Am. Chem. Soc.* **126**, 11800–11801.
 Gopinath, R., Haque, S. J. & Patel, B. K. (2002). *J. Org. Chem.* **67**, 5842–5845.

- Kocienski, P. J. (1994). *Carbonyl Protecting Groups*, in *Protecting Groups*, ch. 5. New York: Thieme Medical Publishers.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Newman, M. S. & Dickson, R. E. (1970). *J. Am. Chem. Soc.* **92**, 6880–6884.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Showler, A. J. & Darley, P. A. (1967). *Chem. Rev.* **67**, 427–440.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Stoe & Cie (2000). *IPDS-I*. Stoe & Cie GmbH, Darmstadt, Germany.
- Thevenet, D. & Neier, R. (2010). In preparation.

supporting information

Acta Cryst. (2010). E66, o473–o474 [https://doi.org/10.1107/S1600536810000644]

2-(2-Naphthyl)-1,3-dioxane

Damien Thevenet, Reinhard Neier and Helen Stoeckli-Evans

S1. Comment

Acetals are the most commonly used protecting groups for carbonyl compounds in organic synthesis (Kocienski, 1994; Showler & Darley, 1967), and many methods have been developed for their deprotection (Cordes & Bull, 1974; Fujioka *et al.*, 2004; Ates *et al.*, 2003). The title 2-naphthaldehyde acetal (Newman & Dickson, 1970; Carmichael & Hug, 1986) was synthesized to investigate the scope of a new photochemical reaction capable of hydrolysing the acetal into an aldehyde (Thevenet & Neier, 2010). The NMR spectra of the unsubstituted 1,3-dioxane ring displays a complicated AA'BB'MN system (Buys & Eliel, 1970), and the X-ray crystal structure was helpful for the interpretation of the NMR spectra (Thevenet & Neier, 2010).

The structure of the title compound is illustrated in Fig. 1, and the geometrical parameters are given in the Supplementary information and the archived CIF. The bond lengths and angles are close to those in three similar compounds located in the Cambridge Crystal Structure Database (CSD, V 5.30, last update Sept. 2009; Allen, 2002). For example, methyl 2,3-di-*O*-acetyl-4,6-*O*-(2-naphthyl)methylene- α -*D*-galactopyranoside (Borbás *et al.*, 2002), which also crystallized in the monoclinic space group *P*2₁, and where the naphthalene ring is planar and the two six-membered rings in the galactopyranoside unit have chair conformations.

In the crystal of the title compound symmetry related molecules are connected *via* a C—H \cdots O interaction (Table 1) giving rise to the formation of helical chains propagating in [010]. These chains are further linked *via* weak C—H \cdots π interactions to form a two-dimensional network in (011) - see Fig. 2 and Table 1 for details.

S2. Experimental

The title compound was synthesized using a modified strategy described by (Gopinath *et al.*, 2002). To a solution of 2-naphthaldehyde (0.64 mmol), trimethylorthoformate (1.41 mmol) and 1,3-propanediol (5.12 mmol) in dry nitromethane (2 ml) was added tetrabutylammonium tribromide (0.025 mmol). The homogeneous reaction mixture was stirred at r.t. and the progress of the reaction monitored by TLC and GC. After completion of the reaction the mixture was poured into a solution of NaHCO₃ (10 ml) and the products were extracted with diethyl ether (3 \times 10 ml). The organic layer was separated, dried over anhydrous Na₂SO₄ and concentrated. The white solid obtained was purified by recrystallization in MeOH, giving colourless thin plate-like crystals of the title compound.

¹H NMR 400 MHz (CDCl₃) δ 7.97 (br s, 1H, H_{1'}), 7.85 (m, 3H, H_{4',5',8'}), 7.60 (dd, 1H, ³J_{3',4'} = 8.5 Hz, ³J_{3',1'} = 1.7 Hz, H_{3'}), 7.48 (m, 2H, H_{6',7'}), 5.68 (s, 1H, H₁), 4.33 (dddd, 2H, ²J_{3e-3a;5e-5a} = -11.7 Hz, ³J_{3e-4a;5e-4a} = 5.0 Hz, ³J_{3e-4e;5e-4e} = 1.5 Hz, ⁴J_{3e-5e} = 3.0 Hz, H_{3e,5e}), 4.06 (ddd, 2H, ²J_{3a-3e;5a-5e} = -11.7 Hz, ³J_{3a-4a;5a-4a} = 12.4 Hz, ³J_{3a-4e;5a-4e} = 2.7 Hz, H_{3a,5a}), 2.29 (dtt, 1H, ²J_{4a-4e} = -13.5 Hz, ³J_{4a-3a;4a-5a} = 12.4 Hz, ³J_{4a-3e;4a-5e} = 5.0 Hz, H_{4a}), 1.50 (dtt, 1H, ²J_{4e-4a} = -13.5 Hz, ³J_{4e-3a;4e-5a} = 2.7 Hz, ³J_{4e-3e;4e-5e} = 1.5 Hz, H_{4e}); ¹³C NMR 100 MHz (CDCl₃) δ 136.1 (C₂), (133.6, 133.1) (C_{9',10'}), (128.4, 128.1, 127.7) (C_{4',5',8'}), (126.2, 126.0) (C_{6',7'}), 125.3 (C_{1'}), 123.8 (C_{3'}), 101.8 (C₁), 67.5 (C_{3,5}), 25.9 (C₄); HRMS (ESI, +): [M + Na]⁺ = 237.09. Note: The same numbering scheme has been used for the crystal structure (Fig. 1). The torsional angles of the 1,3-dioxane ring were

measured to estimate the coupling constants according to the Karplus equation.

S3. Refinement

In the final cycles of refinement, in the absence of significant anomalous scattering effects, 944 (93%) Friedel pairs were merged and $\Delta f'$ set to zero. The H-atoms could all be located in difference electron-density maps. In the final cycles of refinement they were included in calculated positions and treated as riding atoms: C—H = 0.95–1.0 Å, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{parent C-atom})$. Using the one-circle Stoe Image Plate Diffraction System it is not always possible to measure 100% of the Ewald sphere, and here only 93.7% of the data were accessible out to 50° in 2θ . This has little effect on the bond distances and angles when comparing their values with those of the related structure mentioned above (Borbás *et al.*, 2002).




Figure 1

A view of the molecular structure of the title compound, with displacement ellipsoids drawn at the 50% probability level.




Figure 2

A view along the *a* axis of the crystal packing of the title compound. The C—H \cdots O and C—H \cdots π interactions are shown as dotted cyan and black lines, respectively. [The blue balls represent the centroids of the two aromatic rings; H-atoms not involved in the C—H \cdots O and C—H \cdots π interactions have been omitted for clarity; the C—H \cdots π interactions are shown for one molecule only; see Table 1 for details].

2-(2-Naphthyl)-1,3-dioxane

Crystal data

$C_{14}H_{14}O_2$
 $M_r = 214.25$
 Monoclinic, $P2_1$
 Hall symbol: P 2yb
 $a = 7.5351$ (6) Å
 $b = 7.8575$ (8) Å
 $c = 9.4057$ (9) Å
 $\beta = 92.839$ (11)°
 $V = 556.20$ (9) Å³
 $Z = 2$

$F(000) = 228$
 $D_x = 1.279$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 4553 reflections
 $\theta = 2.1$ – 26.0 °
 $\mu = 0.08$ mm⁻¹
 $T = 173$ K
 Plate, colourless
 $0.38 \times 0.30 \times 0.08$ mm

Data collection

Stoe IPDS
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ rotation scans
 4461 measured reflections
 1098 independent reflections

951 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.024$
 $\theta_{max} = 26.0$ °, $\theta_{min} = 2.2$ °
 $h = -8 \rightarrow 8$
 $k = -9 \rightarrow 9$
 $l = -11 \rightarrow 11$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.024$
 $wR(F^2) = 0.061$
 $S = 1.05$
 1098 reflections
 145 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0412P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.13 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.11 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. In the final cycles of refinement, in the absence of significant anomalous scattering effects, 944 (93%) Friedel pairs were merged and $\Delta f''$ set to zero. The H-atoms could all be located in difference electron-density maps. In the final cycles of refinement they were included in calculated positions and treated as riding atoms: C—H = 0.95 - 1.0 Å, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{parent C-atoms})$. Using the one-circle Stoe Image Plate Diffraction System it is not always possible to measure 100% of the Ewald sphere, and here only 93.7% of the data were accessible out to 50° in 2θ .

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O2	1.16506 (19)	0.57058 (16)	0.43163 (12)	0.0337 (4)
O6	1.33418 (18)	0.38156 (15)	0.57014 (12)	0.0295 (4)
C1	1.1616 (2)	0.4392 (2)	0.53276 (17)	0.0251 (5)
C1'	0.9100 (2)	0.46504 (19)	0.69218 (16)	0.0239 (5)
C2'	1.0796 (2)	0.5075 (2)	0.66364 (16)	0.0240 (5)
C3	1.2305 (3)	0.5057 (3)	0.30168 (18)	0.0417 (7)
C3'	1.1796 (3)	0.6163 (2)	0.75664 (17)	0.0277 (6)
C4	1.4131 (3)	0.4321 (3)	0.32815 (19)	0.0400 (7)
C4'	1.1073 (3)	0.6764 (2)	0.87667 (18)	0.0297 (6)
C5	1.4128 (3)	0.3067 (3)	0.44944 (18)	0.0348 (6)
C5'	0.8555 (3)	0.6908 (2)	1.03543 (18)	0.0309 (6)
C6'	0.6865 (3)	0.6467 (2)	1.06380 (18)	0.0316 (6)
C7'	0.5852 (3)	0.5438 (2)	0.96939 (18)	0.0328 (6)
C8'	0.6551 (2)	0.4853 (2)	0.84779 (18)	0.0284 (5)
C9'	0.8310 (2)	0.52609 (19)	0.81562 (16)	0.0236 (5)
C10'	0.9334 (2)	0.63231 (19)	0.91039 (17)	0.0239 (5)
H1	1.08840	0.34250	0.49320	0.0300*
H1'	0.84350	0.39320	0.62810	0.0290*
H3'	1.29720	0.64750	0.73540	0.0330*
H3A	1.14900	0.41670	0.26250	0.0500*
H3E	1.23480	0.59870	0.23090	0.0500*
H4'	1.17550	0.74940	0.93860	0.0360*
H4A	1.45060	0.37410	0.24110	0.0480*
H4E	1.49890	0.52450	0.35160	0.0480*
H5'	0.92210	0.76160	1.10020	0.0370*

H5A	1.53630	0.27170	0.47590	0.0420*
H5E	1.34500	0.20400	0.41910	0.0420*
H6'	0.63660	0.68620	1.14860	0.0380*
H7'	0.46700	0.51460	0.99020	0.0390*
H8'	0.58470	0.41640	0.78400	0.0340*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O2	0.0500 (9)	0.0289 (6)	0.0225 (6)	0.0088 (6)	0.0049 (5)	0.0012 (5)
O6	0.0283 (8)	0.0376 (7)	0.0228 (5)	0.0058 (6)	0.0027 (5)	-0.0007 (5)
C1	0.0269 (11)	0.0239 (8)	0.0244 (8)	-0.0013 (6)	0.0008 (7)	0.0000 (6)
C1'	0.0245 (11)	0.0228 (8)	0.0240 (8)	-0.0016 (6)	-0.0023 (7)	0.0007 (6)
C2'	0.0265 (11)	0.0229 (8)	0.0225 (8)	-0.0003 (7)	0.0012 (7)	0.0021 (7)
C3	0.0679 (17)	0.0358 (9)	0.0220 (8)	0.0095 (10)	0.0084 (9)	0.0013 (8)
C3'	0.0224 (11)	0.0303 (9)	0.0307 (9)	-0.0050 (7)	0.0031 (7)	-0.0032 (7)
C4	0.0533 (16)	0.0374 (10)	0.0306 (9)	-0.0015 (9)	0.0158 (9)	-0.0061 (8)
C4'	0.0273 (12)	0.0299 (9)	0.0317 (9)	-0.0050 (7)	-0.0007 (7)	-0.0058 (7)
C5	0.0371 (13)	0.0395 (10)	0.0283 (9)	0.0069 (8)	0.0077 (8)	-0.0053 (8)
C5'	0.0345 (14)	0.0286 (9)	0.0298 (9)	0.0016 (7)	0.0026 (8)	-0.0023 (7)
C6'	0.0322 (12)	0.0327 (9)	0.0309 (8)	0.0066 (8)	0.0105 (7)	0.0024 (7)
C7'	0.0230 (12)	0.0382 (11)	0.0377 (9)	0.0031 (7)	0.0063 (8)	0.0075 (8)
C8'	0.0222 (11)	0.0316 (9)	0.0313 (8)	-0.0035 (8)	0.0004 (7)	0.0029 (7)
C9'	0.0222 (11)	0.0226 (8)	0.0257 (8)	0.0007 (6)	-0.0005 (7)	0.0051 (6)
C10'	0.0242 (11)	0.0211 (7)	0.0264 (8)	0.0010 (7)	0.0013 (7)	0.0007 (6)

Geometric parameters (Å, °)

O2—C1	1.405 (2)	C8'—C9'	1.411 (2)
O2—C3	1.434 (2)	C9'—C10'	1.421 (2)
O6—C1	1.405 (2)	C1—H1	1.0000
O6—C5	1.433 (2)	C1'—H1'	0.9500
C1—C2'	1.504 (2)	C3—H3A	0.9900
C1'—C2'	1.360 (2)	C3—H3E	0.9900
C1'—C9'	1.415 (2)	C3'—H3'	0.9500
C2'—C3'	1.414 (2)	C4—H4A	0.9900
C3—C4	1.502 (3)	C4—H4E	0.9900
C3'—C4'	1.362 (3)	C4'—H4'	0.9500
C4—C5	1.508 (3)	C5—H5A	0.9900
C4'—C10'	1.407 (3)	C5—H5E	0.9900
C5'—C6'	1.359 (3)	C5'—H5'	0.9500
C5'—C10'	1.417 (2)	C6'—H6'	0.9500
C6'—C7'	1.399 (3)	C7'—H7'	0.9500
C7'—C8'	1.363 (2)	C8'—H8'	0.9500
C1—O2—C3	109.55 (14)	C9'—C1'—H1'	119.00
C1—O6—C5	110.36 (13)	O2—C3—H3A	110.00
O2—C1—O6	110.99 (13)	O2—C3—H3E	110.00

O2—C1—C2'	108.32 (13)	C4—C3—H3A	110.00
O6—C1—C2'	108.84 (13)	C4—C3—H3E	110.00
C2'—C1'—C9'	121.09 (14)	H3A—C3—H3E	108.00
C1—C2'—C1'	120.15 (14)	C2'—C3'—H3'	120.00
C1—C2'—C3'	119.63 (14)	C4'—C3'—H3'	120.00
C1'—C2'—C3'	120.22 (15)	C3—C4—H4A	110.00
O2—C3—C4	110.27 (15)	C3—C4—H4E	110.00
C2'—C3'—C4'	119.98 (19)	C5—C4—H4A	110.00
C3—C4—C5	109.96 (18)	C5—C4—H4E	110.00
C3'—C4'—C10'	121.12 (17)	H4A—C4—H4E	108.00
O6—C5—C4	110.33 (18)	C3'—C4'—H4'	119.00
C6'—C5'—C10'	120.74 (16)	C10'—C4'—H4'	119.00
C5'—C6'—C7'	120.65 (17)	O6—C5—H5A	110.00
C6'—C7'—C8'	120.42 (19)	O6—C5—H5E	110.00
C7'—C8'—C9'	120.64 (16)	C4—C5—H5A	110.00
C1'—C9'—C8'	122.48 (14)	C4—C5—H5E	110.00
C1'—C9'—C10'	118.47 (14)	H5A—C5—H5E	108.00
C8'—C9'—C10'	119.06 (14)	C6'—C5'—H5'	120.00
C4'—C10'—C5'	122.42 (15)	C10'—C5'—H5'	120.00
C4'—C10'—C9'	119.09 (14)	C5'—C6'—H6'	120.00
C5'—C10'—C9'	118.49 (15)	C7'—C6'—H6'	120.00
O2—C1—H1	110.00	C6'—C7'—H7'	120.00
O6—C1—H1	110.00	C8'—C7'—H7'	120.00
C2'—C1—H1	110.00	C7'—C8'—H8'	120.00
C2'—C1'—H1'	119.00	C9'—C8'—H8'	120.00
C3—O2—C1—O6	64.83 (17)	O2—C3—C4—C5	51.9 (2)
C3—O2—C1—C2'	-175.74 (14)	C2'—C3'—C4'—C10'	-0.1 (3)
C1—O2—C3—C4	-58.4 (2)	C3—C4—C5—O6	-50.8 (2)
C5—O6—C1—O2	-64.14 (17)	C3'—C4'—C10'—C5'	179.07 (16)
C5—O6—C1—C2'	176.74 (14)	C3'—C4'—C10'—C9'	-1.3 (2)
C1—O6—C5—C4	56.5 (2)	C10'—C5'—C6'—C7'	-0.6 (3)
O2—C1—C2'—C1'	104.47 (17)	C6'—C5'—C10'—C4'	179.42 (16)
O2—C1—C2'—C3'	-75.40 (18)	C6'—C5'—C10'—C9'	-0.2 (2)
O6—C1—C2'—C1'	-134.75 (15)	C5'—C6'—C7'—C8'	0.4 (3)
O6—C1—C2'—C3'	45.38 (19)	C6'—C7'—C8'—C9'	0.6 (2)
C9'—C1'—C2'—C1	179.19 (14)	C7'—C8'—C9'—C1'	178.60 (15)
C9'—C1'—C2'—C3'	-0.9 (2)	C7'—C8'—C9'—C10'	-1.4 (2)
C2'—C1'—C9'—C8'	179.59 (15)	C1'—C9'—C10'—C4'	1.6 (2)
C2'—C1'—C9'—C10'	-0.5 (2)	C1'—C9'—C10'—C5'	-178.80 (14)
C1—C2'—C3'—C4'	-178.90 (15)	C8'—C9'—C10'—C4'	-178.48 (15)
C1'—C2'—C3'—C4'	1.2 (2)	C8'—C9'—C10'—C5'	1.2 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C1'—H1' \cdots O2 ⁱ	0.95	2.60	3.349 (2)	136
C5'—H5' \cdots Cg1 ⁱⁱ	0.95	2.70	3.555 (2)	151

C4'—H4'...Cg2 ⁱⁱ	0.95	2.92	3.776 (2)	150
C3—H3A...Cg1 ⁱ	0.99	2.99	3.927 (2)	159

Symmetry codes: (i) $-x+2, y-1/2, -z+1$; (ii) $-x+2, y+1/2, -z+2$.