

## (2,7-Dimethoxynaphthalene-1,8-diyl)- bis(4-fluorobenzoyl)dimethanone

Shoji Watanabe,<sup>a</sup> Atsushi Nagasawa,<sup>a</sup> Akiko Okamoto,<sup>a</sup>  
Keiichi Noguchi<sup>b</sup> and Noriyuki Yonezawa<sup>a\*</sup>

<sup>a</sup>Department of Organic and Polymer Materials Chemistry, Tokyo University of Agriculture & Technology, Koganei, Tokyo 184-8588, Japan, and <sup>b</sup>Instrumentation Analysis Center, Tokyo University of Agriculture & Technology, Koganei, Tokyo 184-8588, Japan

Correspondence e-mail: yonezawa@cc.tuat.ac.jp


Received 2 December 2009; accepted 5 January 2010

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.034;  $wR$  factor = 0.095; data-to-parameter ratio = 11.7.

The title compound,  $C_{26}H_{18}F_2O_4$ , is a naphthalene derivative in which the two aryl groups at the 1- and 8-positions (*peri* positions) are *anti* to each other. There is an appreciable difference in the dihedral angles between the naphthalene ring system and the two benzene rings [ $66.88(7)^\circ$  and  $88.09(6)^\circ$ ]. In the crystal, weak C—H···O interactions involving one of the carbonyl groups and an aromatic C—H group *ortho* to the F atom seem to stabilize the packing of the molecules.

### Related literature

Our study on the selective electrophilic aromatic aroylation of 2,7-dimethoxynaphthalene, has shown *peri*-arylnaphthalene compounds to be formed regioselectively with the aid of a suitable acidic mediator, see: (Okamoto & Yonezawa, 2009). For related structures, see: Nakema *et al.* (2007, 2008); Mitsui *et al.* (2009).



### Experimental

#### Crystal data

$C_{26}H_{18}F_2O_4$   
 $M_r = 432.42$

Monoclinic,  $P2_1/c$   
 $a = 9.87444(18)\text{ \AA}$

$b = 17.0275(3)\text{ \AA}$   
 $c = 14.9671(3)\text{ \AA}$   
 $\beta = 126.871(1)^\circ$   
 $V = 2013.19(7)\text{ \AA}^3$   
 $Z = 4$

$\text{Cu } K\alpha$  radiation  
 $\mu = 0.91\text{ mm}^{-1}$   
 $T = 296\text{ K}$   
 $0.40 \times 0.40 \times 0.10\text{ mm}$

#### Data collection

Rigaku R-AXIS RAPID  
diffractometer  
Absorption correction: numerical  
(*NUMABS*; Higashi, 1999)  
 $T_{\min} = 0.713$ ,  $T_{\max} = 0.915$

36825 measured reflections  
3682 independent reflections  
3338 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.095$   
 $S = 1.07$   
3682 reflections  
314 parameters

H atoms treated by a mixture of  
independent and constrained  
refinement  
 $\Delta\rho_{\max} = 0.22\text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.18\text{ e } \text{\AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C16—H7···O1 <sup>i</sup>	1.00	2.54	3.493 (2)	158
C21—H10···O1 <sup>ii</sup>	0.99	2.68	3.636 (2)	161

Symmetry codes: (i)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (ii)  $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$ .

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC, 2004); program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP* (Burnett & Johnson, 1996); software used to prepare material for publication: *SHELXL97*.

This work was partially supported by the Iketani Science and Technology Foundation, Tokyo, Japan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DS2015).

### References

- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Higashi, T. (1999). *NUMABS*. Rigaku Corporation, Tokyo, Japan.
- Mitsui, R., Noguchi, K. & Yonezawa, N. (2009). *Acta Cryst. E65*, o543.
- Nakaema, K., Okamoto, A., Noguchi, K. & Yonezawa, N. (2007). *Acta Cryst. E63*, o4120.
- Nakaema, K., Watanabe, S., Okamoto, A., Noguchi, K. & Yonezawa, N. (2008). *Acta Cryst. E64*, o807.
- Okamoto, A. & Yonezawa, N. (2009). *Chem. Lett.* **38**, 914–915.
- Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2004). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

# supporting information

*Acta Cryst.* (2010). E66, o329 [https://doi.org/10.1107/S1600536810000486]

## (2,7-Dimethoxynaphthalene-1,8-diyl)bis(4-fluorobenzoyl)dimethanone

Shoji Watanabe, Atsushi Nagasawa, Akiko Okamoto, Keiichi Noguchi and Noriyuki Yonezawa

### S1. Comment

In the course of our study on selective electrophilic aromatic arylation of 2,7-dimethoxynaphthalene, *peri*-arylnaphthalene compounds have proved to be formed regioselectively with the aid of suitable acidic mediator (Okamoto & Yonezawa, 2009). The aryl groups at 1,8-positions of the naphthalene rings in these compounds are oriented in opposite fashion and are found to be non-coplanar resulting in partial disruption in  $\pi$ -conjugation systems. Recently, we have reported the X-ray crystal structures of 1,8-bis(4-chlorobenzoyl)-2,7-dimethoxynaphthalene (Nakaema *et al.*, 2007) and 1,8-dibenzoyl-2,7-dimethoxynaphthalene (Nakaema *et al.*, 2008). As a part of the course of our continuous study on the molecular structures of this kind of homologous molecules, the X-ray crystal structure of title compound, *peri*-arylnaphthalene bearing fluoro groups, is discussed in this report.

In the molecule (Fig. 1), the dihedral angle between benzene rings [C12–C17] and [C19–C24] is 32.34 (8) $^\circ$ , which is distinctively larger than that of 1,8-bis(4-chlorobenzoyl)-2,7-dimethoxynaphthalene [7.99 (8) $^\circ$ ]. The dihedral angles between the naphthalene ring [C1–C10] plane and the planes of two benzene rings [C12–C17] and [C19–C24] are 66.88 (7) $^\circ$  and 88.12 (6) $^\circ$ , respectively. The difference between two dihedral angles is larger than that of the analogous compound, which has the corresponding angles of 71.98 (7) $^\circ$  and 71.55 (7) $^\circ$ .

The molecules are packed in the crystal lattice apparently stabilized by C—H $\cdots$ O interactions involving C16, C21 and O1 [2.54 Å, 158 $^\circ$  and 2.68 Å, 161 $^\circ$ ] (Fig. 2 and Table 1).

### S2. Experimental


To a 10 ml flask, 4-fluorobenzoic acid (4.4 mmol, 616.5 mg) and phosphorus pentoxide–methanesulfonic acid (P<sub>2</sub>O<sub>5</sub>–MsOH; 8.8 ml) were placed and stirred at 60°C. To the reaction mixture thus obtained, 2,7-dimethoxynaphthalene (2.0 mmol, 376.4 mg) was added. After the reaction mixture was stirred at 60 °C for 1 h, it was poured into ice-cold water (10 ml) and the mixture was extracted with CHCl<sub>3</sub> (10 ml  $\times$  3). The combined extracts were washed with 2 M aqueous NaOH followed by washing with brine. The organic layers thus obtained were dried over anhydrous MgSO<sub>4</sub>. The solvent was removed under reduced pressure to give cake (98% yield). Crude product was purified by recrystallization from EtOH (77% isolated yield). Furthermore, the isolated product was crystallized from toluene–hexane to give single-crystal.

Spectroscopic Data:


<sup>1</sup>H NMR  $\delta$  (300 MHz, CDCl<sub>3</sub>): 3.70 (6H, s), 7.02 (4H, dd,  $J$  = 8.6 Hz), 7.21 (2H, d,  $J$  = 8.7 Hz), 7.71 (4H, dd,  $J$  = 8.41 Hz), 7.96 (2H, d,  $J$  = 8.7 Hz). <sup>13</sup>C NMR  $\delta$  (300 MHz, CDCl<sub>3</sub>): 56.289, 111.12, 114.92, 115.21, 125.47, 129.76, 131.52, 131.64, 132.25, 135.19, 156.24, 163.81, 167.17, 195.38. IR (KBr): 1596 (C=O), 1270 (Ar—O—Me). m.p. = 196°C. Anal. Calcd for C<sub>26</sub>H<sub>18</sub>F<sub>2</sub>O<sub>4</sub>; C, 70.27; H, 4.20. Found C, 72.05; H, 4.20.

**S3. Refinement**

All the H atoms were found in difference maps and were subsequently refined as riding atoms, with C—H = 0.93 (aromatic) and 0.96 (methyl) Å, and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

**Figure 1**

Molecular structure with displacement ellipsoids at 50% probability for non-H atoms.

**Figure 2**

Weak C–H···O interactions (dotted lines).

**(2,7-Dimethoxynaphthalene-1,8-diyl)bis(4-fluorobenzoyl)dimethanone***Crystal data*

$C_{26}H_{18}F_2O_4$   
 $M_r = 432.42$   
Monoclinic,  $P2_1/c$   
Hall symbol: -P 2ybc  
 $a = 9.87444 (18)$  Å  
 $b = 17.0275 (3)$  Å  
 $c = 14.9671 (3)$  Å  
 $\beta = 126.871 (1)^\circ$   
 $V = 2013.19 (7)$  Å<sup>3</sup>  
 $Z = 4$

$F(000) = 896$   
 $D_x = 1.427 \text{ Mg m}^{-3}$   
Melting point: 469 K  
Cu  $K\alpha$  radiation,  $\lambda = 1.54187$  Å  
Cell parameters from 33844 reflections  
 $\theta = 3.7\text{--}68.2^\circ$   
 $\mu = 0.91 \text{ mm}^{-1}$   
 $T = 296 \text{ K}$   
Platelet, yellow  
0.40 × 0.40 × 0.10 mm

*Data collection*

Rigaku R-AXIS RAPID  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
Detector resolution: 10.00 pixels mm<sup>-1</sup>  
 $\omega$  scans  
Absorption correction: numerical  
(*NUMABS*; Higashi, 1999)  
 $T_{\min} = 0.713$ ,  $T_{\max} = 0.915$

36825 measured reflections  
3682 independent reflections  
3338 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$   
 $\theta_{\max} = 68.2^\circ$ ,  $\theta_{\min} = 4.5^\circ$   
 $h = -11 \rightarrow 11$   
 $k = -20 \rightarrow 20$   
 $l = -18 \rightarrow 18$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.034$$

$$wR(F^2) = 0.095$$

$$S = 1.07$$

3682 reflections

314 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sitesH atoms treated by a mixture of independent  
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0494P)^2 + 0.3061P]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.22 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.18 \text{ e \AA}^{-3}$$

Extinction correction: *SHELXL97* (Sheldrick,  
2008),  $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$ 

Extinction coefficient: 0.0030 (3)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	-0.21053 (13)	0.44424 (5)	-0.09780 (9)	0.0824 (3)
F2	0.35965 (13)	0.21569 (6)	0.65179 (7)	0.0878 (3)
O1	-0.24490 (12)	0.16209 (5)	0.15843 (7)	0.0570 (2)
O2	0.09608 (12)	0.18156 (5)	0.15466 (8)	0.0598 (2)
O3	-0.55074 (11)	0.11265 (6)	-0.11168 (8)	0.0634 (3)
O4	0.31918 (12)	0.01602 (6)	0.31605 (8)	0.0684 (3)
C1	-0.26555 (15)	0.08412 (7)	0.02104 (9)	0.0444 (3)
C2	-0.42192 (16)	0.05978 (7)	-0.07141 (10)	0.0503 (3)
C3	-0.44496 (18)	-0.01516 (8)	-0.11871 (11)	0.0587 (3)
H1	-0.5531	-0.0296	-0.1825	0.070*
C4	-0.31155 (19)	-0.06521 (7)	-0.07028 (11)	0.0580 (3)
H2	-0.3237	-0.1170	-0.1017	0.070*
C5	-0.15072 (17)	-0.04486 (7)	0.02702 (10)	0.0501 (3)
C6	-0.01766 (19)	-0.10000 (7)	0.08034 (12)	0.0574 (3)
H3	-0.0389	-0.1520	0.0463	0.067*
C7	0.13559 (18)	-0.08282 (8)	0.17722 (11)	0.0573 (3)
H4	0.2247	-0.1225	0.2143	0.071*
C8	0.16525 (16)	-0.00701 (7)	0.22300 (10)	0.0516 (3)
C9	0.04104 (15)	0.05019 (7)	0.17254 (9)	0.0450 (3)
C10	-0.12396 (15)	0.03188 (7)	0.07386 (9)	0.0438 (3)
C11	-0.25571 (14)	0.16202 (7)	0.07300 (9)	0.0444 (3)
C12	-0.25352 (14)	0.23685 (7)	0.02252 (9)	0.0449 (3)
C13	-0.20106 (18)	0.30478 (8)	0.08654 (11)	0.0559 (3)

H5	-0.1715	0.3006	0.1573	0.068*
C14	-0.1873 (2)	0.37483 (8)	0.04636 (12)	0.0631 (4)
H6	-0.1474	0.4221	0.0902	0.077*
C15	-0.22767 (18)	0.37560 (8)	-0.05889 (12)	0.0592 (3)
C16	-0.28166 (19)	0.31051 (8)	-0.12562 (11)	0.0614 (3)
H7	-0.3057	0.3146	-0.2008	0.086*
C17	-0.29344 (17)	0.24052 (7)	-0.08343 (10)	0.0526 (3)
H8	-0.3335	0.1938	-0.1306	0.059*
C18	0.09933 (14)	0.13262 (7)	0.21535 (10)	0.0461 (3)
C19	0.16719 (14)	0.15291 (7)	0.33199 (10)	0.0459 (3)
C20	0.29027 (17)	0.21100 (8)	0.38710 (12)	0.0576 (3)
H9	0.3323	0.2342	0.3471	0.074*
C21	0.35501 (18)	0.23239 (9)	0.49493 (12)	0.0642 (4)
H10	0.4460	0.2721	0.5367	0.088*
C22	0.29323 (17)	0.19592 (8)	0.54466 (11)	0.0598 (4)
C23	0.17105 (17)	0.13886 (9)	0.49316 (11)	0.0597 (3)
H11	0.1302	0.1174	0.5288	0.071*
C24	0.10868 (16)	0.11671 (8)	0.38576 (10)	0.0527 (3)
H12	0.0217	0.0758	0.3467	0.062*
C25	-0.71784 (19)	0.08920 (13)	-0.20143 (14)	0.0706 (4)
C26	0.4565 (2)	-0.03775 (11)	0.36866 (15)	0.0697 (4)
H13	-0.751 (3)	0.0383 (12)	-0.1820 (16)	0.098 (6)*
H14	-0.786 (3)	0.1326 (12)	-0.2121 (15)	0.088 (6)*
H15	-0.728 (2)	0.0805 (11)	-0.2710 (16)	0.094 (6)*
H16	0.429 (2)	-0.0860 (11)	0.3956 (14)	0.081 (5)*
H17	0.556 (3)	-0.0075 (11)	0.4306 (16)	0.094 (6)*
H18	0.479 (2)	-0.0552 (10)	0.3166 (15)	0.082 (5)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
F1	0.1025 (7)	0.0541 (5)	0.1006 (7)	0.0052 (4)	0.0663 (6)	0.0177 (4)
F2	0.0833 (6)	0.1105 (8)	0.0550 (5)	-0.0057 (5)	0.0337 (5)	-0.0239 (5)
O1	0.0680 (6)	0.0644 (6)	0.0454 (5)	0.0048 (4)	0.0377 (4)	0.0000 (4)
O2	0.0666 (6)	0.0548 (5)	0.0592 (5)	-0.0018 (4)	0.0384 (5)	0.0083 (4)
O3	0.0468 (5)	0.0667 (6)	0.0569 (5)	-0.0010 (4)	0.0206 (4)	-0.0085 (4)
O4	0.0548 (5)	0.0657 (6)	0.0631 (6)	0.0165 (4)	0.0237 (5)	0.0001 (5)
C1	0.0507 (6)	0.0438 (6)	0.0414 (6)	-0.0025 (5)	0.0291 (5)	-0.0006 (5)
C2	0.0523 (7)	0.0531 (7)	0.0445 (6)	-0.0034 (5)	0.0285 (6)	-0.0011 (5)
C3	0.0631 (8)	0.0568 (7)	0.0490 (7)	-0.0149 (6)	0.0297 (6)	-0.0093 (6)
C4	0.0771 (9)	0.0442 (7)	0.0547 (7)	-0.0094 (6)	0.0406 (7)	-0.0078 (5)
C5	0.0678 (8)	0.0425 (6)	0.0498 (6)	-0.0030 (5)	0.0405 (6)	-0.0011 (5)
C6	0.0821 (9)	0.0425 (6)	0.0613 (8)	0.0044 (6)	0.0503 (8)	0.0006 (6)
C7	0.0729 (9)	0.0493 (7)	0.0589 (8)	0.0160 (6)	0.0444 (7)	0.0089 (6)
C8	0.0579 (7)	0.0525 (7)	0.0489 (6)	0.0078 (6)	0.0345 (6)	0.0051 (5)
C9	0.0527 (7)	0.0435 (6)	0.0443 (6)	0.0026 (5)	0.0321 (5)	0.0020 (5)
C10	0.0548 (7)	0.0418 (6)	0.0416 (6)	-0.0004 (5)	0.0325 (5)	0.0016 (5)
C11	0.0403 (6)	0.0519 (7)	0.0381 (6)	0.0034 (5)	0.0218 (5)	-0.0015 (5)

C12	0.0437 (6)	0.0462 (6)	0.0427 (6)	0.0053 (5)	0.0248 (5)	-0.0016 (5)
C13	0.0647 (8)	0.0530 (7)	0.0477 (7)	0.0040 (6)	0.0325 (6)	-0.0046 (5)
C14	0.0718 (9)	0.0458 (7)	0.0674 (8)	0.0003 (6)	0.0395 (7)	-0.0074 (6)
C15	0.0637 (8)	0.0480 (7)	0.0692 (8)	0.0074 (6)	0.0417 (7)	0.0103 (6)
C16	0.0732 (9)	0.0610 (8)	0.0541 (7)	0.0048 (7)	0.0405 (7)	0.0063 (6)
C17	0.0621 (8)	0.0508 (7)	0.0463 (6)	0.0010 (6)	0.0332 (6)	-0.0035 (5)
C18	0.0426 (6)	0.0462 (6)	0.0491 (6)	0.0034 (5)	0.0272 (5)	0.0038 (5)
C19	0.0427 (6)	0.0425 (6)	0.0486 (6)	0.0023 (5)	0.0253 (5)	0.0003 (5)
C20	0.0559 (7)	0.0527 (7)	0.0665 (8)	-0.0071 (6)	0.0380 (7)	-0.0078 (6)
C21	0.0566 (8)	0.0597 (8)	0.0686 (9)	-0.0104 (6)	0.0334 (7)	-0.0187 (7)
C22	0.0516 (7)	0.0667 (8)	0.0469 (7)	0.0060 (6)	0.0220 (6)	-0.0099 (6)
C23	0.0537 (7)	0.0735 (9)	0.0485 (7)	0.0017 (6)	0.0288 (6)	0.0034 (6)
C24	0.0478 (6)	0.0552 (7)	0.0477 (6)	-0.0050 (5)	0.0246 (5)	0.0001 (5)
C25	0.0494 (8)	0.0904 (12)	0.0578 (9)	-0.0087 (8)	0.0247 (7)	-0.0112 (8)
C26	0.0586 (9)	0.0752 (10)	0.0733 (10)	0.0202 (8)	0.0385 (8)	0.0156 (8)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

F1—C15	1.3610 (15)	C12—C13	1.3888 (17)
F2—C22	1.3611 (15)	C13—C14	1.379 (2)
O1—C11	1.2176 (14)	C13—H5	0.9168
O2—C18	1.2188 (14)	C14—C15	1.374 (2)
O3—C2	1.3681 (16)	C14—H6	0.9600
O3—C25	1.4236 (17)	C15—C16	1.368 (2)
O4—C8	1.3651 (16)	C16—C17	1.3859 (19)
O4—C26	1.4197 (17)	C16—H7	1.0036
C1—C2	1.3808 (17)	C17—H8	0.9758
C1—C10	1.4299 (16)	C18—C19	1.4874 (16)
C1—C11	1.5110 (16)	C19—C24	1.3854 (17)
C2—C3	1.4100 (18)	C19—C20	1.3902 (17)
C3—C4	1.357 (2)	C20—C21	1.381 (2)
C3—H1	0.9430	C20—H9	0.9912
C4—C5	1.4099 (19)	C21—C22	1.363 (2)
C4—H2	0.9717	C21—H10	0.9902
C5—C6	1.4094 (18)	C22—C23	1.370 (2)
C5—C10	1.4305 (16)	C23—C24	1.3847 (18)
C6—C7	1.358 (2)	C23—H11	0.9157
C6—H3	0.9791	C24—H12	0.9809
C7—C8	1.4074 (18)	C25—H13	1.03 (2)
C7—H4	0.9755	C25—H14	0.95 (2)
C8—C9	1.3826 (17)	C25—H15	1.00 (2)
C9—C10	1.4302 (17)	C26—H16	1.018 (19)
C9—C18	1.5065 (16)	C26—H17	1.00 (2)
C11—C12	1.4881 (16)	C26—H18	0.975 (19)
C12—C17	1.3869 (16)		
C2—O3—C25	118.43 (12)	C13—C14—H6	122.3
C8—O4—C26	119.26 (12)	F1—C15—C16	118.51 (13)

C2—C1—C10	120.12 (11)	F1—C15—C14	118.29 (13)
C2—C1—C11	117.85 (11)	C16—C15—C14	123.19 (12)
C10—C1—C11	121.47 (10)	C15—C16—C17	117.81 (12)
O3—C2—C1	115.41 (11)	C15—C16—H7	119.3
O3—C2—C3	123.16 (11)	C17—C16—H7	122.8
C1—C2—C3	121.42 (12)	C16—C17—C12	121.05 (12)
C4—C3—C2	119.20 (12)	C16—C17—H8	118.7
C4—C3—H1	121.6	C12—C17—H8	120.3
C2—C3—H1	119.2	O2—C18—C19	120.52 (11)
C3—C4—C5	121.88 (12)	O2—C18—C9	119.18 (11)
C3—C4—H2	120.8	C19—C18—C9	120.24 (10)
C5—C4—H2	117.3	C24—C19—C20	119.52 (11)
C6—C5—C4	120.71 (11)	C24—C19—C18	121.83 (11)
C6—C5—C10	119.72 (12)	C20—C19—C18	118.64 (11)
C4—C5—C10	119.55 (12)	C21—C20—C19	120.52 (13)
C7—C6—C5	121.59 (12)	C21—C20—H9	122.1
C7—C6—H3	120.2	C19—C20—H9	117.3
C5—C6—H3	118.2	C22—C21—C20	118.23 (13)
C6—C7—C8	119.37 (12)	C22—C21—H10	120.0
C6—C7—H4	120.4	C20—C21—H10	121.7
C8—C7—H4	120.3	F2—C22—C21	118.26 (13)
O4—C8—C9	115.77 (11)	F2—C22—C23	118.51 (14)
O4—C8—C7	122.63 (11)	C21—C22—C23	123.20 (12)
C9—C8—C7	121.52 (12)	C22—C23—C24	118.31 (13)
C8—C9—C10	119.88 (11)	C22—C23—H11	120.2
C8—C9—C18	115.89 (11)	C24—C23—H11	121.5
C10—C9—C18	123.37 (10)	C23—C24—C19	120.21 (12)
C1—C10—C9	124.44 (10)	C23—C24—H12	120.4
C1—C10—C5	117.74 (11)	C19—C24—H12	119.4
C9—C10—C5	117.78 (11)	O3—C25—H13	110.7 (11)
O1—C11—C12	120.91 (10)	O3—C25—H14	104.1 (11)
O1—C11—C1	118.64 (11)	H13—C25—H14	113.3 (16)
C12—C11—C1	120.42 (9)	O3—C25—H15	111.1 (11)
C17—C12—C13	118.93 (11)	H13—C25—H15	108.7 (15)
C17—C12—C11	122.58 (10)	H14—C25—H15	109.0 (15)
C13—C12—C11	118.41 (10)	O4—C26—H16	110.5 (10)
C14—C13—C12	120.87 (12)	O4—C26—H17	105.2 (11)
C14—C13—H5	121.7	H16—C26—H17	113.2 (15)
C12—C13—H5	117.4	O4—C26—H18	110.6 (10)
C15—C14—C13	118.15 (12)	H16—C26—H18	108.1 (14)
C15—C14—H6	119.6	H17—C26—H18	109.3 (16)
C25—O3—C2—C1	-175.20 (12)	C10—C1—C11—O1	-68.78 (15)
C25—O3—C2—C3	3.86 (19)	C2—C1—C11—C12	-79.42 (14)
C10—C1—C2—O3	176.46 (10)	C10—C1—C11—C12	109.20 (12)
C11—C1—C2—O3	4.95 (16)	O1—C11—C12—C17	-168.99 (12)
C10—C1—C2—C3	-2.62 (18)	C1—C11—C12—C17	13.08 (17)
C11—C1—C2—C3	-174.13 (11)	O1—C11—C12—C13	14.35 (17)

O3—C2—C3—C4	-177.06 (12)	C1—C11—C12—C13	-163.58 (11)
C1—C2—C3—C4	2.0 (2)	C17—C12—C13—C14	-0.5 (2)
C2—C3—C4—C5	0.9 (2)	C11—C12—C13—C14	176.26 (12)
C3—C4—C5—C6	175.24 (12)	C12—C13—C14—C15	0.3 (2)
C3—C4—C5—C10	-2.9 (2)	C13—C14—C15—F1	-178.54 (13)
C4—C5—C6—C7	-176.63 (13)	C13—C14—C15—C16	0.4 (2)
C10—C5—C6—C7	1.52 (19)	F1—C15—C16—C17	178.04 (13)
C5—C6—C7—C8	-3.0 (2)	C14—C15—C16—C17	-0.9 (2)
C26—O4—C8—C9	-176.00 (12)	C15—C16—C17—C12	0.7 (2)
C26—O4—C8—C7	0.9 (2)	C13—C12—C17—C16	0.01 (19)
C6—C7—C8—O4	-175.71 (12)	C11—C12—C17—C16	-176.63 (12)
C6—C7—C8—C9	1.0 (2)	C8—C9—C18—O2	111.97 (13)
O4—C8—C9—C10	179.36 (10)	C10—C9—C18—O2	-57.38 (16)
C7—C8—C9—C10	2.44 (18)	C8—C9—C18—C19	-65.30 (14)
O4—C8—C9—C18	9.61 (16)	C10—C9—C18—C19	125.35 (12)
C7—C8—C9—C18	-167.31 (11)	O2—C18—C19—C24	150.27 (12)
C2—C1—C10—C9	-177.21 (11)	C9—C18—C19—C24	-32.49 (17)
C11—C1—C10—C9	-6.01 (17)	O2—C18—C19—C20	-28.41 (17)
C2—C1—C10—C5	0.53 (16)	C9—C18—C19—C20	148.83 (12)
C11—C1—C10—C5	171.73 (10)	C24—C19—C20—C21	0.4 (2)
C8—C9—C10—C1	173.95 (11)	C18—C19—C20—C21	179.11 (12)
C18—C9—C10—C1	-17.11 (17)	C19—C20—C21—C22	-0.9 (2)
C8—C9—C10—C5	-3.79 (16)	C20—C21—C22—F2	178.54 (12)
C18—C9—C10—C5	165.16 (11)	C20—C21—C22—C23	0.4 (2)
C6—C5—C10—C1	-176.01 (10)	F2—C22—C23—C24	-177.46 (12)
C4—C5—C10—C1	2.16 (16)	C21—C22—C23—C24	0.7 (2)
C6—C5—C10—C9	1.88 (16)	C22—C23—C24—C19	-1.2 (2)
C4—C5—C10—C9	-179.95 (11)	C20—C19—C24—C23	0.70 (19)
C2—C1—C11—O1	102.61 (13)	C18—C19—C24—C23	-177.97 (12)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
C16—H7···O1 <sup>i</sup>	1.00	2.54	3.493 (2)	158
C21—H10···O1 <sup>ii</sup>	0.99	2.68	3.636 (2)	161

Symmetry codes: (i)  $x, -y+1/2, z-1/2$ ; (ii)  $x+1, -y+1/2, z+1/2$ .