

Tris{aquabis[3-(2-pyridyl)-1H-pyrazole]-copper(II)} di- μ_9 -arsenato-hexatriaconta- μ_2 -oxido-octadecaoxidoctadeca-molybdate(VI)

Xutang Zhang,^{a,b*} Peihai Wei,^a Congwen Shi,^a Bin Li^a and Bo Hu^a

^aAdvanced Material Institute of Research, Department of Chemistry and Chemical Engineering, ShanDong Institute of Education, Jinan 250013, People's Republic of China, and ^bCollege of Chemistry and Chemical Engineering, LiaoCheng University, LiaoCheng 252059, People's Republic of China

Correspondence e-mail: xiutangzhang@yahoo.com.cn


Received 9 December 2009; accepted 13 January 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.018\text{ \AA}$; disorder in main residue; R factor = 0.046; wR factor = 0.123; data-to-parameter ratio = 10.8.

The title compound, $[\text{Cu}(\text{C}_8\text{H}_7\text{N}_3)_2(\text{H}_2\text{O})]_3[\text{As}_2\text{Mo}_{18}\text{O}_{62}]$, consists of two subunits, *viz.* an α -Dawson-type $[\text{As}_2\text{Mo}_{18}\text{O}_{62}]^{6-}$ anion and a complex $[\text{Cu}(\text{C}_8\text{H}_7\text{N}_3)_2(\text{H}_2\text{O})]^{2+}$ cation. The copper(II) ion (site symmetry .2) is pentacoordinated in a distorted square-pyramidal manner by four N atoms from two chelating 3-(2-pyridyl)pyrazole ligands in equatorial positions and one water molecule in the apical position. In the heteropolyanion, two O atoms of the AsO_4 group (3. symmetry) are equally disordered about the threefold rotation axis. N—H···O and O—H···O hydrogen bonding between the neutral molecules and the water molecules leads to a consolidation of the structure.

Related literature

For background to polyoxometalates, see: Pope & Müller (1991). For polyoxometalates modified with amines, see: Zhang, Dou *et al.* (2009); Zhang, Wei, Shi *et al.* (2010); Zhang, Wei *et al.* (2009); Zhang, Yuan *et al.* (2010). Zhang, Wei, Zhu *et al.* (2010). For another α -Dawson-type anion, see: Li *et al.* (2007).



Experimental

Crystal data

$[\text{Cu}(\text{C}_8\text{H}_7\text{N}_3)_2(\text{H}_2\text{O})]_3[\text{As}_2\text{Mo}_{18}\text{O}_{62}]$	$Z = 6$
$M_r = 3984.45$	Mo $K\alpha$ radiation
Hexagonal, $R\bar{3}c$	$\mu = 3.72\text{ mm}^{-1}$
$a = 21.967 (3)\text{ \AA}$	$T = 293\text{ K}$
$c = 34.411 (7)\text{ \AA}$	$0.12 \times 0.10 \times 0.08\text{ mm}$
$V = 14380 (4)\text{ \AA}^3$	

Data collection

Bruker APEXII CCD diffractometer	25458 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	2750 independent reflections
$(SADABS$; Bruker, 2001)	2053 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.664$, $T_{\max} = 0.755$	$R_{\text{int}} = 0.085$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.123$	$\Delta\rho_{\max} = 2.25\text{ e \AA}^{-3}$
$S = 1.00$	$\Delta\rho_{\min} = -1.15\text{ e \AA}^{-3}$
2750 reflections	
254 parameters	
14 restraints	

Table 1
Selected bond lengths (\AA).

As1—O10A	1.653 (10)	Cu1—N3	1.984 (7)
As1—O10B	1.677 (10)	Cu1—N1	1.985 (8)
As1—O1	1.728 (9)	Cu1—O11	2.29 (2)

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N2—H2A···O6 ⁱ	0.86	2.27	3.097 (13)	162
O11—H1W···O3 ⁱⁱ	0.84 (8)	2.69 (11)	2.860 (10)	94 (8)

Symmetry codes: (i) $x + \frac{1}{3}$, $x - y + \frac{2}{3}$, $z + \frac{1}{6}$; (ii) y , x , $-z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT-Plus* (Bruker, 2001); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Financial support from the Chinese Academy of Sciences ('Hundred Talents Program') and the Ministry of Science and Technology of China (grant No. 2007CB607608), Shandong Provincial Education Department and Shandong Institute of Education are gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2290).

References

- Bruker (2001). *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Li, F. Y., Qu, X. S. & Qiu, Y. F. (2007). *Cryst. Res. Technol.* **42**, 1036–1043.
- Pope, M. T. & Müller, A. (1991). *Angew. Chem. Int. Ed. Engl.* **30**, 34–38.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zhang, X. T., Dou, J. M., Wei, P. H., Li, D. C., Li, B., Shi, C. W. & Hu, B. (2009). *Inorg. Chim. Acta*, **362**, 3325–3332.
- Zhang, X., Wei, P., Shi, C., Li, B. & Hu, B. (2010). *Acta Cryst. E* **66**, m26–m27.
- Zhang, X. T., Wei, P. H., Sun, D. F., Ni, Z. H., Dou, J. M., Li, B., Shi, C. W. & Hu, B. (2009). *Cryst. Growth Des.* **9**, 4424–4428.
- Zhang, X., Wei, P., Zhu, W., Li, B. & Hu, B. (2010). *Acta Cryst. E* **66**, m127–m128.
- Zhang, X., Yuan, D., Wei, P., Li, B. & Hu, B. (2010). *Acta Cryst. E* **66**, m152–m153.

supporting information

Acta Cryst. (2010). E66, m174–m175 [https://doi.org/10.1107/S160053681000156X]

Tris{aquabis[3-(2-pyridyl)-1H-pyrazole]copper(II)} di- μ_9 -arsenato-hexatriaconta- μ_2 -oxido-octadecaoxidoctadecamolybdate(VI)

Xiutang Zhang, Peihai Wei, Congwen Shi, Bin Li and Bo Hu

S1. Comment

The design and synthesis of polyoxometalates has attracted continuous research interest not only because of their appealing structural and topological novelties, but also due to their interesting optical, electronic, magnetic, and catalytic properties, as well as their potential medical applications (Pope *et al.*, 1991). In our group, organic amines, such as 3-(2-pyridyl)pyrazole and pyrazine, are used to effectively modify polyoxomolybdates under hydrothermal condicitions (Zhang, Dou *et al.*, 2009; Zhang, Wei *et al.*, 2009). Here, we describe the synthesis and structural characterization of the title compound.

As shown in Figure 1, the title compound consists of two subunits, *viz.* of an α -Dawson-type $[\text{As}_2\text{Mo}_{18}\text{O}_{62}]^{6-}$ anion (Li *et al.*, 2007) and a complex $[\text{Cu}(\text{H}_2\text{O})(\text{C}_8\text{H}_7\text{N}_3)_2]^{2+}$ cation. The copper ion is penta-coordinated in a distorted square-pyramidal geometry by four N atoms from two 3-(2-pyridyl)pyrazole ligands and by one water molecule. The Cu—N bond lengths are in the range of 1.984 (7)—1.985 (8) Å and the Cu—O bond length is 2.29 (2) Å. In the heteropolyanion, there are four kinds of oxygen atoms according to their coordination manner: (i) 18 terminal O atoms bonded to one Mo atom with their Mo—O distances in the range of 1.651 (6)—1.690 (6) Å; (ii) 36 μ_2 O atoms, the Mo—O distances in the range of 1.797 (5)—2.117 (5) Å; (iii) six μ_3 O atoms shared by one As and two Mo atoms, the Mo—O distances varying from 1.653 (8) to 2.359 (1) Å; (iv) two μ_4 O atoms which are coordinated to one As atom and three Mo atoms, Mo—O distances are between 1.728 (7) and 2.341 (7) Å, respectively. The resulting MoO_6 octahedra are considerably distorted. The AsO_4 group is disordered about a threefold rotation axis and exhibits two sets of short As—O bond lenghts to the disordered O atoms (50% occupation) and one longer As—O bond. N—H···O and O—H···O hydrogen bonding between the neutral molecules and the water molecules leads to a consolidation of the structure (Fig. 2; Table 2) which also contains accessible voids of ca. 136 Å³.

S2. Experimental

A mixture of 3-(2-pyridyl)pyrazole (1 mmol, 0.14 g), sodium molybdate (2 mmol, 0.48 g), sodium arsenate (0.2 mmol, 0.08 g) and copper dichloride dihydrate (1 mmol, 0.28 g) in 14 ml distilled water was sealed in a 25 ml Teflon-lined stainless steel autoclave and was kept at 433 K for three days. Blue crystals suitable for the X-ray experiment were obtained. Anal. Calc. for $\text{C}_{48}\text{H}_{48}\text{As}_2\text{Cu}_3\text{Mo}_{18}\text{N}_{18}\text{O}_{65}$: C 14.46, H 1.20, N 6.32%; Found: C 14.24, H 1.01, N 6.23%.

S3. Refinement

All hydrogen atoms bound to carbon were refined using a riding model with distance C—H = 0.93 Å, $U_{\text{iso}} = 1.2U_{\text{eq}}$ (C) for aromatic atoms. The H atoms of the water molecule were located from difference density maps and were refined with d(O—H) = 0.83 (2) Å, and with a fixed U_{iso} of 0.80 Å². In the AsO_4 unit, two oxygen atoms (O6 and O10) are disordered around a threefold rotation axis. Both positions were refined with split positions and an occupancy ratio of 1:1. In the

final difference Fourier map the highest peak is 3.20 Å from atom O2 and the deepest hole is 0.67 Å from atom O11. The highest peak is located in the voids of the crystal structure and may be associated with an additional water molecule. However, refinement of this position did not result in a reasonable model. Hence this position was excluded from the final refinement.




Figure 1

The cation and anion of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level; H atoms are given as spheres of arbitrary radius.




Figure 2

The crystal packing of the title compound, displayed with hydrogen bonds as dashed lines.

Tris{aquabis[3-(2-pyridyl)-1*H*-pyrazole]copper(II)} di- μ_9 -arsenato-hexatriaconta- μ_2 -oxido-octadecaoxidoctadecamolybdate(VI)

Crystal data

[Cu(C₈H₇N₃)₂(H₂O)]₃[As₂Mo₁₈O₆₂]

$M_r = 3984.45$

Hexagonal, $R\bar{3}c$

Hall symbol: -R 3 2 "c

$a = 21.967 (3) \text{ \AA}$

$c = 34.411 (7) \text{ \AA}$

$V = 14380 (4) \text{ \AA}^3$

$Z = 6$

$F(000) = 11346$

$D_x = 2.761 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2750 reflections

$\theta = 1.6\text{--}25.0^\circ$

$\mu = 3.72 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, blue

$0.12 \times 0.10 \times 0.08 \text{ mm}$

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2001)

$T_{\min} = 0.664$, $T_{\max} = 0.755$

25458 measured reflections

2750 independent reflections

2053 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.085$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.6^\circ$

$h = -25 \rightarrow 26$

$k = -26 \rightarrow 26$

$l = -40 \rightarrow 39$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.046$

$wR(F^2) = 0.123$

$S = 1.00$

2750 reflections

254 parameters

14 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.063P)^2 + 278.6957P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 2.25 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -1.15 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^* / U_{\text{eq}}$	Occ. (<1)
As1	0.0000	0.0000	0.05884 (4)	0.0161 (3)	
C1	0.2641 (5)	0.2974 (5)	0.1696 (3)	0.039 (2)	
C2	0.1872 (5)	0.3163 (5)	0.2056 (3)	0.042 (2)	

H2	0.1698	0.3199	0.2296	0.050*	
C3	0.1552 (7)	0.3216 (5)	0.1730 (3)	0.055 (3)	
H3	0.1192	0.3321	0.1747	0.066*	
C4	0.1768 (7)	0.3112 (5)	0.1380 (3)	0.055 (3)	
H4	0.1534	0.3113	0.1155	0.065*	
C5	0.2326 (6)	0.3008 (5)	0.1359 (3)	0.042 (3)	
H5	0.2493	0.2960	0.1119	0.050*	
C6	0.2831 (5)	0.3216 (5)	0.3283 (2)	0.037 (2)	
C7	0.2761 (7)	0.3645 (7)	0.3554 (4)	0.066 (3)	
H7	0.2809	0.3636	0.3822	0.079*	
C8	0.2605 (7)	0.4092 (6)	0.3343 (3)	0.067 (4)	
H8	0.2523	0.4438	0.3442	0.080*	
Cu1	0.28491 (9)	0.28491 (9)	0.2500	0.0444 (5)	
Mo1	0.06070 (4)	0.10373 (4)	0.14523 (2)	0.0293 (2)	
Mo2	0.03113 (4)	0.17477 (4)	0.05152 (2)	0.0270 (2)	
Mo3	0.17292 (4)	0.14366 (4)	0.05871 (2)	0.0298 (2)	
N1	0.2731 (5)	0.3393 (5)	0.2924 (2)	0.045 (2)	
N2	0.2599 (5)	0.3921 (5)	0.2969 (3)	0.057 (3)	
H2A	0.2518	0.4126	0.2779	0.068*	
N3	0.2426 (4)	0.3063 (4)	0.2048 (2)	0.0381 (19)	
O1	0.0000	0.0000	0.1090 (3)	0.023 (2)	
O2	-0.0297 (3)	0.0513 (3)	0.16857 (15)	0.0250 (12)	
O3	0.1016 (3)	0.1636 (3)	0.18082 (17)	0.0336 (15)	
O4	0.0331 (3)	0.1452 (3)	0.10985 (16)	0.0261 (13)	
O5	0.1395 (3)	0.1213 (3)	0.11134 (16)	0.0271 (13)	
O6	-0.0665 (5)	0.1212 (5)	0.0577 (3)	0.075 (2)	
O7	0.0438 (3)	0.2531 (3)	0.06422 (17)	0.0344 (15)	
O8	0.2534 (3)	0.2085 (3)	0.06733 (18)	0.0412 (17)	
O9	0.0315 (4)	0.1816 (3)	-0.00079 (17)	0.0445 (18)	
O10A	-0.0288 (5)	0.0530 (5)	0.0445 (3)	0.019 (2)	0.50
O10B	-0.0832 (5)	-0.0297 (5)	0.0446 (3)	0.016 (2)	0.50
O11	0.1808 (10)	0.1808 (10)	0.2500	0.142 (2)	
O12A	0.1344 (7)	0.2076 (7)	0.0591 (4)	0.030 (3)	0.50
O12B	0.1111 (7)	0.1708 (8)	0.0481 (4)	0.027 (3)	0.50
H1W	0.1487 (19)	0.179 (2)	0.264 (3)	0.080*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
As1	0.0173 (4)	0.0173 (4)	0.0136 (7)	0.0086 (2)	0.000	0.000
C1	0.043 (6)	0.025 (5)	0.034 (6)	0.005 (4)	-0.004 (4)	0.000 (4)
C2	0.058 (7)	0.042 (6)	0.032 (6)	0.030 (5)	-0.006 (5)	-0.003 (4)
C3	0.081 (9)	0.041 (6)	0.051 (7)	0.038 (6)	-0.015 (6)	-0.005 (5)
C4	0.082 (9)	0.030 (6)	0.055 (8)	0.031 (6)	-0.018 (6)	-0.004 (5)
C5	0.063 (7)	0.032 (5)	0.020 (5)	0.016 (5)	-0.011 (4)	-0.002 (4)
C6	0.045 (6)	0.044 (6)	0.018 (5)	0.019 (5)	0.004 (4)	0.000 (4)
C7	0.076 (9)	0.065 (8)	0.042 (7)	0.025 (7)	0.010 (6)	0.006 (6)
C8	0.099 (10)	0.058 (8)	0.040 (7)	0.036 (7)	0.021 (7)	-0.002 (6)

Cu1	0.0590 (9)	0.0590 (9)	0.0215 (9)	0.0344 (10)	-0.0019 (4)	0.0019 (4)
Mo1	0.0391 (5)	0.0246 (4)	0.0214 (4)	0.0139 (4)	-0.0087 (3)	-0.0058 (3)
Mo2	0.0362 (5)	0.0200 (4)	0.0270 (4)	0.0156 (3)	0.0013 (3)	0.0008 (3)
Mo3	0.0201 (4)	0.0244 (4)	0.0388 (5)	0.0067 (3)	-0.0063 (3)	-0.0045 (3)
N1	0.065 (6)	0.048 (5)	0.022 (4)	0.028 (5)	0.000 (4)	-0.001 (4)
N2	0.086 (7)	0.041 (5)	0.044 (6)	0.033 (5)	0.007 (5)	0.016 (4)
N3	0.048 (5)	0.028 (4)	0.025 (4)	0.009 (4)	-0.007 (4)	0.003 (3)
O1	0.027 (3)	0.027 (3)	0.016 (5)	0.0133 (16)	0.000	0.000
O2	0.028 (3)	0.029 (3)	0.016 (3)	0.013 (3)	0.003 (2)	-0.001 (2)
O3	0.044 (4)	0.028 (3)	0.023 (3)	0.013 (3)	-0.008 (3)	-0.006 (3)
O4	0.028 (3)	0.027 (3)	0.025 (3)	0.015 (3)	0.006 (2)	0.005 (2)
O5	0.030 (3)	0.030 (3)	0.024 (3)	0.017 (3)	0.001 (2)	0.001 (2)
O6	0.074 (3)	0.075 (3)	0.073 (3)	0.0349 (15)	-0.0029 (10)	0.0033 (10)
O7	0.048 (4)	0.025 (3)	0.030 (3)	0.018 (3)	0.001 (3)	-0.001 (3)
O8	0.024 (3)	0.048 (4)	0.035 (4)	0.005 (3)	-0.002 (3)	-0.014 (3)
O9	0.087 (5)	0.030 (4)	0.022 (3)	0.034 (4)	0.001 (3)	0.002 (3)
O10A	0.015 (5)	0.022 (6)	0.020 (6)	0.009 (5)	0.003 (4)	0.001 (4)
O10B	0.017 (5)	0.012 (5)	0.020 (6)	0.009 (4)	0.009 (4)	0.001 (4)
O11	0.142 (2)	0.142 (2)	0.142 (2)	0.0707 (13)	-0.0005 (7)	0.0005 (7)
O12A	0.029 (8)	0.023 (8)	0.037 (9)	0.012 (7)	0.001 (6)	0.001 (6)
O12B	0.028 (8)	0.029 (8)	0.022 (7)	0.012 (7)	0.004 (6)	-0.005 (6)

Geometric parameters (\AA , $^\circ$)

As1—O10A ⁱ	1.653 (10)	Mo1—O1	2.341 (5)
As1—O10A	1.653 (10)	Mo2—O7	1.658 (6)
As1—O10A ⁱⁱ	1.653 (10)	Mo2—O12B	1.806 (13)
As1—O10B	1.677 (10)	Mo2—O9	1.806 (6)
As1—O10B ⁱ	1.677 (10)	Mo2—O6	1.872 (9)
As1—O10B ⁱⁱ	1.677 (10)	Mo2—O12A	2.024 (13)
As1—O1	1.728 (9)	Mo2—O4	2.117 (5)
C1—N3	1.350 (12)	Mo2—O10B ⁱⁱ	2.309 (9)
C1—C5	1.371 (13)	Mo2—O10A	2.330 (10)
C1—C6 ⁱⁱⁱ	1.447 (14)	Mo3—O8	1.651 (6)
C2—N3	1.342 (12)	Mo3—O12B	1.772 (13)
C2—C3	1.359 (14)	Mo3—O6 ⁱⁱ	1.878 (9)
C2—H2	0.9300	Mo3—O5	1.923 (6)
C3—C4	1.355 (16)	Mo3—O12A	1.970 (13)
C3—H3	0.9300	Mo3—O9 ^{iv}	2.000 (6)
C4—C5	1.358 (15)	Mo3—O10B ⁱⁱ	2.325 (9)
C4—H4	0.9300	Mo3—O10A ⁱⁱ	2.359 (10)
C5—H5	0.9300	N1—N2	1.337 (12)
C6—N1	1.345 (11)	N2—H2A	0.8600
C6—C7	1.389 (15)	O1—Mo1 ⁱ	2.341 (5)
C6—C1 ⁱⁱⁱ	1.447 (14)	O1—Mo1 ⁱⁱ	2.341 (5)
C7—C8	1.395 (17)	O2—Mo1 ⁱ	2.053 (5)
C7—H7	0.9300	O6—Mo3 ⁱ	1.878 (9)
C8—N2	1.339 (13)	O9—Mo3 ^v	2.000 (6)

C8—H8	0.9300	O10A—O10B ⁱⁱ	1.582 (13)
Cu1—N3 ⁱⁱⁱ	1.984 (7)	O10A—O10B	1.599 (13)
Cu1—N3	1.984 (7)	O10A—Mo3 ⁱ	2.359 (10)
Cu1—N1	1.985 (8)	O10B—O10A ⁱ	1.582 (13)
Cu1—N1 ⁱⁱⁱ	1.985 (8)	O10B—O12B ⁱ	1.698 (17)
Cu1—O11	2.29 (2)	O10B—Mo2 ⁱ	2.309 (9)
Mo1—O3	1.690 (6)	O10B—Mo3 ⁱ	2.325 (9)
Mo1—O4	1.797 (5)	O11—H1W	0.84 (8)
Mo1—O2	1.904 (5)	O12A—O12B	0.804 (14)
Mo1—O5	1.959 (6)	O12B—O10B ⁱⁱ	1.698 (17)
Mo1—O2 ⁱⁱ	2.053 (5)		
O10A ⁱ —As1—O10A	111.5 (3)	O7—Mo2—O10A	157.6 (3)
O10A ⁱ —As1—O10A ⁱⁱ	111.5 (3)	O12B—Mo2—O10A	86.7 (5)
O10A—As1—O10A ⁱⁱ	111.5 (3)	O9—Mo2—O10A	88.2 (3)
O10A ⁱ —As1—O10B	56.7 (4)	O6—Mo2—O10A	58.5 (4)
O10A—As1—O10B	57.4 (4)	O12A—Mo2—O10A	108.2 (5)
O10A ⁱⁱ —As1—O10B	145.7 (5)	O4—Mo2—O10A	80.3 (3)
O10A ⁱ —As1—O10B ⁱ	57.4 (4)	O10B ⁱⁱ —Mo2—O10A	39.9 (3)
O10A—As1—O10B ⁱ	145.7 (5)	O8—Mo3—O12B	114.7 (5)
O10A ⁱⁱ —As1—O10B ⁱ	56.7 (4)	O8—Mo3—O6 ⁱⁱ	100.9 (4)
O10B—As1—O10B ⁱ	111.8 (3)	O12B—Mo3—O6 ⁱⁱ	143.7 (6)
O10A ⁱ —As1—O10B ⁱⁱ	145.7 (5)	O8—Mo3—O5	99.1 (3)
O10A—As1—O10B ⁱⁱ	56.7 (4)	O12B—Mo3—O5	91.2 (4)
O10A ⁱⁱ —As1—O10B ⁱⁱ	57.4 (4)	O6 ⁱⁱ —Mo3—O5	90.1 (3)
O10B—As1—O10B ⁱⁱ	111.8 (3)	O8—Mo3—O12A	92.3 (5)
O10B ⁱ —As1—O10B ⁱⁱ	111.8 (3)	O12B—Mo3—O12A	24.1 (4)
O10A ⁱ —As1—O1	107.4 (4)	O6 ⁱⁱ —Mo3—O12A	166.7 (5)
O10A—As1—O1	107.4 (4)	O5—Mo3—O12A	86.0 (4)
O10A ⁱⁱ —As1—O1	107.4 (4)	O8—Mo3—O9 ^{iv}	95.5 (3)
O10B—As1—O1	107.0 (3)	O12B—Mo3—O9 ^{iv}	80.4 (5)
O10B ⁱ —As1—O1	107.0 (3)	O6 ⁱⁱ —Mo3—O9 ^{iv}	89.6 (4)
O10B ⁱⁱ —As1—O1	107.0 (3)	O5—Mo3—O9 ^{iv}	165.2 (2)
N3—C1—C5	122.0 (10)	O12A—Mo3—O9 ^{iv}	90.9 (5)
N3—C1—C6 ⁱⁱⁱ	113.1 (8)	O8—Mo3—O10B ⁱⁱ	161.3 (3)
C5—C1—C6 ⁱⁱⁱ	124.9 (9)	O12B—Mo3—O10B ⁱⁱ	46.6 (5)
N3—C2—C3	123.3 (10)	O6 ⁱⁱ —Mo3—O10B ⁱⁱ	97.8 (4)
N3—C2—H2	118.4	O5—Mo3—O10B ⁱⁱ	82.7 (3)
C3—C2—H2	118.4	O12A—Mo3—O10B ⁱⁱ	69.1 (5)
C4—C3—C2	118.6 (11)	O9 ^{iv} —Mo3—O10B ⁱⁱ	82.7 (3)
C4—C3—H3	120.7	O8—Mo3—O10A ⁱⁱ	158.5 (3)
C2—C3—H3	120.7	O12B—Mo3—O10A ⁱⁱ	86.2 (5)
C3—C4—C5	119.9 (11)	O6 ⁱⁱ —Mo3—O10A ⁱⁱ	57.9 (4)
C3—C4—H4	120.1	O5—Mo3—O10A ⁱⁱ	84.9 (3)
C5—C4—H4	120.1	O12A—Mo3—O10A ⁱⁱ	109.0 (5)
C4—C5—C1	119.1 (10)	O9 ^{iv} —Mo3—O10A ⁱⁱ	82.5 (3)
C4—C5—H5	120.4	O10B ⁱⁱ —Mo3—O10A ⁱⁱ	39.9 (3)
C1—C5—H5	120.4	N2—N1—C6	106.6 (8)

N1—C6—C7	109.3 (10)	N2—N1—Cu1	139.1 (7)
N1—C6—C1 ⁱⁱⁱ	115.9 (8)	C6—N1—Cu1	114.2 (7)
C7—C6—C1 ⁱⁱⁱ	134.8 (9)	N1—N2—C8	112.1 (9)
C6—C7—C8	106.0 (10)	N1—N2—H2A	124.0
C6—C7—H7	127.0	C8—N2—H2A	124.0
C8—C7—H7	127.0	C2—N3—C1	116.9 (8)
N2—C8—C7	106.1 (11)	C2—N3—Cu1	126.5 (7)
N2—C8—H8	127.0	C1—N3—Cu1	115.5 (7)
C7—C8—H8	127.0	As1—O1—Mo1 ⁱ	122.13 (18)
N3 ⁱⁱⁱ —Cu1—N3	166.7 (4)	As1—O1—Mo1	122.13 (18)
N3 ⁱⁱⁱ —Cu1—N1	80.7 (3)	Mo1 ⁱ —O1—Mo1	94.3 (2)
N3—Cu1—N1	102.5 (3)	As1—O1—Mo1 ⁱⁱ	122.13 (18)
N3 ⁱⁱⁱ —Cu1—N1 ⁱⁱⁱ	102.5 (3)	Mo1 ⁱ —O1—Mo1 ⁱⁱ	94.3 (2)
N3—Cu1—N1 ⁱⁱⁱ	80.7 (3)	Mo1—O1—Mo1 ⁱⁱ	94.3 (2)
N1—Cu1—N1 ⁱⁱⁱ	152.7 (5)	Mo1—O2—Mo1 ⁱ	120.4 (3)
N3 ⁱⁱⁱ —Cu1—O11	83.4 (2)	Mo1—O4—Mo2	149.5 (3)
N3—Cu1—O11	83.4 (2)	Mo3—O5—Mo1	142.9 (3)
N1—Cu1—O11	103.6 (3)	Mo2—O6—Mo3 ⁱ	144.3 (5)
N1 ⁱⁱⁱ —Cu1—O11	103.6 (3)	Mo2—O9—Mo3 ^v	170.5 (4)
O3—Mo1—O4	106.2 (3)	O10B ⁱⁱ —O10A—O10B	121.6 (9)
O3—Mo1—O2	98.8 (3)	O10B ⁱⁱ —O10A—As1	62.4 (5)
O4—Mo1—O2	94.5 (2)	O10B—O10A—As1	62.0 (5)
O3—Mo1—O5	101.8 (3)	O10B ⁱⁱ —O10A—Mo2	69.3 (5)
O4—Mo1—O5	89.2 (2)	O10B—O10A—Mo2	167.6 (7)
O2—Mo1—O5	157.1 (2)	As1—O10A—Mo2	125.6 (5)
O3—Mo1—O2 ⁱⁱ	95.3 (2)	O10B ⁱⁱ —O10A—Mo3 ⁱ	164.3 (7)
O4—Mo1—O2 ⁱⁱ	158.1 (2)	O10B—O10A—Mo3 ⁱ	68.9 (5)
O2—Mo1—O2 ⁱⁱ	85.9 (3)	As1—O10A—Mo3 ⁱ	121.8 (5)
O5—Mo1—O2 ⁱⁱ	82.5 (2)	Mo2—O10A—Mo3 ⁱ	99.2 (4)
O3—Mo1—O1	164.9 (3)	O10A ⁱ —O10B—O10A	118.4 (9)
O4—Mo1—O1	87.7 (2)	O10A ⁱ —O10B—As1	60.9 (5)
O2—Mo1—O1	73.8 (2)	O10A—O10B—As1	60.5 (5)
O5—Mo1—O1	83.8 (2)	O10A ⁱ —O10B—O12B ⁱ	121.6 (8)
O2 ⁱⁱ —Mo1—O1	71.36 (19)	O10A—O10B—O12B ⁱ	119.9 (8)
O7—Mo2—O12B	113.9 (5)	As1—O10B—O12B ⁱ	159.0 (8)
O7—Mo2—O9	100.7 (3)	O10A ⁱ —O10B—Mo2 ⁱ	70.8 (5)
O12B—Mo2—O9	88.5 (5)	O10A—O10B—Mo2 ⁱ	169.3 (7)
O7—Mo2—O6	99.7 (4)	As1—O10B—Mo2 ⁱ	125.6 (5)
O12B—Mo2—O6	144.4 (5)	O12B ⁱ —O10B—Mo2 ⁱ	50.8 (5)
O9—Mo2—O6	97.0 (4)	O10A ⁱ —O10B—Mo3 ⁱ	165.0 (7)
O7—Mo2—O12A	91.1 (5)	O10A—O10B—Mo3 ⁱ	71.2 (5)
O12B—Mo2—O12A	23.4 (4)	As1—O10B—Mo3 ⁱ	122.5 (5)
O9—Mo2—O12A	98.1 (5)	O12B ⁱ —O10B—Mo3 ⁱ	49.3 (5)
O6—Mo2—O12A	159.5 (5)	Mo2 ⁱ —O10B—Mo3 ⁱ	98.7 (3)
O7—Mo2—O4	92.9 (3)	Cu1—O11—H1W	117 (2)
O12B—Mo2—O4	83.0 (4)	O12B—O12A—Mo3	64.0 (14)
O9—Mo2—O4	166.0 (2)	O12B—O12A—Mo2	63.0 (14)
O6—Mo2—O4	83.8 (3)	Mo3—O12A—Mo2	123.3 (7)

O12A—Mo2—O4	78.2 (4)	O12A—O12B—O10B ⁱⁱ	155.9 (19)
O7—Mo2—O10B ⁱⁱ	158.9 (3)	O12A—O12B—Mo3	91.9 (16)
O12B—Mo2—O10B ⁱⁱ	46.8 (5)	O10B ⁱⁱ —O12B—Mo3	84.1 (7)
O9—Mo2—O10B ⁱⁱ	88.6 (3)	O12A—O12B—Mo2	93.7 (16)
O6—Mo2—O10B ⁱⁱ	98.0 (4)	O10B ⁱⁱ —O12B—Mo2	82.4 (7)
O12A—Mo2—O10B ⁱⁱ	68.6 (5)	Mo3—O12B—Mo2	158.4 (8)
O4—Mo2—O10B ⁱⁱ	77.5 (3)		

Symmetry codes: (i) $-y, x-y, z$; (ii) $-x+y, -x, z$; (iii) $y, x, -z+1/2$; (iv) $y, -x+y, -z$; (v) $x-y, x, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2A \cdots O6 ^{vi}	0.86	2.27	3.097 (13)	162
O11—H1W \cdots O3 ⁱⁱⁱ	0.84 (8)	2.69 (11)	2.860 (10)	94 (8)

Symmetry codes: (iii) $y, x, -z+1/2$; (vi) $x+1/3, x-y+2/3, z+1/6$.