

6-Bromo-1,3-di-2-propynyl-1*H*-imidazo-[4,5-*b*]pyridin-2(3*H*)-one

S. Dahmani,^a A. Haoudi,^a F. Capet,^b El Mokhtar Essassi^c
and Seik Weng Ng^{d*}

^aLaboratoire de Chimie Organique Appliquée, Faculté des Sciences et Techniques, Université Sidi Mohamed Ben Abdallah, Fès, Morocco, ^bUnité de Catalyse et de Chimie du Solide, Ecole Nationale Supérieure de Chimie de Lille, Lille, France,

^cLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, and ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 22 February 2010; accepted 1 March 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.029; wR factor = 0.085; data-to-parameter ratio = 20.9.

The room-temperature reaction of propargyl bromide and 6-bromo-1,3-dihydroimidazo[4,5-*b*]pyridin-2-one in dimethylformamide yields the title compound, $C_{12}H_8\text{BrN}_3\text{O}$, which features nitrogen-bound propynyl substituents. The imidazopyridine fused ring is almost planar (r.m.s. deviation = 0.011 Å); the propynyl chains point in opposite directions relative to the fused ring. One acetylenic H atom is hydrogen bonded to the carbonyl O atom of an inversion-related molecule, forming a dimer; adjacent dimers are linked by a second acetylene–pyridine C–H···N interaction, forming a layer motif.

Related literature

For the crystal structures of other imidazo[4,5-*b*]pyridin-2-ones, see: Kourafalos *et al.* (2002); Meanwell *et al.* (1995).



Experimental

Crystal data

$C_{12}H_8\text{BrN}_3\text{O}$	$V = 1163.56 (6)\text{ \AA}^3$
$M_r = 290.12$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 9.0725 (3)\text{ \AA}$	$\mu = 3.52\text{ mm}^{-1}$
$b = 18.6212 (5)\text{ \AA}$	$T = 293\text{ K}$
$c = 7.0684 (2)\text{ \AA}$	$0.35 \times 0.30 \times 0.15\text{ mm}$
$\beta = 102.995 (1)^\circ$	

Data collection

Bruker X8 APEXII diffractometer	27315 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	3383 independent reflections
($SADABS$; Sheldrick, 1996)	2810 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.372$, $T_{\max} = 0.620$	$R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.085$	$\Delta\rho_{\max} = 0.87\text{ e \AA}^{-3}$
$S = 1.03$	$\Delta\rho_{\min} = -0.80\text{ e \AA}^{-3}$
3383 reflections	
162 parameters	
2 restraints	

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$C8-\text{H}8\cdots\text{O}1^{\text{i}}$	0.95 (1)	2.53 (2)	3.392 (3)	151 (3)
$C12-\text{H}12\cdots\text{N}1^{\text{ii}}$	0.94 (1)	2.51 (2)	3.346 (2)	149 (2)

Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank Université Sidi Mohamed Ben Abdallah, Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2650).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Kourafalos, V. N., Marakos, P., Pouli, N., Terzis, A. & Townsend, L. B. (2002). *Heterocycles*, **57**, 2335–2343.
- Meanwell, N. A., Sit, S. Y., Gao, J. N., Wong, H. S., Gao, Q., St Laurent, D. R. & Balasubramanian, N. (1995). *J. Org. Chem.* **50**, 1565–1582.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *publCIF*. In preparation.

supporting information

Acta Cryst. (2010). E66, o756 [doi:10.1107/S1600536810007701]

6-Bromo-1,3-di-2-propynyl-1*H*-imidazo[4,5-*b*]pyridin-2(3*H*)-one


S. Dahmani, A. Haoudi, F. Capet, El Mokhtar Essassi and Seik Weng Ng

S1. Experimental

To a solution of 6-bromo-1,3-dihydroimidazo[4,5-*b*]pyridin-2-one (1 mmol), potassium carbonate (4 mmol) and tetra-*n*-butylammonium bromide (0.1 mmol) in DMF (20 ml) was added propargyl bromide (2.5 mmol). The solution was stirred for 48 hours. After completion of the reaction (as monitored by TLC), the salt was filtered and the solvent was removed under reduced pressure. The residue was purified by column chromatography on silica gel by using an ethyl acetate/hexane (1/1) mixture as eluent. Slow evaporation of the solvent furnished yellow crystals.

S2. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.97 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to $1.2U(C)$. The terminal acetylenic H-atoms were located in a difference Fourier map, and were refined with a distance restraint of C—H 0.95 ± 0.01 Å; their temperature factors were refined.

**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of $C_{12}H_8BrN_3O$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

6-Bromo-1,3-di-2-propynyl-1*H*-imidazo[4,5-*b*]pyridin- 2(3*H*)-one

Crystal data

$C_{12}H_8BrN_3O$

$M_r = 290.12$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 9.0725 (3)$ Å

$b = 18.6212 (5)$ Å

$c = 7.0684 (2)$ Å

$\beta = 102.995 (1)^\circ$

$V = 1163.56 (6)$ Å³

$Z = 4$

$F(000) = 576$

$D_x = 1.656 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9924 reflections

$\theta = 2.3\text{--}31.8^\circ$

$\mu = 3.52 \text{ mm}^{-1}$

$T = 293$ K

Prism, yellow

$0.35 \times 0.30 \times 0.15$ mm

Data collection

Bruker X8 APEXII
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.372$, $T_{\max} = 0.620$

27315 measured reflections
3383 independent reflections
2810 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$
 $\theta_{\max} = 30.0^\circ$, $\theta_{\min} = 2.6^\circ$
 $h = -12 \rightarrow 12$
 $k = -26 \rightarrow 26$
 $l = -9 \rightarrow 9$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.085$
 $S = 1.03$
3383 reflections
162 parameters
2 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0475P)^2 + 0.4099P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.87 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.80 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.40313 (3)	0.071904 (9)	0.35455 (3)	0.05174 (9)
O1	0.32318 (15)	0.47483 (7)	0.3528 (2)	0.0435 (3)
N1	0.12161 (15)	0.24467 (8)	0.3731 (2)	0.0354 (3)
N2	0.18505 (15)	0.37028 (8)	0.3651 (2)	0.0326 (3)
N3	0.42693 (14)	0.35966 (7)	0.35770 (18)	0.0289 (3)
C1	0.1822 (2)	0.17839 (9)	0.3720 (2)	0.0368 (4)
H1	0.1213	0.1387	0.3786	0.044*
C2	0.3302 (2)	0.16693 (8)	0.3614 (2)	0.0322 (3)
C3	0.43018 (17)	0.22350 (8)	0.3550 (2)	0.0289 (3)
H3	0.5302	0.2161	0.3481	0.035*
C4	0.21635 (17)	0.29782 (8)	0.3662 (2)	0.0279 (3)
C5	0.36927 (16)	0.29091 (8)	0.3596 (2)	0.0251 (3)
C6	0.0418 (2)	0.40256 (12)	0.3799 (3)	0.0457 (4)
H6A	0.0621	0.4471	0.4522	0.055*
H6B	-0.0093	0.3705	0.4524	0.055*
C7	-0.0586 (2)	0.41743 (12)	0.1911 (3)	0.0484 (5)
C8	-0.1389 (3)	0.43130 (16)	0.0410 (4)	0.0699 (8)
H8	-0.212 (3)	0.442 (2)	-0.073 (3)	0.108 (13)*
C9	0.31325 (18)	0.40998 (9)	0.3577 (2)	0.0307 (3)
C10	0.58352 (18)	0.38016 (10)	0.3664 (3)	0.0380 (4)
H10A	0.5957	0.4311	0.3946	0.046*
H10B	0.6068	0.3718	0.2408	0.046*
C11	0.68988 (18)	0.33965 (10)	0.5150 (3)	0.0397 (4)
C12	0.7772 (2)	0.30575 (14)	0.6294 (4)	0.0559 (6)

H12	0.849 (2)	0.2804 (13)	0.721 (3)	0.070 (8)*
-----	-----------	-------------	-----------	------------

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.07831 (18)	0.02651 (10)	0.04492 (13)	0.00475 (8)	0.00231 (10)	-0.00343 (6)
O1	0.0484 (7)	0.0281 (6)	0.0488 (7)	0.0041 (5)	-0.0001 (6)	0.0039 (5)
N1	0.0277 (6)	0.0436 (8)	0.0344 (7)	-0.0079 (6)	0.0059 (5)	-0.0006 (6)
N2	0.0261 (6)	0.0344 (7)	0.0357 (7)	0.0063 (5)	0.0039 (5)	0.0004 (5)
N3	0.0234 (6)	0.0261 (6)	0.0353 (6)	-0.0012 (4)	0.0023 (5)	0.0030 (5)
C1	0.0402 (9)	0.0356 (8)	0.0330 (8)	-0.0124 (7)	0.0050 (7)	-0.0002 (6)
C2	0.0437 (9)	0.0262 (7)	0.0246 (7)	-0.0018 (6)	0.0036 (6)	-0.0010 (5)
C3	0.0291 (7)	0.0294 (7)	0.0276 (7)	0.0028 (5)	0.0051 (6)	0.0003 (5)
C4	0.0248 (7)	0.0332 (7)	0.0240 (6)	0.0005 (5)	0.0022 (5)	-0.0001 (5)
C5	0.0237 (6)	0.0269 (7)	0.0236 (6)	-0.0015 (5)	0.0034 (5)	0.0005 (5)
C6	0.0346 (9)	0.0554 (11)	0.0481 (10)	0.0163 (8)	0.0114 (8)	-0.0029 (9)
C7	0.0306 (9)	0.0535 (11)	0.0610 (12)	0.0143 (8)	0.0101 (8)	0.0079 (9)
C8	0.0452 (13)	0.093 (2)	0.0672 (16)	0.0256 (12)	0.0040 (11)	0.0159 (13)
C9	0.0304 (7)	0.0294 (7)	0.0286 (7)	0.0028 (6)	-0.0014 (6)	0.0018 (6)
C10	0.0274 (7)	0.0392 (9)	0.0457 (9)	-0.0064 (6)	0.0047 (7)	0.0103 (7)
C11	0.0236 (7)	0.0455 (9)	0.0482 (10)	-0.0043 (6)	0.0042 (7)	0.0049 (8)
C12	0.0307 (9)	0.0659 (14)	0.0661 (14)	0.0016 (9)	0.0002 (9)	0.0171 (11)

Geometric parameters (\AA , ^\circ)

Br1—C2	1.8935 (16)	C3—C5	1.375 (2)
O1—C9	1.212 (2)	C3—H3	0.9300
N1—C4	1.319 (2)	C4—C5	1.405 (2)
N1—C1	1.352 (2)	C6—C7	1.463 (3)
N2—C4	1.378 (2)	C6—H6A	0.9700
N2—C9	1.389 (2)	C6—H6B	0.9700
N2—C6	1.457 (2)	C7—C8	1.173 (3)
N3—C5	1.3842 (19)	C8—H8	0.948 (10)
N3—C9	1.394 (2)	C10—C11	1.466 (2)
N3—C10	1.459 (2)	C10—H10A	0.9700
C1—C2	1.379 (3)	C10—H10B	0.9700
C1—H1	0.9300	C11—C12	1.179 (3)
C2—C3	1.397 (2)	C12—H12	0.939 (10)
C4—N1—C1	114.56 (14)	N3—C5—C4	107.09 (13)
C4—N2—C9	110.37 (13)	N2—C6—C7	113.27 (16)
C4—N2—C6	126.11 (16)	N2—C6—H6A	108.9
C9—N2—C6	123.44 (15)	C7—C6—H6A	108.9
C5—N3—C9	109.91 (13)	N2—C6—H6B	108.9
C5—N3—C10	127.46 (14)	C7—C6—H6B	108.9
C9—N3—C10	122.53 (14)	H6A—C6—H6B	107.7
N1—C1—C2	122.96 (15)	C8—C7—C6	178.2 (3)
N1—C1—H1	118.5	C7—C8—H8	174 (2)

C2—C1—H1	118.5	O1—C9—N2	126.84 (15)
C1—C2—C3	122.16 (15)	O1—C9—N3	127.61 (16)
C1—C2—Br1	119.74 (12)	N2—C9—N3	105.55 (13)
C3—C2—Br1	118.10 (13)	N3—C10—C11	111.93 (14)
C5—C3—C2	114.86 (14)	N3—C10—H10A	109.2
C5—C3—H3	122.6	C11—C10—H10A	109.2
C2—C3—H3	122.6	N3—C10—H10B	109.2
N1—C4—N2	126.85 (15)	C11—C10—H10B	109.2
N1—C4—C5	126.10 (15)	H10A—C10—H10B	107.9
N2—C4—C5	107.05 (13)	C12—C11—C10	177.6 (2)
C3—C5—N3	133.58 (14)	C11—C12—H12	177.7 (17)
C3—C5—C4	119.33 (14)		
C4—N1—C1—C2	-1.0 (2)	N1—C4—C5—C3	1.7 (2)
N1—C1—C2—C3	1.4 (3)	N2—C4—C5—C3	-178.61 (13)
N1—C1—C2—Br1	-178.53 (12)	N1—C4—C5—N3	-178.81 (14)
C1—C2—C3—C5	-0.2 (2)	N2—C4—C5—N3	0.92 (16)
Br1—C2—C3—C5	179.73 (10)	C4—N2—C6—C7	-93.3 (2)
C1—N1—C4—N2	179.82 (15)	C9—N2—C6—C7	90.2 (2)
C1—N1—C4—C5	-0.5 (2)	C4—N2—C9—O1	179.15 (16)
C9—N2—C4—N1	179.78 (15)	C6—N2—C9—O1	-3.9 (3)
C6—N2—C4—N1	2.9 (3)	C4—N2—C9—N3	-0.99 (17)
C9—N2—C4—C5	0.05 (17)	C6—N2—C9—N3	175.99 (15)
C6—N2—C4—C5	-176.84 (15)	C5—N3—C9—O1	-178.56 (16)
C2—C3—C5—N3	179.41 (15)	C10—N3—C9—O1	4.8 (3)
C2—C3—C5—C4	-1.2 (2)	C5—N3—C9—N2	1.58 (17)
C9—N3—C5—C3	177.86 (16)	C10—N3—C9—N2	-175.03 (14)
C10—N3—C5—C3	-5.7 (3)	C5—N3—C10—C11	-45.0 (2)
C9—N3—C5—C4	-1.58 (16)	C9—N3—C10—C11	131.00 (17)
C10—N3—C5—C4	174.83 (15)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C8—H8···O1 ⁱ	0.95 (1)	2.53 (2)	3.392 (3)	151 (3)
C12—H12···N1 ⁱⁱ	0.94 (1)	2.51 (2)	3.346 (2)	149 (2)

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $x+1, -y+1/2, z+1/2$.