

2,2'-[**(Propane-1,3-diyl)dinitrilo**]bis(**phenyl-methylidyne**)diphenol

Robert S. Black, David G. Billing,* Agata Bartyzel† and Ewa Cukrowska

Molecular Sciences Institute, School of Chemistry, University of the Witwatersrand, Private Bag 3, PO Wits 2050, South Africa
Correspondence e-mail: dave.billing@wits.ac.za


Received 9 March 2010; accepted 24 March 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.055; wR factor = 0.173; data-to-parameter ratio = 19.7.

In the title molecule, $\text{C}_{29}\text{H}_{26}\text{N}_2\text{O}_2$, there are two strong intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds involving the hydroxy and imine groups, forming $S(6)$ ring motifs. The dihedral angles between adjacent phenyl rings and phenol-containing planes are 85.27 (19) and 91.38 (18) $^\circ$. In the crystal structure, weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds connect molecules into a two-dimensional network.

Related literature

The title compound forms part of the group of Schiff bases with a similar method of synthesis as described in Schilf *et al.* (2007). The intermolecular hydrogen bonds $\text{O}-\text{H}\cdots\text{N}$ between the hydroxy and imine are common to this type of compound as shown with the series of compounds reported by Fernández *et al.* (2001); Kabak (2003); Wojciechowski *et al.* (2001); Dey *et al.* (2001); Koşar, *et al.* (2004); Lu, *et al.* (2008); Qiu & Zhao (2008); Montazerzohori *et al.* (2009); Corden *et al.* (1996). For a description of hydrogen-bond motifs, see: Bernstein *et al.* (1995).



‡ On leave from: Faculty of Chemistry Department of General and Coordination Chemistry Maria Curie-Sklodowska University Maria Curie-Sklodowska Sq. 2 20-031 Lublin Poland.

Experimental

Crystal data

$\text{C}_{29}\text{H}_{26}\text{N}_2\text{O}_2$	$V = 2443.7$ (5) \AA^3
$M_r = 434.52$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 18.226$ (2) \AA	$\mu = 0.07\text{ mm}^{-1}$
$b = 8.2303$ (9) \AA	$T = 293\text{ K}$
$c = 18.642$ (2) \AA	$0.7 \times 0.5 \times 0.28\text{ mm}$
$\beta = 119.086$ (5) $^\circ$	

Data collection

Bruker SMART 1K CCD area-detector diffractometer	5896 independent reflections
21350 measured reflections	3749 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.071$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$	300 parameters
$wR(F^2) = 0.173$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.23\text{ e \AA}^{-3}$
5896 reflections	$\Delta\rho_{\text{min}} = -0.36\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1A…N1	0.82	1.80	2.5344 (17)	148
O2—H2A…N2	0.82	1.81	2.5431 (18)	148
C17—H17…O2 ⁱ	0.93	2.56	3.481 (2)	168
C18—H18…O1 ⁱⁱ	0.93	2.47	3.395 (3)	174
C21—H21…O2 ⁱⁱⁱ	0.93	2.56	3.492 (2)	175

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART-NT* (Bruker, 1998); cell refinement: *SAINT-Plus* (Bruker, 1999); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The University of the Witwatersrand and the National Research Foundation (GUN 2069064) are thanked for providing the infrastructure and for the award of a research grant required to carry out this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5012).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (1998). *SMART-NT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (1999). *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Corden, J. P., Errington, W., Moore, P., Phillips, P. R. & Wallbridge, M. G. H. (1996). *Acta Cryst.* **C52**, 3199–3202.
- Dey, D. K., Dey, S. P., Elmali, A. & Elerman, Y. (2001). *J. Mol. Struct.* **562**, 177–184.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Fernández, G. J. M., del Rio-Portilla, F., Quiroz-Garcia, B., Toscano, R. A. & Salcedo, R. (2001). *J. Mol. Struct.* **561**, 197–207.
- Kabak, M. (2003). *J. Mol. Struct.* **655**, 135–139.
- Koşar, B., Büyükgür, O., Albayrak, Ç. & Odabaşoğlu, M. (2004). *Acta Cryst.* **C60**, o458–o460.
- Lu, J.-F., Min, S.-T., Ji, X.-H. & Dang, Z.-H. (2008). *Acta Cryst. E64*, o1693.

- Montazerozohori, M., Habibi, M. H., Hojjati, A., Mokhtari, R., Yamane, Y. & Suzuki, T. (2009). *Acta Cryst. E*65, o1662–o1663.
- Qiu, F. & Zhao, L.-M. (2008). *Acta Cryst. E*64, o2067.
- Schilf, W., Kamienski, B., Szady-Chełmieniecka, A., Grech, E., Makal, A. & Wozniak, K. (2007). *J. Mol. Struct.* **94–101**, 844–845
- Sheldrick, G. M. (2008). *Acta Cryst. A*64, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D*65, 148–155.
- Wojciechowski, G., Ratajczak-Sitarz, M., Katrusiak, A., Schilf, W., Przybylski, P. & Brzezinski, B. (2001). *J. Mol. Struct.* **650**, 191–199.

supporting information

Acta Cryst. (2010). E66, o1002–o1003 [doi:10.1107/S1600536810011189]

2,2'-(Propane-1,3-diylidinitrilo)bis(phenylmethyldyne)diphenol

Robert S. Black, David G. Billing, Agata Bartyzel and Ewa Cukrowska

S1. Comment


The molecular structure of the title compound is shown in Fig. 1. There are two strong intramolecular O—H..O hydrogen bonds involving the hydroxyl and imine groups forming S(6) ring motifs (Bernstein *et al.*, 1995). These types of hydrogen bonds are common to some reported molecular structures (Schilf *et al.*, 2007; Fernández *et al.*, 2001; Kabak, 2003; Wojciechowski *et al.*, 2001; Dey *et al.*, 2001; Koşar, *et al.*, 2004; Lu, *et al.*, 2008; Qiu & Zhao, 2008; Montazerozohori *et al.*, 2009; Corden *et al.*, 1996). In the crystal structure, weak intermolecular C—H···O hydrogen bond connect molecules to form a two-dimensional network.

S2. Experimental

A mixture of 1.0 mmol (2.00 g) of 2-hydroxybenzophenone, 0.5 mmole (0.42 cm³) of 1,3-propanediamine and 2 drops of glacial acetic acid in 40 ml of methanol was refluxed for 8 h. The excess of solvent (ca. 30 ml) was then evaporated. After cooling to 277 K a yellow solid was produced. The polycrystalline product was collected by filtration, washed with methanol and dried. A yield of 52% was obtained. Recrystallization from an ethanol solution yielded single crystals suitable for x-ray diffraction. Elemental analysis C% 79.67 H% 6.26 N% 6.11%.

S3. Refinement

All H atoms were refined using a riding model, with C—H = 0.93–0.97 Å, O—H = 0.82 Å, and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{O})$.

**Figure 1**

The molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level. Hydrogen bonds are shown as dashed lines.

2,2'-[Propane-1,3-diylidinitrilo]bis(phenylmethyldyne)diphenol

Crystal data

$C_{29}H_{26}N_2O_2$
 $M_r = 434.52$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 18.226 (2)$ Å
 $b = 8.2303 (9)$ Å
 $c = 18.642 (2)$ Å
 $\beta = 119.086 (5)^\circ$
 $V = 2443.7 (5)$ Å³
 $Z = 4$

$F(000) = 920$
 $D_x = 1.181$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 21350 reflections
 $\theta = 2.1\text{--}23.0^\circ$
 $\mu = 0.07$ mm⁻¹
 $T = 293$ K
Block, yellow
 $0.7 \times 0.5 \times 0.28$ mm

Data collection

Bruker SMART 1K CCD area-detector
diffractometer
 φ and ω scans
21350 measured reflections
5896 independent reflections
3749 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.071$
 $\theta_{\text{max}} = 28^\circ, \theta_{\text{min}} = 1.3^\circ$
 $h = -19 \rightarrow 24$
 $k = -10 \rightarrow 10$
 $l = -24 \rightarrow 20$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.173$
 $S = 1.03$
5896 reflections
300 parameters
0 restraints
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0897P)^2 + 0.2258P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$

$$\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.17087 (13)	0.1655 (3)	0.08228 (11)	0.0718 (5)
H1	0.1926	0.2672	0.0813	0.086*
C2	0.15625 (16)	0.0517 (3)	0.02117 (14)	0.0940 (7)
H2	0.1689	0.0776	-0.0201	0.113*
C3	0.12367 (14)	-0.0961 (3)	0.02162 (16)	0.0947 (8)
H3	0.1142	-0.1708	-0.0194	0.114*
C4	0.10480 (16)	-0.1362 (3)	0.08146 (16)	0.0960 (7)
H4	0.0823	-0.2377	0.0812	0.115*
C5	0.11925 (13)	-0.0247 (3)	0.14328 (13)	0.0759 (5)
H5	0.1062	-0.0519	0.1842	0.091*
C6	0.15300 (9)	0.12641 (19)	0.14400 (9)	0.0501 (4)
C7	0.17057 (8)	0.24201 (18)	0.21320 (9)	0.0457 (3)
C8	0.10782 (9)	0.36732 (18)	0.20194 (9)	0.0476 (3)
C9	0.03699 (10)	0.3944 (2)	0.12497 (11)	0.0612 (4)
H9	0.0291	0.3308	0.0805	0.073*
C10	-0.02150 (12)	0.5138 (3)	0.11365 (13)	0.0775 (6)
H10	-0.0675	0.5312	0.062	0.093*
C11	-0.01072 (13)	0.6066 (3)	0.17994 (16)	0.0826 (6)
H11	-0.05	0.6861	0.1726	0.099*
C12	0.05729 (13)	0.5829 (2)	0.25660 (14)	0.0735 (5)
H12	0.0632	0.6461	0.3004	0.088*
C13	0.11810 (10)	0.4642 (2)	0.26941 (10)	0.0548 (4)
C14	0.30624 (10)	0.1168 (2)	0.29798 (11)	0.0637 (4)
H14A	0.3103	0.1037	0.2483	0.076*
H14B	0.2935	0.0117	0.3128	0.076*
C15	0.38907 (10)	0.1788 (2)	0.36662 (10)	0.0590 (4)
H15A	0.385	0.1869	0.4165	0.071*
H15B	0.3992	0.287	0.3528	0.071*
C16	0.46364 (10)	0.0702 (2)	0.38294 (10)	0.0606 (4)
H16A	0.5125	0.1034	0.4336	0.073*
H16B	0.4508	-0.0416	0.3891	0.073*
C17	0.66693 (10)	0.2305 (2)	0.45511 (10)	0.0602 (4)
H17	0.6442	0.3306	0.4316	0.072*
C18	0.73422 (12)	0.2233 (3)	0.53470 (12)	0.0729 (5)
H18	0.756	0.3185	0.5644	0.087*
C19	0.76854 (12)	0.0763 (3)	0.56945 (12)	0.0743 (5)

H19	0.813	0.0721	0.6229	0.089*
C20	0.73734 (12)	-0.0646 (3)	0.52550 (13)	0.0747 (5)
H20	0.7618	-0.1637	0.5489	0.09*
C21	0.66949 (11)	-0.0604 (2)	0.44627 (11)	0.0608 (4)
H21	0.6481	-0.1564	0.4172	0.073*
C22	0.63373 (9)	0.08832 (18)	0.41069 (9)	0.0469 (3)
C23	0.55780 (9)	0.09263 (18)	0.32620 (9)	0.0496 (4)
C24	0.57020 (11)	0.10567 (19)	0.25341 (10)	0.0539 (4)
C25	0.65069 (12)	0.1279 (2)	0.26229 (12)	0.0676 (5)
H25	0.6968	0.134	0.3147	0.081*
C26	0.66269 (15)	0.1409 (3)	0.19441 (15)	0.0808 (6)
H26	0.7163	0.156	0.2012	0.097*
C27	0.59361 (18)	0.1312 (3)	0.11627 (15)	0.0861 (7)
H27	0.6013	0.1387	0.0706	0.103*
C28	0.51399 (16)	0.1106 (2)	0.10547 (12)	0.0769 (6)
H28	0.4686	0.1045	0.0527	0.092*
C29	0.50056 (12)	0.09859 (19)	0.17301 (11)	0.0597 (4)
N1	0.23864 (8)	0.23289 (17)	0.28316 (8)	0.0562 (3)
N2	0.48206 (8)	0.08218 (17)	0.31467 (8)	0.0564 (3)
O1	0.18408 (8)	0.44387 (17)	0.34460 (7)	0.0709 (4)
H1A	0.2167	0.3781	0.3424	0.106*
O2	0.42130 (8)	0.08050 (17)	0.15964 (7)	0.0715 (4)
H2A	0.4218	0.074	0.2038	0.107*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0820 (13)	0.0800 (12)	0.0680 (11)	-0.0146 (10)	0.0478 (10)	-0.0153 (9)
C2	0.1009 (17)	0.125 (2)	0.0795 (14)	-0.0200 (15)	0.0625 (14)	-0.0331 (13)
C3	0.0776 (14)	0.1138 (19)	0.1001 (16)	-0.0262 (13)	0.0490 (13)	-0.0609 (14)
C4	0.0966 (17)	0.0856 (15)	0.1172 (18)	-0.0384 (13)	0.0610 (16)	-0.0465 (14)
C5	0.0807 (13)	0.0769 (12)	0.0816 (13)	-0.0279 (10)	0.0484 (11)	-0.0252 (10)
C6	0.0390 (7)	0.0596 (9)	0.0508 (8)	-0.0030 (6)	0.0212 (6)	-0.0107 (7)
C7	0.0393 (7)	0.0540 (8)	0.0465 (7)	-0.0056 (6)	0.0229 (6)	-0.0073 (6)
C8	0.0400 (7)	0.0531 (8)	0.0529 (8)	-0.0062 (6)	0.0250 (7)	-0.0070 (6)
C9	0.0456 (9)	0.0714 (11)	0.0615 (10)	-0.0002 (7)	0.0220 (8)	-0.0086 (8)
C10	0.0513 (10)	0.0855 (13)	0.0823 (13)	0.0124 (9)	0.0221 (9)	-0.0007 (11)
C11	0.0607 (12)	0.0732 (12)	0.1165 (18)	0.0144 (9)	0.0451 (13)	-0.0065 (12)
C12	0.0679 (12)	0.0709 (12)	0.0926 (14)	0.0000 (9)	0.0475 (11)	-0.0229 (10)
C13	0.0520 (9)	0.0573 (9)	0.0625 (10)	-0.0083 (7)	0.0338 (8)	-0.0127 (7)
C14	0.0450 (9)	0.0724 (11)	0.0627 (10)	0.0065 (7)	0.0175 (8)	-0.0136 (8)
C15	0.0442 (8)	0.0750 (11)	0.0546 (9)	-0.0015 (7)	0.0217 (7)	-0.0083 (8)
C16	0.0435 (8)	0.0848 (12)	0.0528 (9)	0.0052 (8)	0.0229 (7)	0.0092 (8)
C17	0.0553 (9)	0.0568 (9)	0.0662 (10)	-0.0005 (7)	0.0277 (8)	-0.0048 (8)
C18	0.0569 (10)	0.0913 (14)	0.0704 (12)	-0.0188 (10)	0.0310 (9)	-0.0281 (10)
C19	0.0501 (10)	0.1127 (17)	0.0548 (10)	0.0029 (10)	0.0213 (8)	0.0014 (10)
C20	0.0621 (11)	0.0852 (14)	0.0726 (12)	0.0204 (10)	0.0294 (10)	0.0230 (11)
C21	0.0566 (10)	0.0567 (9)	0.0669 (10)	0.0052 (7)	0.0282 (9)	0.0009 (8)

C22	0.0402 (7)	0.0523 (8)	0.0505 (8)	0.0024 (6)	0.0237 (7)	-0.0007 (6)
C23	0.0455 (8)	0.0508 (8)	0.0518 (8)	0.0035 (6)	0.0230 (7)	-0.0005 (6)
C24	0.0605 (9)	0.0511 (8)	0.0554 (9)	0.0042 (7)	0.0324 (8)	-0.0011 (7)
C25	0.0690 (11)	0.0714 (11)	0.0764 (12)	0.0021 (9)	0.0465 (10)	-0.0025 (9)
C26	0.0994 (16)	0.0749 (13)	0.1028 (16)	0.0019 (11)	0.0763 (15)	-0.0001 (11)
C27	0.136 (2)	0.0732 (13)	0.0831 (15)	0.0020 (13)	0.0797 (16)	0.0013 (11)
C28	0.1119 (17)	0.0633 (11)	0.0601 (11)	0.0001 (11)	0.0456 (12)	-0.0017 (8)
C29	0.0754 (12)	0.0501 (9)	0.0562 (9)	0.0028 (8)	0.0340 (9)	-0.0002 (7)
N1	0.0429 (7)	0.0685 (9)	0.0517 (7)	0.0042 (6)	0.0187 (6)	-0.0111 (6)
N2	0.0442 (7)	0.0732 (9)	0.0494 (7)	0.0036 (6)	0.0209 (6)	0.0030 (6)
O1	0.0677 (8)	0.0854 (9)	0.0572 (7)	0.0013 (6)	0.0285 (6)	-0.0219 (6)
O2	0.0667 (8)	0.0858 (9)	0.0505 (6)	-0.0029 (6)	0.0193 (6)	-0.0008 (6)

Geometric parameters (\AA , $^\circ$)

C1—C6	1.378 (2)	C15—H15B	0.97
C1—C2	1.396 (3)	C16—N2	1.469 (2)
C1—H1	0.93	C16—H16A	0.97
C2—C3	1.356 (3)	C16—H16B	0.97
C2—H2	0.93	C17—C22	1.391 (2)
C3—C4	1.358 (3)	C17—C18	1.393 (3)
C3—H3	0.93	C17—H17	0.93
C4—C5	1.394 (3)	C18—C19	1.371 (3)
C4—H4	0.93	C18—H18	0.93
C5—C6	1.385 (2)	C19—C20	1.374 (3)
C5—H5	0.93	C19—H19	0.93
C6—C7	1.507 (2)	C20—C21	1.392 (3)
C7—N1	1.2934 (19)	C20—H20	0.93
C7—C8	1.477 (2)	C21—C22	1.394 (2)
C8—C9	1.405 (2)	C21—H21	0.93
C8—C13	1.422 (2)	C22—C23	1.509 (2)
C9—C10	1.389 (3)	C23—N2	1.293 (2)
C9—H9	0.93	C23—C24	1.484 (2)
C10—C11	1.383 (3)	C24—C25	1.406 (2)
C10—H10	0.93	C24—C29	1.418 (2)
C11—C12	1.376 (3)	C25—C26	1.389 (3)
C11—H11	0.93	C25—H25	0.93
C12—C13	1.408 (2)	C26—C27	1.389 (3)
C12—H12	0.93	C26—H26	0.93
C13—O1	1.342 (2)	C27—C28	1.375 (3)
C14—N1	1.475 (2)	C27—H27	0.93
C14—C15	1.516 (2)	C28—C29	1.398 (3)
C14—H14A	0.97	C28—H28	0.93
C14—H14B	0.97	C29—O2	1.349 (2)
C15—C16	1.528 (2)	O1—H1A	0.82
C15—H15A	0.97	O2—H2A	0.82
C6—C1—C2	119.74 (19)	H15A—C15—H15B	107.7

C6—C1—H1	120.1	N2—C16—C15	110.13 (13)
C2—C1—H1	120.1	N2—C16—H16A	109.6
C3—C2—C1	120.5 (2)	C15—C16—H16A	109.6
C3—C2—H2	119.8	N2—C16—H16B	109.6
C1—C2—H2	119.8	C15—C16—H16B	109.6
C2—C3—C4	120.63 (19)	H16A—C16—H16B	108.1
C2—C3—H3	119.7	C22—C17—C18	120.06 (16)
C4—C3—H3	119.7	C22—C17—H17	120
C3—C4—C5	119.9 (2)	C18—C17—H17	120
C3—C4—H4	120.1	C19—C18—C17	120.31 (18)
C5—C4—H4	120.1	C19—C18—H18	119.8
C6—C5—C4	120.23 (19)	C17—C18—H18	119.8
C6—C5—H5	119.9	C18—C19—C20	120.11 (17)
C4—C5—H5	119.9	C18—C19—H19	119.9
C1—C6—C5	119.07 (16)	C20—C19—H19	119.9
C1—C6—C7	121.98 (15)	C19—C20—C21	120.55 (17)
C5—C6—C7	118.94 (15)	C19—C20—H20	119.7
N1—C7—C8	118.72 (13)	C21—C20—H20	119.7
N1—C7—C6	121.82 (13)	C20—C21—C22	119.74 (16)
C8—C7—C6	119.45 (12)	C20—C21—H21	120.1
C9—C8—C13	118.36 (14)	C22—C21—H21	120.1
C9—C8—C7	121.28 (13)	C17—C22—C21	119.22 (15)
C13—C8—C7	120.37 (13)	C17—C22—C23	120.95 (13)
C10—C9—C8	121.58 (17)	C21—C22—C23	119.78 (13)
C10—C9—H9	119.2	N2—C23—C24	118.54 (14)
C8—C9—H9	119.2	N2—C23—C22	122.36 (14)
C11—C10—C9	119.33 (19)	C24—C23—C22	119.09 (13)
C11—C10—H10	120.3	C25—C24—C29	118.44 (16)
C9—C10—H10	120.3	C25—C24—C23	121.04 (15)
C12—C11—C10	120.99 (17)	C29—C24—C23	120.51 (15)
C12—C11—H11	119.5	C26—C25—C24	121.34 (19)
C10—C11—H11	119.5	C26—C25—H25	119.3
C11—C12—C13	120.75 (17)	C24—C25—H25	119.3
C11—C12—H12	119.6	C27—C26—C25	119.15 (19)
C13—C12—H12	119.6	C27—C26—H26	120.4
O1—C13—C12	119.70 (15)	C25—C26—H26	120.4
O1—C13—C8	121.33 (14)	C28—C27—C26	120.96 (18)
C12—C13—C8	118.98 (16)	C28—C27—H27	119.5
N1—C14—C15	109.80 (14)	C26—C27—H27	119.5
N1—C14—H14A	109.7	C27—C28—C29	120.7 (2)
C15—C14—H14A	109.7	C27—C28—H28	119.6
N1—C14—H14B	109.7	C29—C28—H28	119.6
C15—C14—H14B	109.7	O2—C29—C28	118.80 (17)
H14A—C14—H14B	108.2	O2—C29—C24	121.83 (15)
C14—C15—C16	113.32 (14)	C28—C29—C24	119.37 (18)
C14—C15—H15A	108.9	C7—N1—C14	122.06 (13)
C16—C15—H15A	108.9	C23—N2—C16	122.40 (13)
C14—C15—H15B	108.9	C13—O1—H1A	109.5

C16—C15—H15B	108.9	C29—O2—H2A	109.5
C6—C1—C2—C3	0.8 (4)	C18—C19—C20—C21	-1.6 (3)
C1—C2—C3—C4	-0.1 (4)	C19—C20—C21—C22	1.0 (3)
C2—C3—C4—C5	-0.2 (4)	C18—C17—C22—C21	-1.1 (2)
C3—C4—C5—C6	-0.2 (4)	C18—C17—C22—C23	176.55 (14)
C2—C1—C6—C5	-1.2 (3)	C20—C21—C22—C17	0.3 (2)
C2—C1—C6—C7	177.65 (19)	C20—C21—C22—C23	-177.36 (15)
C4—C5—C6—C1	0.9 (3)	C17—C22—C23—N2	-90.01 (19)
C4—C5—C6—C7	-177.97 (19)	C21—C22—C23—N2	87.61 (19)
C1—C6—C7—N1	-96.1 (2)	C17—C22—C23—C24	91.38 (18)
C5—C6—C7—N1	82.8 (2)	C21—C22—C23—C24	-91.00 (18)
C1—C6—C7—C8	85.27 (19)	N2—C23—C24—C25	175.86 (15)
C5—C6—C7—C8	-95.88 (19)	C22—C23—C24—C25	-5.5 (2)
N1—C7—C8—C9	173.25 (15)	N2—C23—C24—C29	-3.1 (2)
C6—C7—C8—C9	-8.1 (2)	C22—C23—C24—C29	175.57 (13)
N1—C7—C8—C13	-6.6 (2)	C29—C24—C25—C26	-0.7 (3)
C6—C7—C8—C13	172.08 (13)	C23—C24—C25—C26	-179.70 (16)
C13—C8—C9—C10	1.0 (3)	C24—C25—C26—C27	-0.2 (3)
C7—C8—C9—C10	-178.90 (17)	C25—C26—C27—C28	0.7 (3)
C8—C9—C10—C11	-1.2 (3)	C26—C27—C28—C29	-0.1 (3)
C9—C10—C11—C12	0.5 (3)	C27—C28—C29—O2	179.16 (17)
C10—C11—C12—C13	0.4 (3)	C27—C28—C29—C24	-0.9 (3)
C11—C12—C13—O1	179.67 (18)	C25—C24—C29—O2	-178.77 (16)
C11—C12—C13—C8	-0.6 (3)	C23—C24—C29—O2	0.2 (2)
C9—C8—C13—O1	179.63 (15)	C25—C24—C29—C28	1.2 (2)
C7—C8—C13—O1	-0.5 (2)	C23—C24—C29—C28	-179.77 (15)
C9—C8—C13—C12	-0.1 (2)	C8—C7—N1—C14	-176.84 (14)
C7—C8—C13—C12	179.80 (15)	C6—C7—N1—C14	4.5 (2)
N1—C14—C15—C16	-176.82 (14)	C15—C14—N1—C7	155.47 (15)
C14—C15—C16—N2	70.2 (2)	C24—C23—N2—C16	-179.47 (14)
C22—C17—C18—C19	0.5 (3)	C22—C23—N2—C16	1.9 (2)
C17—C18—C19—C20	0.8 (3)	C15—C16—N2—C23	137.49 (16)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1A···N1	0.82	1.80	2.5344 (17)	148
O2—H2A···N2	0.82	1.81	2.5431 (18)	148
C17—H17···O2 ⁱ	0.93	2.56	3.481 (2)	168
C18—H18···O1 ⁱⁱ	0.93	2.47	3.395 (3)	174
C21—H21···O2 ⁱⁱⁱ	0.93	2.56	3.492 (2)	175

Symmetry codes: (i) -x+1, y+1/2, -z+1/2; (ii) -x+1, -y+1, -z+1; (iii) -x+1, y-1/2, -z+1/2.