

## (*E*)-1-[(2-Fluorophenyl)iminomethyl]-2-naphthol-(*Z*)-1-[(2-fluorophenyl)amino-methylidene]naphthalen-2(1*H*)-one (0.57/0.43)

Gökhan Alpaslan,<sup>a</sup> Mustafa Macit,<sup>b</sup> Orhan Büyükgüngör<sup>a\*</sup> and Ahmet Erdönmez<sup>a</sup>

<sup>a</sup>Department of Physics, Faculty of Arts & Science, Ondokuz Mayıs University, TR-55139 Kurupelit-Samsun, Turkey, and <sup>b</sup>Department of Chemistry, Faculty of Arts & Science, Ondokuz Mayıs University, TR-55139 Kurupelit-Samsun, Turkey  
Correspondence e-mail: orhanb@omu.edu.tr


Received 14 April 2010; accepted 22 April 2010

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.006$  Å;  $R$  factor = 0.047;  $wR$  factor = 0.113; data-to-parameter ratio = 8.1.

The title Schiff base compound,  $0.57\text{C}_{17}\text{H}_{12}\text{FNO}\cdot0.43\text{C}_{17}\text{H}_{12}\text{FNO}$ , reveals both the enol (OH) and keto (NH) tautomeric forms with occupancies of 0.57 (6) and 0.43 (6), respectively. The tautomeric forms are stabilized by intramolecular O—H···N (enol) and N—H···O (keto) hydrogen bonds. The dihedral angle between the naphthalene ring system and the benzene ring is 32.76 (1)°.

### Related literature

For the biological properties of Schiff bases, see: Lozier *et al.* (1975). For the coordination chemistry of Schiff bases, see: Kargar *et al.* (2009); Yeap *et al.* (2009). For Schiff base tautomerism, see: Hökelek *et al.* (2000); Kaitner & Pavlovic (1996); Karabiyik *et al.* (2007); Nazir *et al.* (2000); Odabaşoğlu *et al.* (2005); Yıldız *et al.* (1998); Tanak *et al.* (2009). For bond-length data, see: Allen *et al.* (1987).



### Experimental

#### Crystal data

$0.57\text{C}_{17}\text{H}_{12}\text{FNO}\cdot0.43\text{C}_{17}\text{H}_{12}\text{FNO}$   
 $M_r = 265.28$

Orthorhombic,  $P2_12_12_1$   
 $a = 7.2841 (3)$  Å

$b = 12.2158 (6)$  Å  
 $c = 14.5731 (7)$  Å  
 $V = 1296.73 (10)$  Å<sup>3</sup>  
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 0.10$  mm<sup>-1</sup>  
 $T = 296$  K  
 $0.73 \times 0.31 \times 0.10$  mm

#### Data collection

Stoe IPDSII diffractometer  
Absorption correction: integration (*X-RED32*; Stoe & Cie, 2002)  
 $T_{\min} = 0.967$ ,  $T_{\max} = 0.993$

5655 measured reflections  
1477 independent reflections  
951 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.040$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.113$   
 $S = 0.97$   
1477 reflections  
183 parameters

1 restraint  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.23$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.14$  e Å<sup>-3</sup>

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1A···N1	0.82	1.82	2.535 (4)	144
N1—H1B···O1	0.86	1.86	2.535 (4)	134

Data collection: *X-AREA* (Stoe & Cie, 2002); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the Faculty of Arts and Sciences, Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDSII diffractometer (purchased under grant No. F279 of the University Research Fund).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5080).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hökelek, T., Kılıç, Z., İşıklan, M. & Toy, M. (2000). *J. Mol. Struct.* **523**, 61–69.
- Kaitner, B. & Pavlovic, G. (1996). *Acta Cryst. C52*, 2573–2575.
- Karabiyik, H., Güzel, B., Aygün, M., Boğa, G. & Büyükgüngör, O. (2007). *Acta Cryst. C63*, o215–o218.
- Kargar, H., Jamshidvand, A., Fun, H.-K. & Kia, R. (2009). *Acta Cryst. E65*, m403–m404.
- Lozier, R. H., Bogomolni, R. A. & Stoeckenius, W. (1975). *Biophys. J.* **15**, 955–962.
- Nazir, H., Yıldız, M., Yılmaz, H., Tahir, M. N. & Ülkü, D. (2000). *J. Mol. Struct.* **524**, 241–250.
- Odabaşoğlu, M., Albayrak, Ç. & Büyükgüngör, O. (2005). *Acta Cryst. E61*, o425–o426.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Stoe & Cie (2002). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.
- Tanak, H., Bingöl Alpaslan, Y., Yavuz, M., Ağar, E., Erşahin, F. & Büyükgüngör, O. (2009). *Acta Cryst. E65*, o1572.
- Yeap, C. S., Kia, R., Kargar, H. & Fun, H.-K. (2009). *Acta Cryst. E65*, m570–m571.
- Yıldız, M., Kılıç, Z. & Hökelek, T. (1998). *J. Mol. Struct.* **441**, 1–10.

# supporting information

*Acta Cryst.* (2010). E66, o1178 [https://doi.org/10.1107/S1600536810014777]

## (E)-1-[(2-Fluorophenyl)iminomethyl]-2-naphthol-(Z)-1-[(2-fluorophenyl)amino-methylidene]naphthalen-2(1H)-one (0.57/0.43)

Gökhan Alpaslan, Mustafa Macit, Orhan Büyükgüngör and Ahmet Erdönmez

### S1. Comment

Schiff bases often exhibit various biological activities and in many cases were shown to have antibacterial, anticancer, anti-inflammatory and antitoxic properties (Lozier *et al.*, 1975). Schiff bases have also been used as versatile ligands in coordination chemistry (Kargar *et al.*, 2009; Yeap *et al.*, 2009). There are two types of intramolecular hydrogen bonds in Schiff bases, namely N—H···O in keto (NH) (Hökelek *et al.*, 2000) and N···H—O in enol (OH) (Odabaşoğlu *et al.*, 2005) tautomeric forms. In the solid state, while OH tautomeric forms of Schiff bases are predominant in salicylaldimines (Kaitner & Pavlovic, 1996; Yıldız *et al.*, 1998), both NH and OH forms have been found in naphthaldimine Schiff base compounds (Nazır *et al.*, 2000; Karabiyık *et al.*, 2007). Our investigations shows that in the title compound both OH (enol) and NH (keto) tautomers coexist with occupancies of 0.57 (6) and 0.43 (6), respectively. This evidence is also supported by the observed IR vibrational bands given in the experimental section.

An ORTEP-3 (Farrugia, 1997) plot of the molecule of (I) is shown in Fig.1. The C2—O1 [1.314 (5) Å] and C11—N1 [1.306 (4) Å] bond lengths are intermediate between the single and double C—O (1.362 and 1.222 Å, respectively) and C—N bond lengths (1.339 and 1.279 Å, respectively) (Allen *et al.*, 1987). Similar results were observed for 2-[(2,4-di-methylphenyl)iminomethyl]-3,5-dimethoxyphenol (Tanak *et al.*, 2009). The molecule of the title compound is not planar, with a dihedral angle of 32.76 (1)° between naphthalene and benzene rings. The molecular structure is stabilized by O—H···N or N—H···O hydrogen bonds.


### S2. Experimental

The title compound was prepared by refluxing a mixture of a solution containing 2-hydroxy-1-naphthaldehyde (172 mg, 1 mmol) in 50 ml ethanol and a solution containing 2-fluoroaniline (111 mg, 1 mmol) in 30 ml ethanol. The reaction mixture was stirred for 3 h under reflux. The crystals of the title compound were obtained by slow evaporation of ethanol (yield 68%; m.p. 361–363 K). The FT-IR spectra of the title compound was recorded on a KBr pellets with a Schmadzu FT-IR 8900 spectrophotometer. IR (KBr)  $\nu$  = 3558 (O—H), 3420 (N—H), 1698 (C=O) weak, 1620 (C=N) cm<sup>-1</sup>.

### S3. Refinement

The absolute configuration could not be determined from X-ray data, as no strong anomalous scatterer is present; 1058 Friedel pairs were merged before the final refinement. All H atoms (except H1A and H1B) were placed in calculated positions and constrained to ride on their parent atoms, with C—H = 0.93 Å and  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ . Inspection of the tautomeric hydrogen atom's (H1) location in difference Fourier map between O1 and N1 indicated that there is a positional disorder for this H atom. Hence, atoms H1A and H1B bonded with O1 and N1 were positioned geometrically and their occupancies were refined to 0.57 (6) and 0.43 (6), respectively. A restraint (DELU instruction in SHELXL97, Sheldrick, 2008) was used in order to maintain a reasonable geometry and atomic displacement parameters for atoms C13

and F1.




**Figure 1**

The molecular structure of (I), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 40% probability level. Both OH (solid O–H bond) and NH (dashed N–H bond) tautomers are shown.

**(E)-1-[(2-Fluorophenyl)iminomethyl]-2-naphthol-(Z)-1-[(2-fluorophenyl)aminomethylidene]naphthalen-2(1H)-one (0.57/0.43)**

*Crystal data*



$M_r = 265.28$

Orthorhombic,  $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 7.2841 (3) \text{ \AA}$

$b = 12.2158 (6) \text{ \AA}$

$c = 14.5731 (7) \text{ \AA}$

$V = 1296.73 (10) \text{ \AA}^3$

$Z = 4$

$F(000) = 552$

$D_x = 1.359 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4776 reflections

$\theta = 1.4\text{--}27.8^\circ$

$\mu = 0.10 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Needle, yellow

$0.73 \times 0.31 \times 0.10 \text{ mm}$

*Data collection*

Stoe IPDSII

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 6.67 pixels  $\text{mm}^{-1}$

$\omega$  scans

Absorption correction: integration

(*X-RED32*; Stoe & Cie, 2002)

$T_{\min} = 0.967$ ,  $T_{\max} = 0.993$

5655 measured reflections

1477 independent reflections

951 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.040$

$\theta_{\max} = 26.0^\circ$ ,  $\theta_{\min} = 2.2^\circ$

$h = -8 \rightarrow 8$

$k = -13 \rightarrow 15$

$l = -17 \rightarrow 16$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.047$$

$$wR(F^2) = 0.113$$

$$S = 0.97$$

1477 reflections

183 parameters

1 restraint

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0576P)^2]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.23 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.14 \text{ e \AA}^{-3}$$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor wR and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^* / U_{\text{eq}}$	Occ. (<1)
C1	0.3705 (5)	0.8439 (3)	0.5517 (2)	0.0525 (9)	
C2	0.4237 (5)	0.7941 (4)	0.6359 (3)	0.0643 (10)	
C3	0.4381 (5)	0.8594 (4)	0.7158 (3)	0.0736 (12)	
H3	0.4738	0.8268	0.7707	0.088*	
C4	0.4018 (5)	0.9661 (4)	0.7144 (3)	0.0700 (11)	
H4	0.4127	1.0060	0.7685	0.084*	
C5	0.3467 (5)	1.0212 (3)	0.6328 (3)	0.0614 (10)	
C6	0.3091 (6)	1.1331 (4)	0.6324 (3)	0.0780 (12)	
H6	0.3201	1.1728	0.6866	0.094*	
C7	0.2573 (8)	1.1848 (4)	0.5552 (4)	0.0927 (14)	
H7	0.2337	1.2596	0.5562	0.111*	
C8	0.2390 (7)	1.1264 (4)	0.4742 (3)	0.0899 (14)	
H8	0.2024	1.1621	0.4209	0.108*	
C9	0.2744 (6)	1.0171 (3)	0.4721 (3)	0.0706 (11)	
H9	0.2614	0.9793	0.4171	0.085*	
C10	0.3302 (4)	0.9598 (3)	0.5510 (2)	0.0520 (8)	
C11	0.3571 (5)	0.7794 (3)	0.4722 (2)	0.0568 (9)	
H11	0.3241	0.8133	0.4175	0.068*	
C12	0.3841 (5)	0.6111 (3)	0.3898 (3)	0.0577 (9)	
C13	0.3368 (6)	0.5017 (3)	0.3975 (3)	0.0743 (10)	
C14	0.3288 (6)	0.4347 (4)	0.3228 (5)	0.0930 (15)	
H14	0.2959	0.3616	0.3296	0.112*	
C15	0.3693 (7)	0.4754 (5)	0.2380 (4)	0.0916 (16)	
H15	0.3627	0.4302	0.1868	0.110*	
C16	0.4199 (6)	0.5835 (4)	0.2282 (3)	0.0830 (13)	

H16	0.4484	0.6111	0.1704	0.100*	
C17	0.4283 (5)	0.6505 (3)	0.3039 (3)	0.0689 (11)	
H17	0.4642	0.7231	0.2971	0.083*	
F1	0.2925 (4)	0.4629 (2)	0.4811 (2)	0.1113 (10)	
N1	0.3887 (4)	0.6741 (2)	0.4711 (2)	0.0612 (8)	
H1B	0.4135	0.6418	0.5220	0.073*	0.43 (6)
O1	0.4613 (5)	0.6891 (2)	0.6408 (2)	0.0823 (9)	
H1A	0.4242	0.6584	0.5943	0.124*	0.57 (6)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0475 (17)	0.062 (2)	0.048 (2)	-0.0027 (16)	-0.0010 (14)	-0.0002 (18)
C2	0.055 (2)	0.076 (3)	0.062 (3)	-0.003 (2)	0.0028 (18)	0.008 (2)
C3	0.068 (2)	0.106 (4)	0.047 (2)	0.005 (2)	0.0024 (19)	0.007 (2)
C4	0.062 (2)	0.099 (3)	0.049 (2)	-0.007 (2)	0.0014 (19)	-0.007 (2)
C5	0.055 (2)	0.072 (3)	0.057 (2)	-0.0125 (18)	0.0069 (17)	-0.010 (2)
C6	0.081 (3)	0.076 (3)	0.077 (3)	-0.012 (2)	0.015 (2)	-0.020 (3)
C7	0.118 (4)	0.057 (2)	0.103 (4)	-0.001 (3)	0.016 (3)	-0.004 (3)
C8	0.127 (4)	0.062 (3)	0.081 (3)	0.013 (3)	-0.003 (3)	0.006 (2)
C9	0.096 (3)	0.061 (2)	0.055 (2)	0.004 (2)	-0.003 (2)	0.0001 (18)
C10	0.0468 (18)	0.056 (2)	0.053 (2)	-0.0045 (15)	0.0006 (15)	0.0017 (18)
C11	0.055 (2)	0.057 (2)	0.059 (2)	0.0004 (17)	-0.0048 (18)	0.0079 (17)
C12	0.0473 (18)	0.052 (2)	0.073 (3)	0.0021 (16)	-0.0060 (18)	-0.006 (2)
C13	0.062 (2)	0.056 (3)	0.104 (2)	0.0001 (19)	-0.002 (2)	-0.005 (2)
C14	0.074 (3)	0.055 (3)	0.151 (5)	-0.001 (2)	-0.011 (3)	-0.010 (3)
C15	0.075 (3)	0.087 (4)	0.114 (4)	0.014 (3)	-0.007 (3)	-0.040 (3)
C16	0.075 (3)	0.094 (4)	0.080 (3)	0.014 (3)	0.000 (2)	-0.017 (3)
C17	0.067 (2)	0.070 (3)	0.070 (2)	0.007 (2)	0.001 (2)	-0.008 (2)
F1	0.124 (2)	0.0774 (17)	0.133 (2)	-0.0092 (16)	0.007 (2)	0.0304 (16)
N1	0.0596 (18)	0.057 (2)	0.067 (2)	0.0015 (15)	-0.0012 (15)	0.0019 (16)
O1	0.098 (2)	0.0781 (19)	0.0710 (19)	0.0091 (18)	0.0012 (16)	0.0213 (16)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C1—C11	1.405 (5)	C9—H9	0.93
C1—C2	1.424 (5)	C11—N1	1.306 (4)
C1—C10	1.446 (5)	C11—H11	0.93
C2—O1	1.314 (5)	C12—C17	1.380 (5)
C2—C3	1.415 (5)	C12—C13	1.384 (5)
C3—C4	1.330 (6)	C12—N1	1.413 (4)
C3—H3	0.93	C13—F1	1.347 (5)
C4—C5	1.426 (6)	C13—C14	1.363 (6)
C4—H4	0.93	C14—C15	1.365 (7)
C5—C6	1.394 (6)	C14—H14	0.93
C5—C10	1.413 (5)	C15—C16	1.378 (7)
C6—C7	1.344 (6)	C15—H15	0.93
C6—H6	0.93	C16—C17	1.375 (6)

C7—C8	1.387 (6)	C16—H16	0.93
C7—H7	0.93	C17—H17	0.93
C8—C9	1.360 (5)	N1—H1B	0.86
C8—H8	0.93	O1—H1A	0.82
C9—C10	1.406 (5)		
C11—C1—C2	119.3 (3)	C9—C10—C1	123.5 (3)
C11—C1—C10	122.0 (3)	C5—C10—C1	119.7 (3)
C2—C1—C10	118.7 (3)	N1—C11—C1	123.4 (3)
O1—C2—C3	119.4 (4)	N1—C11—H11	118.3
O1—C2—C1	121.3 (4)	C1—C11—H11	118.3
C3—C2—C1	119.3 (4)	C17—C12—C13	117.9 (4)
C4—C3—C2	121.6 (4)	C17—C12—N1	124.4 (3)
C4—C3—H3	119.2	C13—C12—N1	117.7 (4)
C2—C3—H3	119.2	F1—C13—C14	120.0 (4)
C3—C4—C5	122.1 (4)	F1—C13—C12	118.2 (4)
C3—C4—H4	118.9	C14—C13—C12	121.8 (5)
C5—C4—H4	118.9	C13—C14—C15	119.7 (5)
C6—C5—C10	120.1 (4)	C13—C14—H14	120.2
C6—C5—C4	121.4 (4)	C15—C14—H14	120.2
C10—C5—C4	118.5 (4)	C14—C15—C16	120.1 (5)
C7—C6—C5	121.2 (4)	C14—C15—H15	120.0
C7—C6—H6	119.4	C16—C15—H15	120.0
C5—C6—H6	119.4	C17—C16—C15	119.9 (5)
C6—C7—C8	119.9 (4)	C17—C16—H16	120.0
C6—C7—H7	120.1	C15—C16—H16	120.0
C8—C7—H7	120.1	C16—C17—C12	120.7 (4)
C9—C8—C7	120.4 (4)	C16—C17—H17	119.7
C9—C8—H8	119.8	C12—C17—H17	119.7
C7—C8—H8	119.8	C11—N1—C12	122.9 (3)
C8—C9—C10	121.7 (4)	C11—N1—H1B	118.6
C8—C9—H9	119.2	C12—N1—H1B	118.6
C10—C9—H9	119.2	C2—O1—H1A	109.5
C9—C10—C5	116.7 (3)		
C11—C1—C2—O1	0.3 (5)	C11—C1—C10—C9	0.3 (5)
C10—C1—C2—O1	179.8 (3)	C2—C1—C10—C9	-179.2 (4)
C11—C1—C2—C3	179.7 (3)	C11—C1—C10—C5	-179.7 (3)
C10—C1—C2—C3	-0.9 (5)	C2—C1—C10—C5	0.8 (5)
O1—C2—C3—C4	179.9 (4)	C2—C1—C11—N1	1.0 (5)
C1—C2—C3—C4	0.5 (6)	C10—C1—C11—N1	-178.5 (3)
C2—C3—C4—C5	-0.1 (6)	C17—C12—C13—F1	-179.7 (4)
C3—C4—C5—C6	180.0 (4)	N1—C12—C13—F1	2.0 (5)
C3—C4—C5—C10	0.1 (5)	C17—C12—C13—C14	-1.7 (6)
C10—C5—C6—C7	-0.1 (6)	N1—C12—C13—C14	-180.0 (4)
C4—C5—C6—C7	180.0 (4)	F1—C13—C14—C15	178.3 (4)
C5—C6—C7—C8	0.5 (8)	C12—C13—C14—C15	0.4 (6)
C6—C7—C8—C9	-0.4 (8)	C13—C14—C15—C16	0.7 (7)

C7—C8—C9—C10	0.0 (8)	C14—C15—C16—C17	-0.5 (7)
C8—C9—C10—C5	0.4 (6)	C15—C16—C17—C12	-0.9 (7)
C8—C9—C10—C1	-179.5 (4)	C13—C12—C17—C16	1.9 (6)
C6—C5—C10—C9	-0.3 (5)	N1—C12—C17—C16	-179.9 (4)
C4—C5—C10—C9	179.6 (3)	C1—C11—N1—C12	-176.7 (3)
C6—C5—C10—C1	179.6 (3)	C17—C12—N1—C11	30.5 (5)
C4—C5—C10—C1	-0.5 (5)	C13—C12—N1—C11	-151.4 (3)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1A···N1	0.82	1.82	2.535 (4)	144
N1—H1B···O1	0.86	1.86	2.535 (4)	134