

## Methyl 2-ethyl-4-hydroxy-2H-1,2-benzothiazine-3-carboxylate 1,1-dioxide

Muhammad Nadeem Arshad,<sup>a</sup>‡ Muhammad Zia-ur-Rehman<sup>b</sup> and Islam Ullah Khan<sup>a\*</sup>

<sup>a</sup>Materials Chemistry Laboratory, Department of Chemistry, GC University, Lahore, Pakistan, and <sup>b</sup>Applied Chemistry Research Centre, PCSIR Laboratories Complex, Lahore 54600, Pakistan

Correspondence e-mail: iukhan.gcu@gmail.com


Received 31 March 2010; accepted 1 April 2010

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$ ;  $R$  factor = 0.054;  $wR$  factor = 0.102; data-to-parameter ratio = 17.9.

In the title compound,  $\text{C}_{12}\text{H}_{13}\text{NO}_5\text{S}$ , the thiazine ring adopts a half chair conformation and an intramolecular  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bond generates an S(6) ring. In the crystal, the molecules are linked by  $\text{C}-\text{H}\cdots\text{O}$  interactions, leading to zigzag chains along the  $b$  axis.

### Related literature

For background to the biological properties of thiazines, see: Zia-ur-Rehman *et al.* (2005, 2006). For related structures, see: Arshad *et al.* (2009a,b). For graph-set notation, see: Bernstein, *et al.* (1995).



### Experimental

#### Crystal data

$\text{C}_{12}\text{H}_{13}\text{NO}_5\text{S}$

$M_r = 283.29$

Orthorhombic,  $Pna2_1$

$a = 7.2460 (6)\text{ \AA}$

$b = 20.548 (2)\text{ \AA}$

$c = 8.5710 (8)\text{ \AA}$

$V = 1276.1 (2)\text{ \AA}^3$

$Z = 4$

Mo  $K\alpha$  radiation

$\mu = 0.27\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.24 \times 0.14 \times 0.10\text{ mm}$

#### Data collection

Bruker KAPPA APEXII CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2007)

$T_{\min} = 0.938$ ,  $T_{\max} = 0.974$

14253 measured reflections

3148 independent reflections

1270 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.133$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$

$wR(F^2) = 0.102$

$S = 0.95$

3148 reflections

176 parameters

1 restraint

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.22\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.25\text{ e \AA}^{-3}$

Absolute structure: Flack (1983),

1464 Friedel pairs

Flack parameter: 0.07 (12)

**Table 1**

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3—H3 $\cdots$ O4	0.82	1.92	2.628 (5)	144
C4—H4 $\cdots$ O4 <sup>i</sup>	0.93	2.55	3.395 (5)	152
C10—H10B $\cdots$ O2 <sup>ii</sup>	0.96	2.52	3.320 (5)	141

Symmetry codes: (i)  $x - \frac{1}{2}, -y + \frac{1}{2}, z + 1$ ; (ii)  $-x + 1, -y, z - \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *X-SEED* (Barbur, 2001); *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the Higher Education Commission (HEC) of Pakistan for providing a grant under the project ‘Strengthening the Materials Chemistry Laboratory at GC University, Lahore’. MNA also acknowledges the HEC for providing a fellowship under the International Research Support Initiative Program (IRSIP).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5390).

### References

- Arshad, M. N., Zia-ur-Rehman, M. & Khan, I. U. (2009a). *Acta Cryst. E65*, o3025.
- Arshad, M. N., Zia-ur-Rehman, M. & Khan, I. U. (2009b). *Acta Cryst. E65*, o3077.
- Barbur, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2007). *SADABS*, *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Zia-ur-Rehman, M., Choudary, J. A. & Ahmad, S. (2005). *Bull. Korean Chem. Soc.* **26**, 1771–1775.
- Zia-ur-Rehman, M., Choudary, J. A., Ahmad, S. & Siddiqui, H. L. (2006). *Chem. Pharm. Bull.* **54**, 1175–1178.

‡ Current address: Department of Chemistry, Georgetown University, 37th and O St NW, Washington DC 20057-212 USA.

# supporting information

*Acta Cryst.* (2010). E66, o1070 [https://doi.org/10.1107/S1600536810012365]

## Methyl 2-ethyl-4-hydroxy-2H-1,2-benzothiazine-3-carboxylate 1,1-dioxide


**Muhammad Nadeem Arshad, Muhammad Zia-ur-Rehman and Islam Ullah Khan**

### S1. Comment

Under the heading of synthesis and X-ray studies and biological evaluation of thiazine related heterocycles our group has already reported biological applications (Zia-ur-Rehman *et al.*, 2005, 2006) and the crystal structures of 1,2-benzothiazine derivatives (Arshad *et al.*, 2009a,b) II & III. The title compound Methyl 2-ethyl-4-hydroxy-2H-1,2-benzothiazine-3-carboxylate 1,1-dioxide (I) is different only in H-alkylation. The hydrogen bonding interactions in I, II and III are pretty much common. The intramolecular O—H···O interaction tend to rise the six membered ring motif R<sub>1</sub>(6) (Bernstein, *et al.*, 1995) which is almost planar with the r.m.s deviaton of 0.0156 Å and inclined at dihedral angle of 25.1 (6)<sup>o</sup> & 17.5 (6)<sup>o</sup> with respect to the thiazine and benzene ring respectively. The half chair shaped thiazine ring exhibits a maximum deviation from the least square plane measure 0.347 (2) Å for S1 and 0.345 (2) Å for C1. The intermolecular C—H···O hydrogen bonding forms zig-zag network along the *b* axes. The bond lengths and bond angles are compareable with the molecules II and III.


### S2. Experimental

Ethyl iodide (250 mg, 1.6 mmol) was added drop wise to the mixture of methyl 4-hydroxy-2H-1,2-benzothiazine-3-carboxylate-1,1-dioxide (350 mg, 1.37 mmol), anhydrous potassium carbonate (161 mg, 1.6 mmol) and dimethyl-formamide (5 ml) in a round bottom flask. Contents were stirred at room temperature for 5 h under nitrogen atmosphere and poured over ice cooled water (100 ml) resulting white precipitates, which was filtered and washed with cold water. Colourless needles of (I) were obtained by re-crystallization from a methanol solution under slow evaporation.



**Figure 1**

The molecular structure of (I) showing 50% displacement ellipsoids.

**Figure 2**

Unit cell packing for (I) showing the inter and intramolecular hydrogen bondings using dashed lines. Hydrogen atoms have been omitted for clarity.

### Methyl 2-ethyl-4-hydroxy-2*H*-1,2-benzothiazine-3-carboxylate 1,1-dioxide

#### Crystal data

$C_{12}H_{13}NO_5S$   
 $M_r = 283.29$   
Orthorhombic,  $Pna2_1$   
Hall symbol: P 2c -2n  
 $a = 7.2460 (6) \text{ \AA}$   
 $b = 20.548 (2) \text{ \AA}$   
 $c = 8.5710 (8) \text{ \AA}$   
 $V = 1276.1 (2) \text{ \AA}^3$   
 $Z = 4$

$F(000) = 592$   
 $D_x = 1.475 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
Cell parameters from 733 reflections  
 $\theta = 2.6\text{--}17.3^\circ$   
 $\mu = 0.27 \text{ mm}^{-1}$   
 $T = 296 \text{ K}$   
Cut needle, colourless  
 $0.24 \times 0.14 \times 0.10 \text{ mm}$

*Data collection*

Bruker KAPPA APEXII CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(*SADABS*; Bruker, 2007)  
 $T_{\min} = 0.938$ ,  $T_{\max} = 0.974$

14253 measured reflections  
3148 independent reflections  
1270 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.133$   
 $\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 2.0^\circ$   
 $h = -9 \rightarrow 9$   
 $k = -26 \rightarrow 27$   
 $l = -11 \rightarrow 11$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.054$   
 $wR(F^2) = 0.102$   
 $S = 0.95$   
3148 reflections  
176 parameters  
1 restraint  
Primary atom site location: structure-invariant  
direct methods  
Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0219P)^2]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$   
Extinction correction: *SHELXL*,  
 $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$   
Extinction coefficient: 0.0062 (9)  
Absolute structure: Flack (1983), 1464 Friedel  
pairs  
Absolute structure parameter: 0.07 (12)

*Special details*

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.19292 (16)	0.06952 (6)	0.49242 (15)	0.0514 (4)
O1	0.0578 (4)	0.10023 (16)	0.3936 (4)	0.0648 (10)
O2	0.1809 (4)	0.00107 (13)	0.5198 (4)	0.0716 (10)
O3	0.3931 (4)	0.26033 (12)	0.5272 (4)	0.0532 (8)
H3	0.4444	0.2709	0.4458	0.080*
O4	0.5638 (4)	0.23848 (15)	0.2618 (4)	0.0563 (10)
O5	0.5818 (4)	0.13380 (14)	0.1843 (4)	0.0550 (9)
N1	0.3974 (4)	0.08716 (17)	0.4241 (4)	0.0413 (9)
C1	0.1976 (6)	0.1103 (2)	0.6695 (5)	0.0422 (11)
C2	0.1285 (6)	0.0824 (2)	0.8054 (6)	0.0536 (14)
H2	0.0732	0.0416	0.8027	0.064*
C3	0.1429 (6)	0.1160 (3)	0.9446 (6)	0.0560 (15)
H3A	0.1030	0.0970	1.0370	0.067*

C4	0.2163 (6)	0.1776 (2)	0.9459 (5)	0.0589 (15)
H4	0.2201	0.2009	1.0389	0.071*
C5	0.2849 (6)	0.2056 (2)	0.8105 (6)	0.0560 (14)
H5	0.3341	0.2474	0.8139	0.067*
C6	0.2811 (6)	0.1722 (2)	0.6708 (6)	0.0396 (11)
C7	0.3701 (5)	0.1955 (2)	0.5296 (6)	0.0406 (11)
C8	0.4294 (6)	0.1557 (2)	0.4160 (5)	0.0372 (11)
C9	0.5296 (6)	0.1805 (3)	0.2822 (5)	0.0471 (12)
C10	0.6951 (6)	0.1538 (2)	0.0543 (5)	0.0707 (16)
H10A	0.6288	0.1848	-0.0079	0.106*
H10B	0.7252	0.1165	-0.0083	0.106*
H10C	0.8066	0.1733	0.0926	0.106*
C11	0.5572 (6)	0.0441 (2)	0.4634 (5)	0.0548 (15)
H11A	0.6478	0.0475	0.3805	0.066*
H11B	0.5143	-0.0006	0.4657	0.066*
C12	0.6504 (6)	0.0588 (3)	0.6154 (7)	0.0840 (19)
H12A	0.6928	0.1031	0.6152	0.126*
H12B	0.7537	0.0301	0.6290	0.126*
H12C	0.5646	0.0527	0.6994	0.126*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0554 (7)	0.0474 (7)	0.0516 (7)	-0.0144 (7)	0.0019 (9)	-0.0055 (8)
O1	0.051 (2)	0.083 (3)	0.060 (2)	-0.0095 (18)	-0.010 (2)	-0.003 (2)
O2	0.097 (2)	0.0399 (19)	0.078 (3)	-0.0283 (16)	0.014 (2)	-0.010 (2)
O3	0.065 (2)	0.033 (2)	0.061 (3)	-0.0045 (14)	0.0097 (19)	0.0007 (19)
O4	0.067 (2)	0.042 (2)	0.059 (2)	-0.0064 (17)	0.0121 (19)	0.0070 (18)
O5	0.068 (2)	0.048 (2)	0.050 (2)	0.0000 (17)	0.0208 (19)	-0.0001 (19)
N1	0.044 (2)	0.037 (2)	0.043 (2)	-0.0036 (17)	0.0052 (18)	-0.0020 (18)
C1	0.038 (3)	0.048 (3)	0.040 (3)	0.000 (2)	0.000 (2)	-0.006 (3)
C2	0.038 (3)	0.056 (4)	0.066 (4)	-0.003 (2)	0.009 (3)	0.005 (3)
C3	0.056 (3)	0.062 (4)	0.050 (4)	0.010 (3)	0.012 (3)	0.008 (3)
C4	0.062 (3)	0.075 (4)	0.039 (4)	0.017 (3)	0.006 (3)	-0.010 (3)
C5	0.061 (3)	0.051 (3)	0.056 (4)	0.002 (3)	0.003 (3)	-0.011 (3)
C6	0.039 (3)	0.035 (3)	0.045 (3)	0.002 (2)	-0.001 (3)	0.000 (3)
C7	0.044 (3)	0.033 (3)	0.045 (3)	-0.005 (2)	-0.003 (2)	-0.001 (3)
C8	0.040 (3)	0.027 (3)	0.045 (3)	-0.004 (2)	-0.005 (2)	0.002 (2)
C9	0.043 (3)	0.058 (3)	0.040 (3)	-0.003 (3)	-0.006 (3)	-0.001 (3)
C10	0.086 (4)	0.065 (4)	0.061 (4)	-0.004 (3)	0.036 (3)	0.002 (3)
C11	0.057 (3)	0.037 (3)	0.070 (4)	0.010 (2)	0.005 (3)	0.010 (3)
C12	0.069 (4)	0.102 (5)	0.081 (5)	0.019 (3)	-0.013 (3)	0.015 (4)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

S1—O2	1.429 (3)	C4—C5	1.388 (6)
S1—O1	1.441 (3)	C4—H4	0.9300
S1—N1	1.634 (3)	C5—C6	1.381 (6)

S1—C1	1.734 (5)	C5—H5	0.9300
O3—C7	1.343 (4)	C6—C7	1.452 (6)
O3—H3	0.8200	C7—C8	1.343 (5)
O4—C9	1.229 (5)	C8—C9	1.450 (6)
O5—C9	1.330 (5)	C10—H10A	0.9600
O5—C10	1.444 (5)	C10—H10B	0.9600
N1—C8	1.428 (5)	C10—H10C	0.9600
N1—C11	1.495 (5)	C11—C12	1.499 (6)
C1—C2	1.392 (6)	C11—H11A	0.9700
C1—C6	1.408 (5)	C11—H11B	0.9700
C2—C3	1.383 (6)	C12—H12A	0.9600
C2—H2	0.9300	C12—H12B	0.9600
C3—C4	1.372 (6)	C12—H12C	0.9600
C3—H3A	0.9300		
O2—S1—O1	119.1 (2)	C1—C6—C7	118.8 (4)
O2—S1—N1	109.44 (18)	C8—C7—O3	123.6 (4)
O1—S1—N1	107.97 (19)	C8—C7—C6	123.0 (4)
O2—S1—C1	109.4 (2)	O3—C7—C6	113.3 (4)
O1—S1—C1	108.4 (2)	C7—C8—N1	120.9 (4)
N1—S1—C1	100.9 (2)	C7—C8—C9	121.3 (4)
C7—O3—H3	109.5	N1—C8—C9	117.9 (4)
C9—O5—C10	116.3 (3)	O4—C9—O5	123.6 (4)
C8—N1—C11	117.9 (3)	O4—C9—C8	123.7 (4)
C8—N1—S1	112.5 (3)	O5—C9—C8	112.7 (4)
C11—N1—S1	119.4 (3)	O5—C10—H10A	109.5
C2—C1—C6	121.3 (4)	O5—C10—H10B	109.5
C2—C1—S1	121.8 (4)	H10A—C10—H10B	109.5
C6—C1—S1	116.9 (4)	O5—C10—H10C	109.5
C3—C2—C1	119.3 (4)	H10A—C10—H10C	109.5
C3—C2—H2	120.4	H10B—C10—H10C	109.5
C1—C2—H2	120.4	N1—C11—C12	115.2 (4)
C4—C3—C2	119.8 (5)	N1—C11—H11A	108.5
C4—C3—H3A	120.1	C12—C11—H11A	108.5
C2—C3—H3A	120.1	N1—C11—H11B	108.5
C3—C4—C5	120.9 (4)	C12—C11—H11B	108.5
C3—C4—H4	119.5	H11A—C11—H11B	107.5
C5—C4—H4	119.5	C11—C12—H12A	109.5
C6—C5—C4	120.8 (4)	C11—C12—H12B	109.5
C6—C5—H5	119.6	H12A—C12—H12B	109.5
C4—C5—H5	119.6	C11—C12—H12C	109.5
C5—C6—C1	117.7 (4)	H12A—C12—H12C	109.5
C5—C6—C7	123.3 (4)	H12B—C12—H12C	109.5
O2—S1—N1—C8	-171.2 (3)	S1—C1—C6—C7	-4.9 (5)
O1—S1—N1—C8	57.8 (3)	C5—C6—C7—C8	153.4 (4)
C1—S1—N1—C8	-55.9 (3)	C1—C6—C7—C8	-21.9 (6)
O2—S1—N1—C11	-26.6 (4)	C5—C6—C7—O3	-24.2 (6)

O1—S1—N1—C11	−157.6 (3)	C1—C6—C7—O3	160.5 (4)
C1—S1—N1—C11	88.7 (3)	O3—C7—C8—N1	−178.8 (4)
O2—S1—C1—C2	−22.5 (4)	C6—C7—C8—N1	3.8 (6)
O1—S1—C1—C2	108.9 (4)	O3—C7—C8—C9	2.0 (6)
N1—S1—C1—C2	−137.8 (4)	C6—C7—C8—C9	−175.4 (4)
O2—S1—C1—C6	155.2 (3)	C11—N1—C8—C7	−105.7 (4)
O1—S1—C1—C6	−73.4 (4)	S1—N1—C8—C7	39.5 (5)
N1—S1—C1—C6	39.9 (4)	C11—N1—C8—C9	73.6 (5)
C6—C1—C2—C3	−0.3 (7)	S1—N1—C8—C9	−141.2 (3)
S1—C1—C2—C3	177.3 (3)	C10—O5—C9—O4	4.3 (7)
C1—C2—C3—C4	3.3 (7)	C10—O5—C9—C8	−175.3 (4)
C2—C3—C4—C5	−3.2 (7)	C7—C8—C9—O4	−1.5 (7)
C3—C4—C5—C6	0.0 (7)	N1—C8—C9—O4	179.3 (4)
C4—C5—C6—C1	2.9 (6)	C7—C8—C9—O5	178.1 (4)
C4—C5—C6—C7	−172.4 (4)	N1—C8—C9—O5	−1.1 (5)
C2—C1—C6—C5	−2.8 (6)	C8—N1—C11—C12	57.6 (5)
S1—C1—C6—C5	179.5 (3)	S1—N1—C11—C12	−85.1 (4)
C2—C1—C6—C7	172.8 (4)		

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
O3—H3···O4	0.82	1.92	2.628 (5)	144
C4—H4···O4 <sup>i</sup>	0.93	2.55	3.395 (5)	152
C10—H10B···O2 <sup>ii</sup>	0.96	2.52	3.320 (5)	141

Symmetry codes: (i)  $x-1/2, -y+1/2, z+1$ ; (ii)  $-x+1, -y, z-1/2$ .