

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(2,6-Dihydroxy-4-methoxyphenyl)-3-phenylpropan-1-one¹Suchada Chantrapromma,^{a*}§ Jutatip Jeerapong,^b
Thongchai Kruahong,^b Surat Laphookhieo^c and
Hoong-Kun Fun^{d¶}

^aCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand, ^bDepartment of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science and Technology, Suratthani Rajabhat University, Mueang, Surat Thani 84100, Thailand, ^cNatural Products Research Laboratory, School of Science, Mae Fah Luang University, Muang, Chiang Rai 57100, Thailand, and ^dX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia
Correspondence e-mail: suchada.c@psu.ac.th

Received 2 April 2010; accepted 13 April 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.034; wR factor = 0.097; data-to-parameter ratio = 16.4.

The title compound, $\text{C}_{16}\text{H}_{16}\text{O}_4$, a dihydrochalcone, was isolated from the rhizomes of *Etilingera littoralis*. The molecule is twisted with a dihedral angle of $71.69(6)^\circ$ between the two aromatic rings. The propanone unit makes dihedral angles of $4.07(6)$ and $73.56(7)^\circ$, respectively, with the 2,6-dihydroxy-4-methoxyphenyl and phenyl rings. The methoxy group is approximately coplanar with the attached benzene ring with a dihedral angle of $1.74(10)^\circ$. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond generates an $S(6)$ ring motif. In the crystal, intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into chains along $[201]$. A $\pi-\pi$ interaction with a centroid-centroid distance of $3.5185(6)$ Å is also observed.


Related literature

For details of hydrogen-bond motifs, see: Bernstein *et al.* (1995). For bond-length data, see: Allen *et al.* (1987). For background to dihydrochalcones and their activities, see: Nilsson (1961); Nowakowska (2007); Portet *et al.* (2007). For Zingiberaceae plants, see: Chuakul & Boonpleng (2003); Reanmongkol *et al.* (2006); Sirirugsa (1999); Tewtrakul, Subhadhirasakul & Kummee (2003); Tewtrakul, Subhadhirasakul, Puripattanavong & Panphadung (2003). For a related structure, see: Ng *et al.* (2005). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).

¹This paper is dedicated to Her Royal Highness Princess Maha Chakri Sirindhorn of Thailand on the occasion of her 55th Birthday Anniversary which fell on April 2nd, 2010.

§ Thomson Reuters ResearcherID: A-5085-2009.

¶ Additional correspondence author, e-mail: hkfun@usm.my. Thomson Reuters ResearcherID: A-3561-2009.



Experimental

Crystal data

$\text{C}_{16}\text{H}_{16}\text{O}_4$
 $M_r = 272.29$
Monoclinic, Cc
 $a = 7.2142(6)$ Å
 $b = 30.522(2)$ Å
 $c = 6.5587(5)$ Å
 $\beta = 107.267(2)^\circ$
 $V = 1379.09(18)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 100$ K
 $0.46 \times 0.34 \times 0.18$ mm

Data collection

Bruker APEX DUO CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.958$, $T_{\max} = 0.983$
17744 measured reflections
3044 independent reflections
2940 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.097$
 $S = 1.08$
3044 reflections
186 parameters
2 restraints
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.34$ e Å⁻³
 $\Delta\rho_{\min} = -0.40$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}2-\text{H}1\text{O}2\cdots\text{O}1$	0.82	1.71	2.4576 (11)	150
$\text{O}4-\text{H}1\text{O}4\cdots\text{O}2^i$	0.80 (3)	1.90 (3)	2.6920 (10)	175 (3)

Symmetry code: (i) $x + 1, -y, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

JJ and TK thank the Center of Excellence for Innovation in Chemistry (PERCH-CIC) and the Faculty of Science and Technology, Suratthani Rajabhat University, for financial support. The Natural Products Research Laboratory, School of Science, Mae Fah Luang University, is gratefully acknowledged for laboratory facilities. SC thanks the Prince of Songkla University for financial support. Mr Nawong Boonnak is acknowledged for providing useful information. The authors also thank the Malaysian Government and Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2537).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2009). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chuakul, W. & Boonpleng, A. (2003). *J. Med. Plants*, **10**, 33–35.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Ng, S.-L., Razak, I. A., Fun, H.-K., Boonsri, S., Chantrapromma, S. & Prawat, U. (2005). *Acta Cryst.* **E61**, o3234–o3236.
- Nilsson, M. (1961). *Acta Chem. Scand.* **15**, 154–158.
- Nowakowska, Z. (2007). *Eur. J. Med. Chem.* **42**, 125–137.
- Portet, B., Fabre, N., Roumy, V., Gornitzka, H., Bourdy, G., Chevalley, S., Sauvain, M., Valentin, A. & Moulis, C. (2007). *Phytochemistry*, **68**, 1312–1320.
- Reanmongkol, W., Subhadhirasakul, S., Khaisombat, N., Fuengnawakit, P., Jantasila, S. & Khamjun, A. (2006). *Songklanakarin J. Sci. Technol.* **25**, 999–1008.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sirirugsa, P. (1999). *Thai Zingiberaceae: Species diversity and their uses*, <http://www.iupac.org/symposia/proceedings/phuket97/sirirugsa.html>.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Tewtrakul, S., Subhadhirasakul, S. & Kummee, S. (2003). *Songklanakarin J. Sci. Technol.* **25**, 239–243.
- Tewtrakul, S., Subhadhirasakul, S., Puripattanavong, J. & Panphadung, T. (2003). *Songklanakarin J. Sci. Technol.* **25**, 503–508.

supporting information

Acta Cryst. (2010). E66, o1120–o1121 [https://doi.org/10.1107/S1600536810013590]

1-(2,6-Dihydroxy-4-methoxyphenyl)-3-phenylpropan-1-one

Suchada Chantrapromma, Jutatip Jeerapong, Thongchai Kruahong, Surat Laphookhieo and Hoong-Kun Fun

S1. Comment

Zingiberaceae plants are the ground plants of tropical forests. Many of them are used for food, spices, medicines, dyes, perfume and aesthetics (Sirirugsa, 1999). Secondary metabolites from Zingiberaceae plants have found to be anti-inflammatory (Reanmongkol *et al.*, 2006), HIV-1 protease inhibitory (Tewtrakul, Subhadhirasakul & Kummee, 2003; Tewtrakul, Subhadhirasakul, Puripattanavong & Panphadung, 2003). *Etilingera littoralis* is one of the Zingiberaceae plants and its decoction of the rhizomes has been used for the treatment of stomachache, carminative and heart tonic (Chuakul & Boonpleng, 2003). As part of our study of chemical constituents and bioactive compounds from the rhizomes of *Etilingera littoralis* which were collected from Surat Thani province in the southern of Thailand, the title dihydrochalcone, (I), was isolated. Herein we report its crystal structure. The title compound was found to possess antibacterial (Nowakowska, 2007) and antiplasmodial activities (Portet *et al.*, 2007).

The molecule of the title dihydrochalcone (Fig. 1), C₁₆H₁₆O₄, is twisted as the dihedral angle between the 2,6-dihydroxy-4-methoxyphenyl and phenyl rings is 71.69 (6)°. Whereas the 1-propanone unit (C7–C9/O1) makes the dihedral angles of 4.07 (6) and 73.56 (7)° with the C1–C6 benzene and C10–C15 phenyl rings, respectively. The two hydroxy and a methoxy groups are co-planar with the attached benzene ring with the *r.m.s.* of 0.0078 (1) Å for the ten non H atoms and the torsion angle C16–O3–C3–C2 = -1.66 (15)°. An intramolecular O2—H1O2···O1 hydrogen bond generates an S(6) ring motif (Bernstein *et al.*, 1995) (Fig. 1 and Table 1). The bond distances are of normal values (Allen *et al.*, 1987) and are comparable with the closely related structure (Ng *et al.*, 2005).

In the crystal packing (Fig. 2), O—H···O hydrogen bonds (Table 1) formed between the two hydroxy groups link the molecules into chains along the [201] direction in which the adjacent chains are in anti-parallel manner. A π - π interaction with Cg1···Cg1 distance of 3.5185 (6) Å was observed (symmetry code x, -y, -1/2+z); Cg1 is the centroid of the C1–C6 benzene ring.

S2. Experimental

The fresh rhizomes of *E. littoralis* (3.89 kg) were chopped and extracted with 50% CH₂Cl₂-MeOH, over the period of 3 days at room temperature. The extraction was filtered and evaporated to dryness under reduced pressure to give crude extract which was further partitioned with water and CH₂Cl₂ to afford the dichloromethane extract (22.88 g). The portion of dichloromethane extract (11.80 g) was subjected to quick column chromatography (QCC) on silica gel eluting with a gradient of EtOAc-hexane to give thirteen fractions. Fraction F8 (322.4 mg) was washed with 20% CH₂Cl₂-hexane yielding solid which was further separated by column chromatography on silica gel with 70% CH₂Cl₂-hexane to give compound (I) (50.1 mg). Yellow block-shaped single crystals of the compound (I) suitable for X-ray structure determination were obtained from ethyl acetate by slow evaporation at room temperature after a few days, Mp 443 K. The NMR spectral data were consistent with the X-ray structure.

S3. Refinement

Hydroxy H atoms attached to O4 was located from a difference map and isotropically refined. The remaining H atoms were placed in calculated positions, with $d(\text{O—H}) = 0.82 \text{ \AA}$ and $d(\text{C—H}) = 0.93 \text{ \AA}$ for aromatic, 0.97 \AA for CH_2 and 0.96 \AA for CH_3 atoms. The $U_{\text{iso}}(\text{H})$ values were constrained to be $1.5U_{\text{eq}}$ of the carrier atom for hydroxy and methyl H atoms and $1.2U_{\text{eq}}$ for the remaining H atoms. A rotating group model was used for the methyl groups. The highest residual electron density peak is located at 0.69 \AA from C1 and the deepest hole is located at 0.84 \AA from C7. A total of 2607 Friedel pairs were merged before final refinement as there is no large anomalous dispersion for the determination of the absolute configuration.




Figure 1

The molecular structure of the title compound, with 50% probability displacement ellipsoids and the atom-numbering scheme.




Figure 2

The crystal packing of the title compound viewed down the b axis, showing chains running along the $[201]$ direction. Hydrogen bonds are shown as dashed lines.

1-(2,6-Dihydroxy-4-methoxyphenyl)-3-phenylpropan-1-one

Crystal data

$C_{16}H_{16}O_4$

$M_r = 272.29$

Monoclinic, Cc

Hall symbol: $C -2yc$

$a = 7.2142$ (6) Å

$b = 30.522$ (2) Å

$c = 6.5587$ (5) Å

$\beta = 107.267$ (2)°

$V = 1379.09$ (18) Å³

$Z = 4$

$F(000) = 576$

$D_x = 1.311$ Mg m⁻³

Melting point: 443 K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3044 reflections

$\theta = 2.7\text{--}35.0^\circ$

$\mu = 0.09$ mm⁻¹

$T = 100$ K

Block, yellow

$0.46 \times 0.34 \times 0.18$ mm

Data collection

Bruker APEX DUO CCD area-detector
diffractometer
Radiation source: sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)
 $T_{\min} = 0.958$, $T_{\max} = 0.983$

17744 measured reflections
3044 independent reflections
2940 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$
 $\theta_{\max} = 35.0^\circ$, $\theta_{\min} = 2.7^\circ$
 $h = -11 \rightarrow 11$
 $k = -49 \rightarrow 48$
 $l = -10 \rightarrow 10$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.097$
 $S = 1.08$
3044 reflections
186 parameters
2 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0648P)^2 + 0.1812P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.40 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.57929 (10)	0.09273 (3)	0.16110 (12)	0.01772 (14)
O2	0.40256 (10)	0.02283 (2)	0.09183 (12)	0.01659 (14)
H1O2	0.4212	0.0494	0.0997	0.025*
O3	0.76008 (12)	-0.11076 (2)	0.22402 (13)	0.01827 (13)
O4	1.09458 (10)	0.02313 (2)	0.35682 (12)	0.01540 (13)
H1O4	1.186 (4)	0.0088 (8)	0.419 (4)	0.043 (6)*
C1	0.57479 (11)	0.00170 (3)	0.15654 (13)	0.01230 (14)
C2	0.56979 (13)	-0.04392 (3)	0.15433 (14)	0.01400 (15)
H2A	0.4524	-0.0589	0.1102	0.017*
C3	0.74533 (14)	-0.06642 (3)	0.21997 (15)	0.01307 (13)
C4	0.92225 (13)	-0.04409 (3)	0.28721 (14)	0.01272 (14)
H4A	1.0381	-0.0597	0.3298	0.015*
C5	0.92521 (11)	0.00137 (3)	0.29051 (12)	0.01140 (13)
C6	0.75073 (14)	0.02604 (3)	0.22318 (15)	0.01112 (13)

C7	0.74075 (13)	0.07385 (3)	0.21838 (14)	0.01271 (13)
C8	0.91965 (13)	0.10221 (3)	0.27702 (14)	0.01421 (14)
H8A	1.0023	0.0936	0.1911	0.017*
H8B	0.9911	0.0971	0.4255	0.017*
C9	0.87479 (16)	0.15107 (3)	0.24454 (17)	0.01880 (16)
H9A	0.7893	0.1559	0.1013	0.023*
H9B	0.8081	0.1608	0.3448	0.023*
C10	1.05757 (16)	0.17743 (3)	0.27659 (18)	0.01983 (18)
C11	1.1302 (2)	0.20381 (4)	0.4562 (2)	0.0303 (2)
H11A	1.0662	0.2049	0.5601	0.036*
C12	1.2988 (3)	0.22868 (4)	0.4815 (3)	0.0440 (4)
H12A	1.3455	0.2463	0.6013	0.053*
C13	1.3961 (2)	0.22708 (5)	0.3287 (3)	0.0453 (4)
H13A	1.5074	0.2438	0.3455	0.054*
C14	1.3276 (2)	0.20066 (5)	0.1517 (3)	0.0396 (3)
H14A	1.3938	0.1992	0.0499	0.048*
C15	1.15876 (19)	0.17616 (4)	0.1255 (2)	0.0273 (2)
H15A	1.1128	0.1587	0.0049	0.033*
C16	0.58350 (17)	-0.13568 (3)	0.16199 (18)	0.0230 (2)
H16A	0.6136	-0.1664	0.1775	0.034*
H16B	0.5142	-0.1294	0.0157	0.034*
H16C	0.5046	-0.1279	0.2513	0.034*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0124 (3)	0.0166 (3)	0.0227 (3)	0.0027 (2)	0.0030 (2)	0.0000 (3)
O2	0.0085 (3)	0.0182 (3)	0.0212 (3)	0.0004 (2)	0.0016 (2)	-0.0017 (2)
O3	0.0201 (3)	0.0120 (3)	0.0216 (3)	-0.0026 (2)	0.0044 (2)	-0.0005 (2)
O4	0.0072 (2)	0.0147 (3)	0.0221 (3)	-0.0008 (2)	0.0010 (2)	0.0007 (2)
C1	0.0087 (3)	0.0161 (3)	0.0118 (3)	-0.0003 (3)	0.0025 (2)	-0.0004 (3)
C2	0.0119 (3)	0.0157 (3)	0.0139 (3)	-0.0030 (3)	0.0030 (3)	-0.0011 (3)
C3	0.0144 (3)	0.0129 (3)	0.0116 (3)	-0.0014 (3)	0.0034 (2)	-0.0004 (3)
C4	0.0111 (3)	0.0132 (3)	0.0134 (3)	0.0002 (3)	0.0029 (2)	0.0002 (3)
C5	0.0089 (3)	0.0135 (3)	0.0114 (3)	-0.0005 (3)	0.0025 (2)	0.0002 (3)
C6	0.0087 (3)	0.0128 (3)	0.0114 (3)	-0.0007 (3)	0.0022 (2)	-0.0004 (3)
C7	0.0117 (3)	0.0137 (3)	0.0124 (3)	0.0003 (3)	0.0032 (2)	-0.0002 (3)
C8	0.0130 (3)	0.0125 (3)	0.0161 (3)	-0.0013 (3)	0.0028 (3)	-0.0007 (3)
C9	0.0179 (4)	0.0126 (3)	0.0242 (4)	0.0011 (3)	0.0036 (3)	-0.0009 (3)
C10	0.0203 (4)	0.0112 (3)	0.0232 (4)	-0.0016 (3)	-0.0010 (3)	0.0012 (3)
C11	0.0361 (6)	0.0168 (4)	0.0293 (5)	-0.0027 (4)	-0.0035 (4)	-0.0040 (4)
C12	0.0459 (8)	0.0190 (5)	0.0472 (8)	-0.0115 (5)	-0.0170 (6)	0.0002 (5)
C13	0.0297 (6)	0.0271 (6)	0.0637 (10)	-0.0136 (5)	-0.0098 (6)	0.0190 (6)
C14	0.0266 (5)	0.0360 (7)	0.0532 (8)	-0.0074 (5)	0.0073 (6)	0.0202 (6)
C15	0.0251 (5)	0.0238 (5)	0.0308 (5)	-0.0043 (4)	0.0052 (4)	0.0054 (4)
C16	0.0263 (5)	0.0178 (4)	0.0227 (4)	-0.0090 (4)	0.0040 (4)	-0.0014 (3)

Geometric parameters (Å, °)

O1—C7	1.2531 (11)	C8—H8B	0.9700
O2—C1	1.3516 (11)	C9—C10	1.5052 (14)
O2—H1O2	0.8200	C9—H9A	0.9700
O3—C3	1.3574 (11)	C9—H9B	0.9700
O3—C16	1.4349 (13)	C10—C11	1.3943 (15)
O4—C5	1.3445 (10)	C10—C15	1.3951 (17)
O4—H1O4	0.79 (3)	C11—C12	1.401 (2)
C1—C2	1.3928 (13)	C11—H11A	0.9300
C1—C6	1.4230 (12)	C12—C13	1.384 (3)
C2—C3	1.3917 (14)	C12—H12A	0.9300
C2—H2A	0.9300	C13—C14	1.379 (3)
C3—C4	1.3978 (13)	C13—H13A	0.9300
C4—C5	1.3877 (12)	C14—C15	1.3956 (18)
C4—H4A	0.9300	C14—H14A	0.9300
C5—C6	1.4201 (12)	C15—H15A	0.9300
C6—C7	1.4607 (11)	C16—H16A	0.9600
C7—C8	1.5061 (13)	C16—H16B	0.9600
C8—C9	1.5276 (13)	C16—H16C	0.9600
C8—H8A	0.9700		
C1—O2—H1O2	109.5	C10—C9—C8	111.21 (8)
C3—O3—C16	117.72 (9)	C10—C9—H9A	109.4
C5—O4—H1O4	115.4 (18)	C8—C9—H9A	109.4
O2—C1—C2	117.08 (8)	C10—C9—H9B	109.4
O2—C1—C6	120.02 (8)	C8—C9—H9B	109.4
C2—C1—C6	122.90 (8)	H9A—C9—H9B	108.0
C3—C2—C1	118.14 (8)	C11—C10—C15	118.14 (11)
C3—C2—H2A	120.9	C11—C10—C9	121.36 (11)
C1—C2—H2A	120.9	C15—C10—C9	120.50 (9)
O3—C3—C2	123.84 (9)	C10—C11—C12	120.52 (15)
O3—C3—C4	114.90 (9)	C10—C11—H11A	119.7
C2—C3—C4	121.26 (7)	C12—C11—H11A	119.7
C5—C4—C3	120.05 (8)	C13—C12—C11	120.27 (14)
C5—C4—H4A	120.0	C13—C12—H12A	119.9
C3—C4—H4A	120.0	C11—C12—H12A	119.9
O4—C5—C4	120.48 (8)	C14—C13—C12	119.88 (13)
O4—C5—C6	118.37 (7)	C14—C13—H13A	120.1
C4—C5—C6	121.15 (8)	C12—C13—H13A	120.1
C5—C6—C1	116.50 (7)	C13—C14—C15	119.88 (16)
C5—C6—C7	124.74 (8)	C13—C14—H14A	120.1
C1—C6—C7	118.76 (8)	C15—C14—H14A	120.1
O1—C7—C6	120.09 (8)	C10—C15—C14	121.29 (13)
O1—C7—C8	117.52 (7)	C10—C15—H15A	119.4
C6—C7—C8	122.38 (8)	C14—C15—H15A	119.4
C7—C8—C9	113.30 (7)	O3—C16—H16A	109.5
C7—C8—H8A	108.9	O3—C16—H16B	109.5

C9—C8—H8A	108.9	H16A—C16—H16B	109.5
C7—C8—H8B	108.9	O3—C16—H16C	109.5
C9—C8—H8B	108.9	H16A—C16—H16C	109.5
H8A—C8—H8B	107.7	H16B—C16—H16C	109.5
O2—C1—C2—C3	179.91 (8)	C5—C6—C7—O1	-178.29 (8)
C6—C1—C2—C3	-0.16 (14)	C1—C6—C7—O1	1.64 (14)
C16—O3—C3—C2	-1.66 (15)	C5—C6—C7—C8	2.74 (15)
C16—O3—C3—C4	178.47 (8)	C1—C6—C7—C8	-177.33 (8)
C1—C2—C3—O3	-179.62 (8)	O1—C7—C8—C9	-3.30 (12)
C1—C2—C3—C4	0.24 (14)	C6—C7—C8—C9	175.70 (8)
O3—C3—C4—C5	-179.79 (8)	C7—C8—C9—C10	-172.32 (8)
C2—C3—C4—C5	0.34 (14)	C8—C9—C10—C11	-108.05 (11)
C3—C4—C5—O4	179.38 (7)	C8—C9—C10—C15	72.21 (12)
C3—C4—C5—C6	-1.01 (14)	C15—C10—C11—C12	0.83 (17)
O4—C5—C6—C1	-179.33 (7)	C9—C10—C11—C12	-178.92 (11)
C4—C5—C6—C1	1.05 (13)	C10—C11—C12—C13	-0.5 (2)
O4—C5—C6—C7	0.60 (14)	C11—C12—C13—C14	-0.4 (2)
C4—C5—C6—C7	-179.02 (8)	C12—C13—C14—C15	1.0 (2)
O2—C1—C6—C5	179.46 (8)	C11—C10—C15—C14	-0.27 (17)
C2—C1—C6—C5	-0.46 (14)	C9—C10—C15—C14	179.47 (11)
O2—C1—C6—C7	-0.47 (13)	C13—C14—C15—C10	-0.6 (2)
C2—C1—C6—C7	179.60 (8)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H1O2 \cdots O1	0.82	1.71	2.4576 (11)	150
O4—H1O4 \cdots O2 ⁱ	0.80 (3)	1.90 (3)	2.6920 (10)	175 (3)

Symmetry code: (i) $x+1, -y, z+1/2$.