

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## *tert*-Butyl 3-amino-2-methyl-6,7-dihydro-2*H*-pyrazolo[4,3-*c*]pyridine-5(4*H*)-carboxylate

Xin Guo, Xiao Guang Bai, Yi Liang Li and Yu Cheng Wang\*

Institute of Medicinal Biotechnology, Chinese Academy of Medical Sciences, and Peking Union Medical College, Beijing 100050, People's Republic of China  
Correspondence e-mail: hongzhaoupr@yahoo.com


Received 30 March 2010; accepted 9 April 2010

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.064;  $wR$  factor = 0.170; data-to-parameter ratio = 18.0.

In the molecule of the title compound,  $\text{C}_{12}\text{H}_{20}\text{N}_4\text{O}_2$ , the dihydropiperidine ring assumes a half-chair conformation. In the crystal, classical  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{N}-\text{H}\cdots\text{N}$  intermolecular hydrogen bonds link molecules into double chains along the  $a$  axis.

### Related literature

For the synthesis and properties of related kinase inhibitors, see: Fancelli *et al.* (2005); Gadekar *et al.* (1968).



### Experimental

#### Crystal data

$\text{C}_{12}\text{H}_{20}\text{N}_4\text{O}_2$   
 $M_r = 252.32$   
Triclinic,  $P\bar{1}$   
 $a = 6.3151$  (13) Å  
 $b = 9.3615$  (19) Å  
 $c = 11.215$  (2) Å  
 $\alpha = 85.837$  (4)°  
 $\beta = 86.794$  (4)°  
 $\gamma = 87.733$  (4)°  
 $V = 659.8$  (2) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.09$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.30 \times 0.26 \times 0.16$  mm

#### Data collection

Rigaku, SCXmini diffractometer  
Absorption correction: multi-scan  
(*CrystalClear*; Rigaku, 2005)  
 $T_{\min} = 0.972$ ,  $T_{\max} = 0.985$   
6859 measured reflections  
3008 independent reflections  
1737 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.052$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.064$   
 $wR(F^2) = 0.170$   
 $S = 1.01$   
3008 reflections  
167 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.28$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.36$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N4}-\text{H4A}\cdots\text{O2}^i$	0.86	2.32	3.093 (3)	149
$\text{N4}-\text{H4B}\cdots\text{N2}^{ii}$	0.86	2.57	3.420 (3)	172

Symmetry codes: (i)  $-x, -y + 1, -z + 1$ ; (ii)  $x - 1, y, z$ .

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL/PC* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL/PC*.

This work was supported by the National S&T Major Special Project on Major New Drug Innovation (grant No. 2009ZX09301-003).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2434).

### References

- Fancelli, D. *et al.* (2005). *J. Med. Chem.* **48**, 3080–3084.  
Gadekar, S. M., Johnson, B. D. & Cohen, E. (1968). *J. Med. Chem.* **11**, 616–618.  
Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supporting information

*Acta Cryst.* (2010). E66, o1108 [https://doi.org/10.1107/S1600536810013218]

## *tert*-Butyl 3-amino-2-methyl-6,7-dihydro-2*H*-pyrazolo[4,3-*c*]pyridine-5(4*H*)-carboxylate

Xin Guo, Xiao Guang Bai, Yi Liang Li and Yu Cheng Wang

### S1. Comment

In our ongoing project aimed at the development of potential anticancer kinase inhibitors (Fancelli *et al.*, 2005; Gadekar *et al.*, 1968), we have synthesized the title compound and report its crystal structure herein.

In the molecule of the title compound (Fig. 1), bond lengths and angles are within the expected range. The dehydro-piperidine ring assumes a half-chair conformation, with atoms N1 and C1 displaced from the C2–C5 mean plane by  $-0.4493$  (19) and  $0.293$  (3)° respectively. In the crystal packing (Fig. 2), classical N—H⋯O and N—H⋯N intermolecular hydrogen bonds (Table 1) link molecules into double chains along the *a* axis.

### S2. Experimental

A mixture of *tert*-butyl 3-cyano-4-oxopyrrolidine-1-carboxylate (2.1 g, 10.0 mmol) and methylhydrazine (0.46 g, 10.0 mol) was dissolved in ethanol (50 ml) and stirred at room temperature for 12 hours to give a white precipitate of the title compound. Colourless block crystals suitable for X-ray diffraction were obtained in 5 days by slow evaporation of a methanol solution (15 ml) of 100 mg of the crude product.

### S3. Refinement

All H atoms were placed at calculated positions and refined as riding, with C—H = 0.96–0.97 Å, N—H = 0.86 Å, and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$  or  $1.5U_{\text{eq}}(\text{C})$  for methyl H atoms.




Figure 1

The molecular structure of the title compound, showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.




Figure 2

Packing diagram of the title compound viewed along the  $b$  axis. Intermolecular hydrogen bonds are shown as dashed lines.

*tert*-Butyl 3-amino-2-methyl-6,7-dihydro-2*H*-pyrazolo[4,3-*c*]pyridine-5(4*H*)-carboxylate

*Crystal data*

$C_{12}H_{20}N_4O_2$

$M_r = 252.32$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 6.3151\ (13)\ \text{\AA}$

$b = 9.3615\ (19)\ \text{\AA}$

$c = 11.215\ (2)\ \text{\AA}$

$\alpha = 85.837\ (4)^\circ$

$\beta = 86.794\ (4)^\circ$

$\gamma = 87.733\ (4)^\circ$

$V = 659.8\ (2)\ \text{\AA}^3$

$Z = 2$

$F(000) = 272$

$D_x = 1.270\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 5123 reflections

$\theta = 3.2\text{--}27.5^\circ$

$\mu = 0.09\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Block, colourless

$0.30 \times 0.26 \times 0.16\ \text{mm}$

*Data collection*

Rigaku, SCXmini  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution:  $13.6612\ \text{pixels mm}^{-1}$

$\omega$  scans

Absorption correction: multi-scan

(*CrystalClear*; Rigaku, 2005)

$T_{\min} = 0.972$ ,  $T_{\max} = 0.985$

6859 measured reflections

3008 independent reflections

1737 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.052$

$\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 3.2^\circ$

$h = -8 \rightarrow 8$

$k = -12 \rightarrow 12$

$l = -14 \rightarrow 14$

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.064$

$wR(F^2) = 0.170$

$S = 1.01$

3008 reflections

167 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.081P)^2]$$

where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$$

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3109 (4)	0.0363 (3)	0.6957 (2)	0.0386 (6)
H1A	0.2825	-0.0408	0.6462	0.046*
H1B	0.3436	-0.0061	0.7744	0.046*
C2	0.5027 (4)	0.1173 (3)	0.6406 (2)	0.0398 (6)
H2A	0.5619	0.1713	0.7007	0.048*
H2B	0.6113	0.0498	0.6125	0.048*
C3	0.4344 (4)	0.2180 (2)	0.5375 (2)	0.0331 (6)
C4	0.2211 (4)	0.2453 (3)	0.5144 (2)	0.0329 (5)
C5	0.0400 (4)	0.1836 (3)	0.59148 (19)	0.0385 (6)
H5A	-0.0710	0.2567	0.6030	0.046*
H5B	-0.0191	0.1057	0.5531	0.046*
C6	0.2214 (4)	0.3437 (3)	0.4173 (2)	0.0346 (6)
C7	0.5130 (4)	0.4687 (3)	0.2914 (2)	0.0479 (7)
H7A	0.3984	0.5194	0.2522	0.072*
H7B	0.5974	0.4162	0.2344	0.072*
H7C	0.5997	0.5358	0.3251	0.072*
C8	0.0740 (4)	0.2085 (3)	0.8027 (2)	0.0327 (5)
C9	0.1596 (4)	0.2266 (3)	1.0122 (2)	0.0371 (6)
C10	0.2487 (5)	0.3756 (3)	0.9930 (3)	0.0561 (8)
H10A	0.3934	0.3684	0.9622	0.084*
H10B	0.2431	0.4197	1.0679	0.084*
H10C	0.1660	0.4329	0.9369	0.084*
C11	-0.0678 (4)	0.2267 (3)	1.0624 (2)	0.0486 (7)
H11A	-0.1532	0.2919	1.0136	0.073*
H11B	-0.0739	0.2563	1.1427	0.073*
H11C	-0.1206	0.1319	1.0627	0.073*
C12	0.2991 (5)	0.1310 (3)	1.0927 (2)	0.0564 (8)
H12A	0.2457	0.0360	1.1008	0.085*
H12B	0.2984	0.1687	1.1701	0.085*
H12C	0.4416	0.1279	1.0580	0.085*
N1	0.1203 (3)	0.1310 (2)	0.70667 (17)	0.0353 (5)
N2	0.5648 (3)	0.2928 (2)	0.46032 (17)	0.0388 (5)

N3	0.4281 (3)	0.3700 (2)	0.38587 (17)	0.0386 (5)
N4	0.0539 (3)	0.4037 (2)	0.35407 (19)	0.0507 (6)
H4A	0.0781	0.4606	0.2915	0.061*
H4B	-0.0744	0.3840	0.3773	0.061*
O1	0.1814 (3)	0.15709 (17)	0.89818 (14)	0.0398 (4)
O2	-0.0535 (3)	0.31087 (19)	0.80320 (15)	0.0480 (5)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0362 (14)	0.0381 (14)	0.0400 (14)	0.0063 (11)	0.0030 (11)	-0.0011 (11)
C2	0.0327 (14)	0.0447 (15)	0.0402 (14)	0.0078 (11)	0.0001 (11)	0.0032 (12)
C3	0.0273 (12)	0.0387 (14)	0.0327 (12)	0.0038 (10)	-0.0009 (10)	-0.0015 (11)
C4	0.0268 (12)	0.0436 (14)	0.0285 (12)	0.0024 (10)	-0.0019 (9)	-0.0044 (10)
C5	0.0308 (13)	0.0529 (16)	0.0319 (13)	-0.0022 (11)	-0.0021 (10)	-0.0040 (12)
C6	0.0296 (13)	0.0445 (14)	0.0299 (12)	0.0033 (11)	-0.0042 (10)	-0.0053 (11)
C7	0.0437 (16)	0.0557 (17)	0.0420 (15)	-0.0021 (13)	0.0000 (12)	0.0110 (13)
C8	0.0271 (12)	0.0374 (13)	0.0323 (12)	-0.0002 (11)	0.0005 (10)	0.0037 (11)
C9	0.0413 (15)	0.0385 (14)	0.0312 (13)	-0.0011 (11)	-0.0013 (10)	-0.0018 (11)
C10	0.0587 (19)	0.0517 (18)	0.0591 (18)	-0.0151 (14)	-0.0054 (14)	-0.0026 (14)
C11	0.0493 (17)	0.0514 (17)	0.0440 (15)	-0.0054 (13)	0.0103 (13)	-0.0028 (13)
C12	0.065 (2)	0.0602 (19)	0.0440 (16)	0.0080 (15)	-0.0120 (14)	0.0000 (14)
N1	0.0302 (11)	0.0431 (12)	0.0317 (10)	0.0020 (9)	0.0024 (8)	-0.0015 (9)
N2	0.0274 (11)	0.0496 (13)	0.0376 (11)	0.0047 (9)	-0.0007 (9)	0.0052 (10)
N3	0.0292 (11)	0.0487 (13)	0.0360 (11)	0.0037 (9)	-0.0020 (9)	0.0067 (10)
N4	0.0365 (13)	0.0750 (17)	0.0386 (12)	0.0059 (11)	-0.0059 (10)	0.0092 (11)
O1	0.0475 (11)	0.0406 (10)	0.0307 (9)	0.0096 (8)	-0.0045 (7)	-0.0031 (7)
O2	0.0519 (12)	0.0496 (11)	0.0403 (10)	0.0201 (9)	-0.0040 (8)	0.0006 (8)

*Geometric parameters (Å, °)*

C1—N1	1.471 (3)	C8—O2	1.227 (3)
C1—C2	1.532 (3)	C8—O1	1.349 (3)
C1—H1A	0.9700	C8—N1	1.354 (3)
C1—H1B	0.9700	C9—O1	1.474 (3)
C2—C3	1.509 (3)	C9—C11	1.513 (3)
C2—H2A	0.9700	C9—C12	1.516 (3)
C2—H2B	0.9700	C9—C10	1.521 (3)
C3—N2	1.340 (3)	C10—H10A	0.9600
C3—C4	1.396 (3)	C10—H10B	0.9600
C4—C6	1.375 (3)	C10—H10C	0.9600
C4—C5	1.502 (3)	C11—H11A	0.9600
C5—N1	1.460 (3)	C11—H11B	0.9600
C5—H5A	0.9700	C11—H11C	0.9600
C5—H5B	0.9700	C12—H12A	0.9600
C6—N3	1.360 (3)	C12—H12B	0.9600
C6—N4	1.384 (3)	C12—H12C	0.9600
C7—N3	1.448 (3)	N2—N3	1.382 (3)

C7—H7A	0.9600	N4—H4A	0.8600
C7—H7B	0.9600	N4—H4B	0.8600
C7—H7C	0.9600		
N1—C1—C2	111.82 (19)	O1—C9—C11	111.03 (19)
N1—C1—H1A	109.3	O1—C9—C12	102.7 (2)
C2—C1—H1A	109.3	C11—C9—C12	110.1 (2)
N1—C1—H1B	109.3	O1—C9—C10	108.8 (2)
C2—C1—H1B	109.3	C11—C9—C10	113.5 (2)
H1A—C1—H1B	107.9	C12—C9—C10	110.2 (2)
C3—C2—C1	109.5 (2)	C9—C10—H10A	109.5
C3—C2—H2A	109.8	C9—C10—H10B	109.5
C1—C2—H2A	109.8	H10A—C10—H10B	109.5
C3—C2—H2B	109.8	C9—C10—H10C	109.5
C1—C2—H2B	109.8	H10A—C10—H10C	109.5
H2A—C2—H2B	108.2	H10B—C10—H10C	109.5
N2—C3—C4	112.4 (2)	C9—C11—H11A	109.5
N2—C3—C2	125.5 (2)	C9—C11—H11B	109.5
C4—C3—C2	122.1 (2)	H11A—C11—H11B	109.5
C6—C4—C3	105.4 (2)	C9—C11—H11C	109.5
C6—C4—C5	130.6 (2)	H11A—C11—H11C	109.5
C3—C4—C5	123.8 (2)	H11B—C11—H11C	109.5
N1—C5—C4	108.28 (18)	C9—C12—H12A	109.5
N1—C5—H5A	110.0	C9—C12—H12B	109.5
C4—C5—H5A	110.0	H12A—C12—H12B	109.5
N1—C5—H5B	110.0	C9—C12—H12C	109.5
C4—C5—H5B	110.0	H12A—C12—H12C	109.5
H5A—C5—H5B	108.4	H12B—C12—H12C	109.5
N3—C6—C4	106.7 (2)	C8—N1—C5	118.53 (19)
N3—C6—N4	123.2 (2)	C8—N1—C1	122.99 (19)
C4—C6—N4	129.9 (2)	C5—N1—C1	113.39 (19)
N3—C7—H7A	109.5	C3—N2—N3	103.51 (18)
N3—C7—H7B	109.5	C6—N3—N2	111.98 (19)
H7A—C7—H7B	109.5	C6—N3—C7	128.4 (2)
N3—C7—H7C	109.5	N2—N3—C7	119.56 (19)
H7A—C7—H7C	109.5	C6—N4—H4A	120.0
H7B—C7—H7C	109.5	C6—N4—H4B	120.0
O2—C8—O1	124.3 (2)	H4A—N4—H4B	120.0
O2—C8—N1	124.0 (2)	C8—O1—C9	121.18 (18)
O1—C8—N1	111.59 (19)		
N1—C1—C2—C3	-42.3 (3)	C4—C5—N1—C8	104.8 (2)
C1—C2—C3—N2	-171.7 (2)	C4—C5—N1—C1	-50.8 (3)
C1—C2—C3—C4	9.9 (3)	C2—C1—N1—C8	-86.9 (3)
N2—C3—C4—C6	-0.3 (3)	C2—C1—N1—C5	67.4 (2)
C2—C3—C4—C6	178.3 (2)	C4—C3—N2—N3	0.0 (3)
N2—C3—C4—C5	-175.9 (2)	C2—C3—N2—N3	-178.6 (2)
C2—C3—C4—C5	2.8 (4)	C4—C6—N3—N2	-0.7 (3)

C6—C4—C5—N1	-157.6 (2)	N4—C6—N3—N2	-177.5 (2)
C3—C4—C5—N1	16.7 (3)	C4—C6—N3—C7	-178.1 (2)
C3—C4—C6—N3	0.6 (3)	N4—C6—N3—C7	5.1 (4)
C5—C4—C6—N3	175.7 (2)	C3—N2—N3—C6	0.4 (2)
C3—C4—C6—N4	177.1 (2)	C3—N2—N3—C7	178.1 (2)
C5—C4—C6—N4	-7.8 (4)	O2—C8—O1—C9	-2.9 (3)
O2—C8—N1—C5	9.7 (3)	N1—C8—O1—C9	178.62 (19)
O1—C8—N1—C5	-171.83 (18)	C11—C9—O1—C8	60.3 (3)
O2—C8—N1—C1	162.8 (2)	C12—C9—O1—C8	177.9 (2)
O1—C8—N1—C1	-18.7 (3)	C10—C9—O1—C8	-65.3 (3)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N4—H4 <i>A</i> $\cdots$ O2 <sup>i</sup>	0.86	2.32	3.093 (3)	149
N4—H4 <i>B</i> $\cdots$ N2 <sup>ii</sup>	0.86	2.57	3.420 (3)	172

Symmetry codes: (i)  $-x, -y+1, -z+1$ ; (ii)  $x-1, y, z$ .