

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 4-Ethyl-1-(4-methylbenzylidene)thiosemicarbazide

 Yu-Feng Li<sup>a</sup> and Fang-Fang Jian<sup>b\*</sup>

<sup>a</sup>Microscale Science Institute, Department of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China, and <sup>b</sup>Microscale Science Institute, Weifang University, Weifang 261061, People's Republic of China  
Correspondence e-mail: liyufeng8111@163.com


Received 13 May 2010; accepted 15 May 2010

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.045;  $wR$  factor = 0.144; data-to-parameter ratio = 20.0.

In the title compound,  $\text{C}_{11}\text{H}_{15}\text{N}_3\text{S}$ , an intramolecular  $\text{N}-\text{H}\cdots\text{N}$  hydrogen bond generates an  $S(5)$  ring. In the crystal, inversion dimers linked by pairs of  $\text{N}-\text{H}\cdots\text{S}$  bonds occur, generating an  $R_2^2(8)$  loop.

### Related literature

For a related structure, see: Li &amp; Jian (2010).



### Experimental

#### Crystal data

$\text{C}_{11}\text{H}_{15}\text{N}_3\text{S}$   
 $M_r = 221.32$

Monoclinic,  $P2_1/c$   
 $a = 8.5777$  (17) Å

$b = 13.620$  (3) Å  
 $c = 10.364$  (2) Å  
 $\beta = 90.00$  (3)°  
 $V = 1210.8$  (4) Å<sup>3</sup>  
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 0.24$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.23 \times 0.20 \times 0.18$  mm

#### Data collection

Bruker SMART CCD diffractometer  
11225 measured reflections

2724 independent reflections  
1811 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.035$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.144$   
 $S = 1.10$   
2724 reflections

136 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.22$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.21$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H2A}\cdots\text{S1}^i$	0.86	2.69	3.505 (2)	158
$\text{N3}-\text{H3A}\cdots\text{N1}$	0.86	2.21	2.605 (2)	108

 Symmetry code: (i)  $-x, -y + 1, -z$ .

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINTE* (Bruker, 1997); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5448).

### References

- Bruker (1997). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Li, Y.-F. & Jian, F.-F. (2010). *Acta Cryst.* **E66**, o1397.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supporting information

*Acta Cryst.* (2010). E66, o1399 [https://doi.org/10.1107/S1600536810017988]

## 4-Ethyl-1-(4-methylbenzylidene)thiosemicarbazide

Yu-Feng Li and Fang-Fang Jian

## S1. Experimental

A mixture of 4-methylbenzaldehyde (0.10 mol), and 4-ethylthiosemicarbazide (0.10 mol) was stirred in refluxing ethanol (10 ml) for 4 h to afford the title compound (0.078 mol, yield 78%). Colourless blocks of (I) were obtained by recrystallization from ethanol at room temperature.

## S2. Refinement

H atoms were fixed geometrically and allowed to ride on their attached atoms, with C—H distances = 0.93–0.97 Å; N—H = 0.86 Å and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$  or  $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ .




Figure 1

The structure of (I) showing 30% probability displacement ellipsoids.

## 4-Ethyl-1-(4-methylbenzylidene)thiosemicarbazide

*Crystal data*

$\text{C}_{11}\text{H}_{15}\text{N}_3\text{S}$

$M_r = 221.32$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2_1/c$

$a = 8.5777$  (17) Å

$b = 13.620$  (3) Å

$c = 10.364$  (2) Å

$\beta = 90.00$  (3)°

$V = 1210.8$  (4) Å<sup>3</sup>

$Z = 4$

$F(000) = 472$

$D_x = 1.214$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1811 reflections

$\theta = 3.7$ – $25.4$ °

$\mu = 0.24$  mm<sup>-1</sup>

$T = 293$  K  
Block, colorless

$0.23 \times 0.20 \times 0.18$  mm

#### Data collection

Bruker SMART CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\phi$  and  $\omega$  scans  
11225 measured reflections  
2724 independent reflections

1811 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.035$   
 $\theta_{\text{max}} = 27.5^\circ$ ,  $\theta_{\text{min}} = 3.4^\circ$   
 $h = -10 \rightarrow 11$   
 $k = -17 \rightarrow 17$   
 $l = -13 \rightarrow 12$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.144$   
 $S = 1.10$   
2724 reflections  
136 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0722P)^2 + 0.0957P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 0.22 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.21 \text{ e } \text{\AA}^{-3}$

#### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.18659 (7)	0.61350 (4)	0.05705 (5)	0.0646 (2)
N2	0.02730 (19)	0.59199 (13)	-0.15745 (14)	0.0517 (4)
H2A	-0.0314	0.5527	-0.1144	0.062*
N1	-0.00273 (18)	0.61045 (12)	-0.28521 (14)	0.0471 (4)
C5	-0.1597 (2)	0.57835 (14)	-0.47004 (17)	0.0454 (4)
C6	-0.0800 (2)	0.64307 (15)	-0.55008 (18)	0.0514 (5)
H6A	0.0041	0.6785	-0.5179	0.062*
C8	-0.2494 (2)	0.60447 (16)	-0.72820 (18)	0.0535 (5)
C4	-0.1162 (2)	0.56386 (15)	-0.33565 (18)	0.0494 (5)
H4A	-0.1724	0.5196	-0.2855	0.059*
C3	0.1500 (2)	0.63616 (15)	-0.09996 (17)	0.0486 (5)
N3	0.2320 (2)	0.69703 (14)	-0.17353 (16)	0.0586 (5)
H3A	0.1966	0.7095	-0.2494	0.070*
C7	-0.1245 (2)	0.65502 (17)	-0.67644 (18)	0.0556 (5)
H7A	-0.0691	0.6983	-0.7285	0.067*

C10	-0.2838 (2)	0.52628 (17)	-0.52182 (19)	0.0582 (5)
H10A	-0.3384	0.4819	-0.4706	0.070*
C9	-0.3270 (2)	0.53994 (18)	-0.6490 (2)	0.0618 (6)
H9A	-0.4107	0.5045	-0.6819	0.074*
C11	-0.2989 (3)	0.6213 (2)	-0.8660 (2)	0.0747 (7)
H11A	-0.3868	0.5803	-0.8857	0.112*
H11B	-0.2142	0.6054	-0.9228	0.112*
H11C	-0.3272	0.6890	-0.8774	0.112*
C2	0.3779 (3)	0.7445 (2)	-0.1356 (2)	0.0759 (7)
H2B	0.3760	0.7574	-0.0436	0.091*
H2C	0.3869	0.8071	-0.1798	0.091*
C1	0.5171 (3)	0.6822 (2)	-0.1673 (3)	0.0875 (9)
H1B	0.6104	0.7152	-0.1403	0.131*
H1C	0.5210	0.6710	-0.2587	0.131*
H1D	0.5089	0.6204	-0.1232	0.131*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0854 (4)	0.0693 (4)	0.0392 (3)	-0.0124 (3)	-0.0082 (2)	0.0033 (2)
N2	0.0574 (10)	0.0600 (11)	0.0377 (8)	-0.0058 (8)	-0.0014 (7)	0.0049 (7)
N1	0.0559 (9)	0.0484 (9)	0.0372 (8)	0.0019 (7)	-0.0015 (7)	0.0003 (6)
C5	0.0449 (10)	0.0467 (11)	0.0445 (9)	0.0024 (8)	-0.0011 (7)	-0.0003 (7)
C6	0.0552 (11)	0.0497 (11)	0.0493 (10)	-0.0063 (9)	-0.0042 (8)	0.0026 (8)
C8	0.0596 (12)	0.0557 (12)	0.0451 (10)	0.0138 (10)	-0.0061 (9)	-0.0064 (8)
C4	0.0509 (11)	0.0510 (12)	0.0463 (10)	-0.0008 (9)	0.0016 (8)	0.0047 (8)
C3	0.0561 (11)	0.0490 (11)	0.0406 (9)	0.0019 (9)	-0.0004 (8)	-0.0009 (8)
N3	0.0645 (10)	0.0658 (11)	0.0455 (8)	-0.0144 (9)	-0.0061 (7)	0.0078 (7)
C7	0.0654 (12)	0.0525 (12)	0.0488 (10)	0.0008 (10)	0.0009 (9)	0.0059 (8)
C10	0.0551 (11)	0.0631 (14)	0.0563 (11)	-0.0121 (10)	-0.0008 (9)	0.0036 (9)
C9	0.0572 (12)	0.0670 (15)	0.0611 (12)	-0.0059 (10)	-0.0120 (10)	-0.0078 (10)
C11	0.0885 (16)	0.0861 (19)	0.0494 (12)	0.0190 (14)	-0.0140 (11)	-0.0061 (11)
C2	0.0977 (18)	0.0733 (17)	0.0569 (13)	-0.0359 (15)	-0.0120 (12)	0.0062 (11)
C1	0.0677 (16)	0.095 (2)	0.099 (2)	-0.0246 (15)	-0.0174 (14)	0.0220 (15)

*Geometric parameters (Å, °)*

S1—C3	1.6858 (18)	N3—C2	1.462 (3)
N2—C3	1.351 (2)	N3—H3A	0.8600
N2—N1	1.372 (2)	C7—H7A	0.9300
N2—H2A	0.8600	C10—C9	1.381 (3)
N1—C4	1.274 (2)	C10—H10A	0.9300
C5—C10	1.388 (3)	C9—H9A	0.9300
C5—C6	1.390 (3)	C11—H11A	0.9600
C5—C4	1.455 (2)	C11—H11B	0.9600
C6—C7	1.374 (3)	C11—H11C	0.9600
C6—H6A	0.9300	C2—C1	1.502 (4)
C8—C9	1.375 (3)	C2—H2B	0.9700

C8—C7	1.381 (3)	C2—H2C	0.9700
C8—C11	1.508 (3)	C1—H1B	0.9600
C4—H4A	0.9300	C1—H1C	0.9600
C3—N3	1.328 (3)	C1—H1D	0.9600
C3—N2—N1	119.36 (16)	C9—C10—C5	120.4 (2)
C3—N2—H2A	120.3	C9—C10—H10A	119.8
N1—N2—H2A	120.3	C5—C10—H10A	119.8
C4—N1—N2	116.63 (16)	C8—C9—C10	121.75 (19)
C10—C5—C6	118.08 (17)	C8—C9—H9A	119.1
C10—C5—C4	119.85 (18)	C10—C9—H9A	119.1
C6—C5—C4	122.08 (17)	C8—C11—H11A	109.5
C7—C6—C5	120.48 (18)	C8—C11—H11B	109.5
C7—C6—H6A	119.8	H11A—C11—H11B	109.5
C5—C6—H6A	119.8	C8—C11—H11C	109.5
C9—C8—C7	117.52 (18)	H11A—C11—H11C	109.5
C9—C8—C11	121.8 (2)	H11B—C11—H11C	109.5
C7—C8—C11	120.7 (2)	N3—C2—C1	111.8 (2)
N1—C4—C5	121.40 (18)	N3—C2—H2B	109.3
N1—C4—H4A	119.3	C1—C2—H2B	109.3
C5—C4—H4A	119.3	N3—C2—H2C	109.3
N3—C3—N2	115.95 (16)	C1—C2—H2C	109.3
N3—C3—S1	124.73 (15)	H2B—C2—H2C	107.9
N2—C3—S1	119.30 (15)	C2—C1—H1B	109.5
C3—N3—C2	125.12 (17)	C2—C1—H1C	109.5
C3—N3—H3A	117.4	H1B—C1—H1C	109.5
C2—N3—H3A	117.4	C2—C1—H1D	109.5
C6—C7—C8	121.8 (2)	H1B—C1—H1D	109.5
C6—C7—H7A	119.1	H1C—C1—H1D	109.5
C8—C7—H7A	119.1		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2A...S1 <sup>i</sup>	0.86	2.69	3.505 (2)	158
N3—H3A...N1	0.86	2.21	2.605 (2)	108

Symmetry code: (i)  $-x, -y+1, -z$ .