

## Spiro[1,3-dioxolane-2,3'-indolin]-2'-one

Yan Meng<sup>a\*</sup> and Yanqing Miao<sup>b</sup>

<sup>a</sup>School of Environmental Engineering, Chang'an University, South Second Cycle Road 368#, Xi'an 710054, Shannxi, People's Republic of China, and <sup>b</sup>Department of Pharmacy, Xi'an Medical University, Hanguang Round No. 137, Xi'an 710021, Xi'an, People's Republic of China  
Correspondence e-mail: cg1014@126.com


Received 28 March 2010; accepted 2 May 2010

Key indicators: single-crystal X-ray study;  $T = 273\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.048;  $wR$  factor = 0.137; data-to-parameter ratio = 11.7.

The title compound,  $\text{C}_{10}\text{H}_9\text{NO}_3$ , was synthesized by the condensation reaction of isatin (systematic name 1*H*-indole-2,3-dione) with glycol in presence of *p*-toluenesulfonic acid. The indol-2-one ring system is essentially planar [ $\text{N}-\text{C}=\text{C}-\text{C}$  torsion angle =  $3.1(2)^\circ$ ], and the 1,3-dioxolane ring is slightly distorted. The crystal structure exhibits intermolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds.

### Related literature

For the synthesis of the title compound, see: Santos *et al.* (2008). For the bioactivity of the title compound, see: Demosthenes *et al.* (1998); Rajopadhye & Popp (1988).



### Experimental

#### Crystal data

$\text{C}_{10}\text{H}_9\text{NO}_3$   
 $M_r = 191.18$

Monoclinic,  $P_{\bar{2}}1/c$   
 $a = 7.484(2)\text{ \AA}$

$b = 5.650(1)\text{ \AA}$   
 $c = 20.942(5)\text{ \AA}$   
 $\beta = 97.889(8)^\circ$   
 $V = 877.1(4)\text{ \AA}^3$   
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 0.11\text{ mm}^{-1}$   
 $T = 273\text{ K}$   
 $0.36 \times 0.27 \times 0.21\text{ mm}$

#### Data collection

Bruker SMART CCD  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Bruker, 2002)  
 $T_{\min} = 0.963$ ,  $T_{\max} = 0.989$

4056 measured reflections  
1534 independent reflections  
1093 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.070$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.137$   
 $S = 1.09$   
1534 reflections  
131 parameters

H atoms treated by a mixture of  
independent and constrained  
refinement  
 $\Delta\rho_{\max} = 0.23\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.20\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 $\cdots$ O1 <sup>i</sup>	0.87 (3)	2.07 (3)	2.941 (3)	174 (2)

Symmetry code: (i)  $-x + 1, -y, -z + 1$ .

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINT-Plus* (Bruker, 2002); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2143).

### References

- Bruker (2002). *SMART*, *SAINT-Plus* and *SADABS*. Bruker AXS Inc, Madison, Wisconsin, USA.
- Demosthenes, F., William, J. R., David, S. C. & David, L. C. (1998). *Tetrahedron Lett.* **39**, 2235–2238.
- Rajopadhye, M. & Popp, F. D. (1988). *J. Med. Chem.* **31**, 1001–1005.
- Santos, E. L., Gomes, W. A. Jr, Ribeiro, N. M. & Andrade, H. M. C. (2008). *J. Mol. Catal. A*, **295**, 18–23.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

# supporting information

*Acta Cryst.* (2010). E66, o1305 [https://doi.org/10.1107/S1600536810016132]

## Spiro[1,3-dioxolane-2,3'-indolin]-2'-one

Yan Meng and Yanqing Miao

### S1. Comment

Isatin derivatives, has caught great attention of many researchers as a versatile lead molecule for designing of potential drugs for the variety of biological activities, such as anti-bacteria, anti-virus, anti-tumor and neuroprotection. Among these compounds, spiro-oxindol analogues have received considerable attention as potential anti-bacteria and neuroprotection agents(Demosthenes *et al.* 1998; Rajopadhye *et al.* 1988; Santos *et al.* 2008).


The X-ray structural analysis confirmed the assignment of its structure from spectroscopic data. The molecular structure is depicted in Fig. 1, and a diagram of interactions between the title compounds is depicted in Fig. 2. Geometric parameters of the title compound are in the usual ranges. The crystal packing (Fig. 2) is stabilized by intermolecular N—H···O hydrogen bonds between the indoline H atom and the oxygen of the C=O unit, with a N1—H1···O1<sup>i</sup> (Table 1).

### S2. Experimental


Isatin (1 mmol) and glycol (1 mmol) was dissolved in cyclohexane (20 ml), and 0.01 mmol TsOH was added. The mixture was stirred under reflux. After completion of the reaction, it was evaporated to dryness, followed by chromatography to the pure title compound. <sup>1</sup>H-NMR (D<sub>6</sub>-Acetone, 400 MHz) delta: 10.44 (1H, s), 7.33 (2H, m), 7.00 (1H, td, *J* = 7.2, 0.8 Hz), 6.82 (1H, d, *J* = 7.6 Hz), 4.33 (2H, m), 4.23 (2H, m); EI-MS, m/z (%): 233 (M<sup>+</sup>)

### S3. Refinement

The H atom bound N atom was located from difference Fourier map and refined freely. All H atoms of C atoms were positioned geometrically and refined using a riding model, with C—H = 0.93 Å for aryl and 0.97 Å for methylene H atoms. *U*<sub>iso</sub>(H)= 1.2*U*<sub>eq</sub>(C) for all H atoms.

**Figure 1**

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are presented as a small cycles of arbitrary radius.

**Figure 2**

N–H···O interactions (dotted lines) in the crystal structure of the title compound.

### Spiro[1,3-dioxolane-2,3'-indolin]-2'-one

#### Crystal data

$C_{10}H_9NO_3$   
 $M_r = 191.18$   
Monoclinic,  $P2_1/c$   
Hall symbol: -P 2ybc  
 $a = 7.484 (2)$  Å  
 $b = 5.650 (1)$  Å  
 $c = 20.942 (5)$  Å  
 $\beta = 97.889 (8)^\circ$   
 $V = 877.1 (4)$  Å<sup>3</sup>  
 $Z = 4$

$F(000) = 400$   
 $D_x = 1.448 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 7304 reflections  
 $\theta = 1.5\text{--}25.0^\circ$   
 $\mu = 0.11 \text{ mm}^{-1}$   
 $T = 273 \text{ K}$   
Block, colourless  
 $0.36 \times 0.27 \times 0.21$  mm

*Data collection*

Bruker SMART CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
phi and  $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Bruker, 2002)  
 $T_{\min} = 0.963$ ,  $T_{\max} = 0.989$

4056 measured reflections  
1534 independent reflections  
1093 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.070$   
 $\theta_{\max} = 25.0^\circ$ ,  $\theta_{\min} = 2.0^\circ$   
 $h = -8 \rightarrow 8$   
 $k = -6 \rightarrow 6$   
 $l = -21 \rightarrow 24$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.137$   
 $S = 1.09$   
1534 reflections  
131 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: difference Fourier map  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0648P)^2 + 0.151P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.6983 (2)	0.2241 (3)	0.50491 (8)	0.0600 (5)
O3	0.6788 (2)	0.6884 (3)	0.43906 (8)	0.0551 (5)
O2	0.8104 (2)	0.4151 (3)	0.38133 (8)	0.0560 (5)
N1	0.4575 (3)	0.1613 (4)	0.42609 (9)	0.0469 (6)
H1	0.407 (4)	0.044 (5)	0.4439 (13)	0.070 (9)*
C7	0.3813 (3)	0.2722 (4)	0.36864 (10)	0.0400 (6)
C2	0.6494 (3)	0.4723 (4)	0.40675 (10)	0.0417 (6)
C8	0.4869 (3)	0.4639 (4)	0.35588 (10)	0.0402 (6)
C3	0.4343 (3)	0.6053 (4)	0.30311 (11)	0.0506 (6)
H3A	0.5032	0.7355	0.2945	0.061*
C6	0.2254 (3)	0.2139 (4)	0.32859 (11)	0.0512 (6)
H6A	0.1566	0.0834	0.3370	0.061*
C1	0.6076 (3)	0.2731 (4)	0.45352 (10)	0.0447 (6)
C5	0.1752 (3)	0.3571 (5)	0.27544 (11)	0.0550 (7)
H5A	0.0707	0.3216	0.2477	0.066*
C4	0.2758 (3)	0.5498 (5)	0.26285 (11)	0.0542 (7)

H4A	0.2381	0.6443	0.2272	0.065*
C10	0.8530 (4)	0.7641 (6)	0.4348 (2)	0.0911 (11)
H10A	0.9164	0.7972	0.4774	0.109*
H10B	0.8504	0.9075	0.4092	0.109*
C9	0.9435 (4)	0.5760 (6)	0.40444 (16)	0.0809 (10)
H9B	1.0019	0.6383	0.3694	0.097*
H9C	1.0340	0.5011	0.4355	0.097*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0568 (12)	0.0592 (12)	0.0595 (10)	-0.0091 (9)	-0.0079 (9)	0.0079 (8)
O3	0.0577 (11)	0.0369 (9)	0.0735 (11)	-0.0087 (8)	0.0196 (9)	-0.0187 (8)
O2	0.0441 (10)	0.0489 (10)	0.0777 (11)	-0.0054 (8)	0.0176 (8)	-0.0210 (8)
N1	0.0488 (13)	0.0362 (11)	0.0535 (11)	-0.0084 (10)	-0.0008 (9)	0.0106 (9)
C7	0.0435 (14)	0.0317 (12)	0.0452 (12)	0.0029 (10)	0.0081 (10)	0.0008 (9)
C2	0.0424 (13)	0.0305 (12)	0.0538 (12)	-0.0037 (10)	0.0127 (10)	-0.0069 (10)
C8	0.0435 (13)	0.0297 (12)	0.0490 (12)	0.0015 (10)	0.0122 (10)	-0.0006 (9)
C3	0.0542 (15)	0.0397 (13)	0.0605 (14)	0.0000 (11)	0.0174 (12)	0.0095 (11)
C6	0.0495 (16)	0.0433 (14)	0.0596 (14)	-0.0058 (11)	0.0032 (11)	0.0019 (11)
C1	0.0464 (14)	0.0368 (13)	0.0497 (12)	0.0005 (11)	0.0022 (11)	-0.0019 (10)
C5	0.0483 (15)	0.0612 (17)	0.0540 (13)	0.0034 (13)	0.0014 (11)	0.0042 (12)
C4	0.0558 (16)	0.0569 (16)	0.0505 (13)	0.0139 (14)	0.0095 (11)	0.0114 (11)
C10	0.059 (2)	0.0579 (19)	0.160 (3)	-0.0178 (16)	0.028 (2)	-0.043 (2)
C9	0.0602 (18)	0.084 (2)	0.102 (2)	-0.0233 (17)	0.0217 (16)	-0.0377 (19)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

O1—C1	1.223 (2)	C3—C4	1.393 (3)
O3—C10	1.387 (4)	C3—H3A	0.9300
O3—C2	1.399 (3)	C6—C5	1.386 (3)
O2—C9	1.386 (3)	C6—H6A	0.9300
O2—C2	1.420 (3)	C5—C4	1.370 (3)
N1—C1	1.347 (3)	C5—H5A	0.9300
N1—C7	1.406 (3)	C4—H4A	0.9300
N1—H1	0.87 (3)	C10—C9	1.452 (4)
C7—C6	1.380 (3)	C10—H10A	0.9700
C7—C8	1.388 (3)	C10—H10B	0.9700
C2—C8	1.503 (3)	C9—H9B	0.9700
C2—C1	1.552 (3)	C9—H9C	0.9700
C8—C3	1.377 (3)		
C10—O3—C2	108.95 (19)	C5—C6—H6A	121.2
C9—O2—C2	109.03 (19)	O1—C1—N1	126.8 (2)
C1—N1—C7	111.8 (2)	O1—C1—C2	125.8 (2)
C1—N1—H1	124.0 (17)	N1—C1—C2	107.44 (18)
C7—N1—H1	123.8 (17)	C4—C5—C6	121.5 (2)
C6—C7—C8	121.7 (2)	C4—C5—H5A	119.2

C6—C7—N1	128.6 (2)	C6—C5—H5A	119.2
C8—C7—N1	109.75 (19)	C5—C4—C3	120.5 (2)
O3—C2—O2	107.13 (17)	C5—C4—H4A	119.8
O3—C2—C8	115.39 (18)	C3—C4—H4A	119.8
O2—C2—C8	111.88 (17)	O3—C10—C9	107.6 (2)
O3—C2—C1	111.09 (17)	O3—C10—H10A	110.2
O2—C2—C1	109.05 (18)	C9—C10—H10A	110.2
C8—C2—C1	102.17 (17)	O3—C10—H10B	110.2
C3—C8—C7	120.0 (2)	C9—C10—H10B	110.2
C3—C8—C2	131.7 (2)	H10A—C10—H10B	108.5
C7—C8—C2	108.34 (17)	O2—C9—C10	106.1 (2)
C8—C3—C4	118.7 (2)	O2—C9—H9B	110.5
C8—C3—H3A	120.6	C10—C9—H9B	110.5
C4—C3—H3A	120.6	O2—C9—H9C	110.5
C7—C6—C5	117.6 (2)	C10—C9—H9C	110.5
C7—C6—H6A	121.2	H9B—C9—H9C	108.7
C1—N1—C7—C6	-177.7 (2)	C7—C8—C3—C4	-0.9 (3)
C1—N1—C7—C8	1.5 (3)	C2—C8—C3—C4	178.4 (2)
C10—O3—C2—O2	-0.6 (3)	C8—C7—C6—C5	-1.3 (4)
C10—O3—C2—C8	-125.9 (3)	N1—C7—C6—C5	177.8 (2)
C10—O3—C2—C1	118.4 (3)	C7—N1—C1—O1	176.7 (2)
C9—O2—C2—O3	7.5 (3)	C7—N1—C1—C2	-5.3 (3)
C9—O2—C2—C8	134.9 (2)	O3—C2—C1—O1	-51.7 (3)
C9—O2—C2—C1	-112.8 (2)	O2—C2—C1—O1	66.2 (3)
C6—C7—C8—C3	1.8 (3)	C8—C2—C1—O1	-175.3 (2)
N1—C7—C8—C3	-177.5 (2)	O3—C2—C1—N1	130.3 (2)
C6—C7—C8—C2	-177.7 (2)	O2—C2—C1—N1	-111.9 (2)
N1—C7—C8—C2	3.1 (2)	C8—C2—C1—N1	6.7 (2)
O3—C2—C8—C3	54.2 (3)	C7—C6—C5—C4	-0.1 (4)
O2—C2—C8—C3	-68.6 (3)	C6—C5—C4—C3	0.9 (4)
C1—C2—C8—C3	174.9 (2)	C8—C3—C4—C5	-0.4 (4)
O3—C2—C8—C7	-126.4 (2)	C2—O3—C10—C9	-6.2 (4)
O2—C2—C8—C7	110.7 (2)	C2—O2—C9—C10	-11.1 (4)
C1—C2—C8—C7	-5.8 (2)	O3—C10—C9—O2	10.7 (4)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···O1 <sup>i</sup>	0.87 (3)	2.07 (3)	2.941 (3)	174 (2)

Symmetry code: (i)  $-x+1, -y, -z+1$ .