

## Methyl 1-benzyl-1*H*-1,2,3-triazole-4-carboxylate

Chiung-Cheng Huang,<sup>a</sup> Feng-Ling Wu,<sup>b</sup> Yih Hsing Lo,<sup>b\*</sup>  
Wen-Rong Lai<sup>b</sup> and Chia-Her Lin<sup>c</sup>

<sup>a</sup>Department of Chemical Engineering, Tatung University, Taipei 104, Taiwan,

<sup>b</sup>Department of Natural Science, Taipei Municipal University of Education, Taipei 10048, Taiwan, and <sup>c</sup>Department of Chemistry, Chung-Yuan Christian University, Chung-Li 320, Taiwan

Correspondence e-mail: yhlo@mail.tmue.edu.tw


Received 8 June 2010; accepted 12 June 2010

Key indicators: single-crystal X-ray study;  $T = 200\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.034;  $wR$  factor = 0.091; data-to-parameter ratio = 12.7.

In the title compound,  $\text{C}_{11}\text{H}_{11}\text{N}_3\text{O}_2$ , prepared by the [3+2] cycloaddition reaction of benzyl azide with methyl propiolate, the dihedral angle between the ring planes is  $67.87(11)^\circ$ .

### Related literature

For catalytic transformations of organic alkynes mediated by ruthenium complexes, see: Naota *et al.* (1998); Bruneau & Dixneuf (1999); Trost *et al.* (2001); Chen *et al.* (2009); Cheng *et al.* (2009). For the synthesis of triazoles, see: Padwa (1976). For applications of triazoles, see: Krivopalov & Shkurko (2005).



### Experimental

#### Crystal data

$\text{C}_{11}\text{H}_{11}\text{N}_3\text{O}_2$

$M_r = 217.23$

Monoclinic,  $P2_1/c$

$a = 12.0551(6)\text{ \AA}$

$b = 5.6285(3)\text{ \AA}$

$c = 16.7578(10)\text{ \AA}$

$\beta = 110.664(3)^\circ$

$V = 1063.90(10)\text{ \AA}^3$

$Z = 4$

Mo  $K\alpha$  radiation

$\mu = 0.10\text{ mm}^{-1}$   
 $T = 200\text{ K}$

$0.55 \times 0.40 \times 0.35\text{ mm}$

#### Data collection

Nonius KappaCCD diffractometer  
Absorption correction: multi-scan  
(*SORTAV*; Blessing, 1995)  
 $T_{\min} = 0.949$ ,  $T_{\max} = 0.967$

7021 measured reflections  
1845 independent reflections  
1615 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.023$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.091$   
 $S = 1.01$   
1845 reflections

145 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.15\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.16\text{ e \AA}^{-3}$

Data collection: *COLLECT* (Nonius, 1999); cell refinement: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We gratefully acknowledge financial support from the National Science Council, Taiwan (NSC 97-2113-M-133-001-MY2). We also thank Mr Ting Shen Kuo (Department of Chemistry, National Taiwan Normal University) for his assistance with the X-ray structure analysis and the project of the specific research fields in the Chung Yuan Christian University, Taiwan, under grant CYCU-98-CR—CH.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2294).

### References

- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Bruneau, C. & Dixneuf, P. H. (1999). *Acc. Chem. Res.* **32**, 311–323.
- Chen, C.-K., Tong, H.-C., Chen Hsu, C.-Y., Lee, C.-Y., Fong, Y. H., Chuang, Y.-S., Lo, Y.-H., Lin, Y.-C. & Wang, Y. (2009). *Organometallics*, **28**, 3358–3368.
- Cheng, C.-J., Tong, H.-C., Fong, Y.-H., Wang, P.-Y., Kuo, Y.-L., Lo, Y.-H. & Lin, C.-H. (2009). *Dalton Trans.* pp. 4435–4438.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Krivopalov, V. P. & Shkurko, O. P. (2005). *Russ. Chem. Rev.* **74**, 339–379.
- Naota, T., Takaya, H. & Murahashi, S.-I. (1998). *Chem. Rev.* **98**, 2599–2660.
- Nonius (1999). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Padwa, A. (1976). *Angew. Chem. Int. Ed. Engl.* **15**, 123–136.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Trost, B. M., Toste, F. D. & Pinkerton, A. B. (2001). *Chem. Rev.* **101**, 2067–2096.

# supporting information

*Acta Cryst.* (2010). E66, o1690 [doi:10.1107/S1600536810022531]

## Methyl 1-benzyl-1*H*-1,2,3-triazole-4-carboxylate

**Chiung-Cheng Huang, Feng-Ling Wu, Yih Hsing Lo, Wen-Rong Lai and Chia-Her Lin**

### S1. Comment

Catalytic transformations of organic alkynes mediated by ruthenium complexes are well known, and confirmation for the intermediacy of ruthenium(II) acetylide and vinylidene complexes has been provided (Bruneau & Dixneuf, 1999; Cheng *et al.*, 2009; Naota *et al.*, 1998; Trost *et al.*, 2001). Therefore, ruthenium was a logical choice in our search for a new catalyst of click reaction (Chen *et al.*, 2009). Organic azides are synthetically useful reagents. Amongst many reactions, perhaps the most significant are the 1,3-dipolar cycloaddition reactions with alkynes to synthesize triazoles (Padwa, 1976). Triazoles are nitrogen heteroarenes which have found a range of important applications in the pharmaceutical and agricultural industries (Krivopalov & Shkurko, 2005).

A mixture of benzyl azide and methyl propiolate (1:1.5 equiv, respectively) in toluene was refluxed for 24 h in the presence of 5% moles of  $\{(Tp)(PPh_3)_2Ru(N_3)\}$ , leading to the title compound [Tp is hydridotris(pyrazolyl)borate]. Single crystals of the title compound suitable for X-ray structure analysis were obtained by recrystallization of the crude product from dichloromethane–ether.


In the title compound (Fig. 1), phenyl and triazole are linked together through a methylene group. Of major interest is the methylene C atom, which presents a C—CH<sub>2</sub>—N angle of 112.13 (11) $^{\circ}$ , larger than the ideal tetrahedral value of 109.47 $^{\circ}$ . The N3—C4, C4—C3, C3—N1, N1—N2, and N2—N3 bond lengths are 1.3367 (17), 1.3722 (17), 1.3621 (16), 1.3092 (15) and 1.3560 (15) Å, respectively, which compare with those found for C=C, N=N and C—N bonds in related compounds.

### S2. Experimental

A mixture of benzyl azide, methyl propiolate and  $\{(Tp)(PPh_3)_2Ru(N_3)\}$  in toluene was refluxed for 24 h. The solvent was removed under vacuum and the product was purified by silica gel chromatography. The unreacted alkyne and traces of side products were first eluted out with ether. The pure 1,4-disubstituted triazole product was then obtained by elution with CH<sub>2</sub>Cl<sub>2</sub>.

### S3. Refinement

H atoms were placed in idealized positions and constrained to ride on their parent atoms, with C—H = 0.95–0.99 Å and  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  or  $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C})$ .

**Figure 1**

Molecular structure of the title compound with labelling and displacement ellipsoids drawn at the 30% probability level (H atoms are shown as spheres of arbitrary radius).

### Methyl 1-benzyl-1*H*-1,2,3-triazole-4-carboxylate

#### Crystal data

$C_{11}H_{11}N_3O_2$   
 $M_r = 217.23$   
Monoclinic,  $P2_1/c$   
Hall symbol: -P 2ybc  
 $a = 12.0551 (6)$  Å  
 $b = 5.6285 (3)$  Å  
 $c = 16.7578 (10)$  Å  
 $\beta = 110.664 (3)^\circ$   
 $V = 1063.90 (10)$  Å<sup>3</sup>  
 $Z = 4$

$F(000) = 456$   
 $D_x = 1.356 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 3680 reflections  
 $\theta = 2.6\text{--}25.0^\circ$   
 $\mu = 0.10 \text{ mm}^{-1}$   
 $T = 200$  K  
Prism, colorless  
 $0.55 \times 0.40 \times 0.35$  mm

#### Data collection

Nonius KappaCCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
Detector resolution: 9 pixels mm<sup>-1</sup>  
CCD rotation images, thick slices scans  
Absorption correction: multi-scan  
(SORTAV; Blessing, 1995)  
 $T_{\min} = 0.949$ ,  $T_{\max} = 0.967$

7021 measured reflections  
1845 independent reflections  
1615 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.023$   
 $\theta_{\max} = 25.0^\circ$ ,  $\theta_{\min} = 1.8^\circ$   
 $h = -14 \rightarrow 14$   
 $k = -6 \rightarrow 6$   
 $l = -19 \rightarrow 18$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.091$   
 $S = 1.01$   
1845 reflections  
145 parameters  
0 restraints  
0 constraints  
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
Hydrogen site location: inferred from neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0442P)^2 + 0.2782P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.11251 (12)	1.1370 (3)	0.66802 (10)	0.0462 (4)
H1A	-0.1102	1.3069	0.6814	0.069*
H1B	-0.1054	1.0444	0.7191	0.069*
H1C	-0.1878	1.0991	0.6226	0.069*
C2	-0.00887 (11)	0.8505 (2)	0.61959 (8)	0.0359 (3)
C3	0.09145 (10)	0.8061 (2)	0.59130 (8)	0.0344 (3)
C4	0.13667 (11)	0.5915 (2)	0.57855 (8)	0.0379 (3)
H4	0.1102	0.4373	0.5864	0.045*
C5	0.30857 (12)	0.4920 (2)	0.53001 (10)	0.0441 (3)
H5A	0.3067	0.5334	0.4721	0.053*
H5B	0.2820	0.3252	0.5288	0.053*
C6	0.43422 (11)	0.5135 (2)	0.59233 (8)	0.0359 (3)
C7	0.48209 (14)	0.3404 (3)	0.65326 (10)	0.0508 (4)
H7	0.4357	0.2066	0.6564	0.061*
C8	0.59814 (16)	0.3616 (4)	0.71015 (11)	0.0662 (5)
H8	0.6306	0.2417	0.7518	0.079*
C9	0.66570 (14)	0.5532 (4)	0.70663 (11)	0.0647 (5)
H9	0.7448	0.5669	0.7456	0.078*
C10	0.61865 (14)	0.7251 (3)	0.64659 (12)	0.0630 (5)
H10	0.6653	0.8590	0.6442	0.076*
C11	0.50372 (12)	0.7062 (3)	0.58917 (10)	0.0488 (4)
H11	0.4724	0.8262	0.5474	0.059*
N1	0.15518 (9)	0.98335 (18)	0.57253 (7)	0.0379 (3)
N2	0.23722 (9)	0.88701 (19)	0.54855 (8)	0.0404 (3)
N3	0.22627 (9)	0.64798 (18)	0.55266 (7)	0.0380 (3)
O1	-0.01523 (8)	1.07905 (16)	0.63991 (6)	0.0417 (3)
O2	-0.07666 (8)	0.69845 (17)	0.62483 (7)	0.0480 (3)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0442 (8)	0.0438 (8)	0.0558 (9)	-0.0033 (6)	0.0239 (7)	-0.0050 (6)
C2	0.0330 (6)	0.0314 (6)	0.0372 (7)	-0.0041 (5)	0.0047 (5)	0.0008 (5)
C3	0.0298 (6)	0.0293 (6)	0.0373 (7)	-0.0038 (5)	0.0032 (5)	-0.0005 (5)
C4	0.0326 (6)	0.0298 (6)	0.0456 (8)	-0.0047 (5)	0.0067 (6)	-0.0009 (5)
C5	0.0396 (7)	0.0351 (7)	0.0554 (8)	0.0013 (6)	0.0143 (6)	-0.0090 (6)
C6	0.0361 (7)	0.0332 (7)	0.0416 (7)	0.0016 (5)	0.0177 (6)	-0.0016 (5)
C7	0.0564 (9)	0.0444 (8)	0.0557 (9)	0.0048 (7)	0.0248 (7)	0.0087 (7)
C8	0.0688 (11)	0.0781 (13)	0.0459 (9)	0.0277 (10)	0.0131 (8)	0.0094 (8)
C9	0.0395 (8)	0.0891 (14)	0.0590 (10)	0.0112 (9)	0.0091 (7)	-0.0207 (10)
C10	0.0409 (8)	0.0662 (11)	0.0854 (13)	-0.0141 (8)	0.0266 (8)	-0.0157 (9)
C11	0.0439 (8)	0.0435 (8)	0.0618 (9)	-0.0029 (6)	0.0221 (7)	0.0054 (7)
N1	0.0323 (5)	0.0309 (6)	0.0472 (7)	-0.0012 (4)	0.0098 (5)	-0.0001 (5)
N2	0.0358 (6)	0.0300 (6)	0.0533 (7)	-0.0010 (4)	0.0134 (5)	-0.0008 (5)
N3	0.0329 (5)	0.0283 (5)	0.0474 (7)	-0.0015 (4)	0.0076 (5)	-0.0037 (5)

O1	0.0387 (5)	0.0336 (5)	0.0555 (6)	-0.0053 (4)	0.0199 (4)	-0.0067 (4)
O2	0.0437 (5)	0.0365 (5)	0.0641 (7)	-0.0083 (4)	0.0196 (5)	0.0017 (4)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

C1—O1	1.4469 (16)	C5—H5B	0.9900
C1—H1A	0.9800	C6—C7	1.3817 (19)
C1—H1B	0.9800	C6—C11	1.3830 (19)
C1—H1C	0.9800	C7—C8	1.392 (2)
C2—O2	1.2076 (15)	C7—H7	0.9500
C2—O1	1.3400 (15)	C8—C9	1.365 (3)
C2—C3	1.4678 (19)	C8—H8	0.9500
C3—N1	1.3621 (16)	C9—C10	1.367 (3)
C3—C4	1.3722 (17)	C9—H9	0.9500
C4—N3	1.3367 (17)	C10—C11	1.384 (2)
C4—H4	0.9500	C10—H10	0.9500
C5—N3	1.4713 (17)	C11—H11	0.9500
C5—C6	1.5110 (18)	N1—N2	1.3092 (15)
C5—H5A	0.9900	N2—N3	1.3560 (15)
O1—C1—H1A	109.5	C11—C6—C5	120.62 (12)
O1—C1—H1B	109.5	C6—C7—C8	120.16 (15)
H1A—C1—H1B	109.5	C6—C7—H7	119.9
O1—C1—H1C	109.5	C8—C7—H7	119.9
H1A—C1—H1C	109.5	C9—C8—C7	120.57 (16)
H1B—C1—H1C	109.5	C9—C8—H8	119.7
O2—C2—O1	124.12 (13)	C7—C8—H8	119.7
O2—C2—C3	123.97 (12)	C8—C9—C10	119.48 (15)
O1—C2—C3	111.90 (10)	C8—C9—H9	120.3
N1—C3—C4	108.74 (11)	C10—C9—H9	120.3
N1—C3—C2	123.10 (11)	C9—C10—C11	120.74 (16)
C4—C3—C2	128.16 (11)	C9—C10—H10	119.6
N3—C4—C3	104.61 (11)	C11—C10—H10	119.6
N3—C4—H4	127.7	C6—C11—C10	120.32 (15)
C3—C4—H4	127.7	C6—C11—H11	119.8
N3—C5—C6	112.13 (11)	C10—C11—H11	119.8
N3—C5—H5A	109.2	N2—N1—C3	108.43 (10)
C6—C5—H5A	109.2	N1—N2—N3	107.32 (10)
N3—C5—H5B	109.2	C4—N3—N2	110.89 (11)
C6—C5—H5B	109.2	C4—N3—C5	129.62 (11)
H5A—C5—H5B	107.9	N2—N3—C5	119.49 (11)
C7—C6—C11	118.73 (13)	C2—O1—C1	115.14 (10)
C7—C6—C5	120.65 (12)		