

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## (2E)-1-(2,5-Dichloro-3-thienyl)-3-(6-methoxy-2-naphthyl)prop-2-en-1-one

 Jerry P. Jasinski,<sup>a\*</sup> Albert E. Pek,<sup>a</sup> C. S. Chidan Kumar,<sup>b</sup>  
 H. S. Yathirajan<sup>b</sup> and A. N. Mayekar<sup>b,c</sup>
<sup>a</sup>Department of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, <sup>b</sup>Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and <sup>c</sup>SeQuent Scientific Ltd, Baikampady, New Mangalore, 575 011, India

Correspondence e-mail: jjasinski@keene.edu


Received 10 June 2010; accepted 13 June 2010

 Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.030;  $wR$  factor = 0.077; data-to-parameter ratio = 22.9.

In the title compound,  $\text{C}_{18}\text{H}_{12}\text{Cl}_2\text{O}_2\text{S}$ , the dihedral angle between the thiophene ring and the naphthalene ring system is  $2.13(4)^\circ$ . In the crystal, pairs of weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds form centrosymmetric dimers.

## Related literature

For the biological activity of thiophene-containing compounds, see: Ferreira *et al.* (2006); Bonini *et al.* (2005); Kulikova *et al.* (1980). For the antiradiation activity of thiophenes, see: Hassan *et al.* (1998). For the synthesis and antimicrobial evaluation of new chalcones, see: Tomar *et al.* (2007). For the biological activity of chalcone derivatives, see: Nowakowska *et al.* (2007). For related structures, see: Butcher *et al.* (2007); Harrison *et al.* (2007a,b); Li *et al.* (2009); Yathirajan *et al.* (2006).



## Experimental

## Crystal data

 $\text{C}_{18}\text{H}_{12}\text{Cl}_2\text{O}_2\text{S}$   
 $M_r = 363.24$   
 Monoclinic,  $P2_1/c$   
 $a = 7.3237(5)$  Å  
 $b = 9.4919(6)$  Å  
 $c = 22.4037(15)$  Å  
 $\beta = 96.183(1)^\circ$ 
 $V = 1548.35(18)$  Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.56$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.55 \times 0.40 \times 0.39$  mm

## Data collection

 Bruker APEXII CCD  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2008)  
 $T_{\min} = 0.748$ ,  $T_{\max} = 0.811$ 

 17429 measured reflections  
 4780 independent reflections  
 4373 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.030$   
 $wR(F^2) = 0.077$   
 $S = 0.97$   
 4780 reflections

 209 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.51$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.26$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}2-\text{H}18\cdots\text{O}2^i$	0.93	2.56	3.2051 (14)	127

 Symmetry code: (i)  $-x + 2, -y + 2, -z + 1$ .

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

JPJ thanks Dr Matthias Zeller and the Department of Chemistry, Youngstown State University (YSU), for their assistance with the data collection. The diffractometer was funded by NSF grant No. 0087210, by Ohio Board of Regents grant CAP-491 and by YSU. CSC thanks the University of Mysore for research facilities and HSY thanks the University of Mysore for sabbatical leave.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5100).

## References

- Bonini, C., Chiummiento, L., Bonis, M. D., Funicello, M., Lupattelli, P., Suanno, G., Brault, L., Migianu, E., Naguesque, A., Battaglia, E., Bagrel, D. & Kirsch, G. (2005). *Eur. J. Med. Chem.* **40**, 757–763.
- Bruker (2008). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Butcher, R. J., Yathirajan, H. S., Ashalatha, B. V., Narayana, B. & Sarojini, B. K. (2007). *Acta Cryst.* **E63**, o1430–o1431.
- Ferreira, I. C. F. R., Queiroz, M. R. P., Vilas-Boas, M., Estevinho, L. M., Begouin, A. & Kirsch, G. (2006). *Bioorg. Med. Chem. Lett.* **16**, 1384–1387.
- Harrison, W. T. A., Ashalatha, B. V., Narayana, B., Sarojini, B. K. & Yathirajan, H. S. (2007a). *Acta Cryst.* **E63**, o4183.
- Harrison, W. T. A., Ashalatha, B. V., Narayana, B., Sarojini, B. K. & Yathirajan, H. S. (2007b). *Acta Cryst.* **E63**, o3898.
- Hassan, A. Y., Ghorab, M. M. & Nassar, O. M. (1998). *Phosphorus Sulfur Silicon Relat. Elem.* **134**, 77–86.
- Kulikova, D. A., Churkin, Y. D. & Panfilova, L. V. (1980). *Pharm. Chem. J.* **14**, 227–229.
- Li, H., Mayekar, A. N., Narayana, B., Yathirajan, H. S. & Harrison, W. T. A. (2009). *Acta Cryst.* **E65**, o1533.
- Nowakowska, Z. (2007). *Eur. J. Med. Chem.* **42**, 125–137.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Tomar, V., Bhattacharjee, G., Kamaluddin & Kumar, A. (2007). *Bioorg. Med. Chem. Lett.* **17**, 5321–5324.
- Yathirajan, H. S., Narayana, B., Ashalatha, B., Sarojini, B. K. & Bolte, M. (2006). *Acta Cryst.* **E62**, o4440–o4441.

## supporting information

*Acta Cryst.* (2010). E66, o1717 [doi:10.1107/S1600536810022725]

**(2E)-1-(2,5-Dichloro-3-thienyl)-3-(6-methoxy-2-naphthyl)prop-2-en-1-one**

**Jerry P. Jasinski, Albert E. Pek, C. S. Chidan Kumar, H. S. Yathirajan and A. N Mayekar**

**S1. Comment**

Thiophenes are important heterocyclic compounds that are widely used as building blocks in many agrochemicals and pharmaceuticals. Thiophene containing compounds are well known to exhibit various biological activities such as antioxidant activity (Ferreira *et al.*, 2006), anti-inflammatory agents and anti-HIV PR inhibitors (Bonini *et al.*, 2005). Thiophene derivatives not only being biologically active (Kulikova *et al.*, 1980), but also show antiradiation activity (Hassan *et al.*, 1998).

The synthesis and antimicrobial evaluation of new chalcones containing 2,5-dichlorothiophene moiety is reported (Tomar *et al.*, 2007). Chalcones have been reported to possess many useful properties, including anti-inflammatory, antimicrobial, antifungal, antioxidant, cytotoxic, antitumor and anticancer activities (Nowakowska *et al.*, 2007). In view of the importance of thiophenes, we report here the crystal structure of the title compound.


In the title molecule, the 2,5-dichloro-3-thienyl and 6-methoxy-2-naphthyl rings are bonded at the opposite ends of the propenone group, the biologically active region (Fig.1). The dihedral angle between mean planes of the dichlorothieryl and naphthyl rings is 2.13 (4)°. The angles between the mean plane of the prop-2-en-1-one group and the mean planes of the thienyl and naphthyl rings are 3.08 (4)°, and 2.88 (4)° respectively. In the crystal, pairs of weak intermolecular C2—H18···O2 hydrogen bonds form dimers and contribute to crystal stability.

**S2. Experimental**

1-(2,5-Dichlorothiophen-3-yl)ethanone (1.95 g, 0.01 mol) was mixed with 6-methoxy-2-naphthaldehyde (1.86 g, 0.01 mol) and dissolved in ethanol (30 ml). To this, 3 ml of KOH (50%) was added (Fig. 3). The reaction mixture was stirred for 6 h. The resulting crude solid was filtered, washed successively with distilled water and finally recrystallized from ethanol (95%) to give the pure chalcone. Single crystals suitable for X-ray diffraction studies were grown by the slow evaporation of the acetone-toluene (1:1) solution (m.p. 401–403 K).


**S3. Refinement**

H atoms were placed in their calculated positions and then refined using the riding model with C–H = 0.93 or 0.96 Å, and with  $U_{\text{iso}}(\text{H}) = 1.18\text{--}1.50U_{\text{eq}}(\text{C})$ .




**Figure 1**

Molecular structure of the title compound, showing the atom labeling scheme and 50% probability displacement ellipsoids.

**Figure 2**

Packing diagram of the title compound, viewed down the *a* axis. Dashed lines indicate weak C—H...O intermolecular interactions which form dimers.

**Figure 3**

Reaction scheme for the title compound.

**(2E)-1-(2,5-Dichloro-3-thienyl)-3-(6-methoxy-2-naphthyl)prop-2-en-1-one***Crystal data*C<sub>18</sub>H<sub>12</sub>Cl<sub>2</sub>O<sub>2</sub>S $M_r = 363.24$ Monoclinic,  $P2_1/c$ 

Hall symbol: -P 2ybc

 $a = 7.3237$  (5) Å $b = 9.4919$  (6) Å $c = 22.4037$  (15) Å $\beta = 96.183$  (1)° $V = 1548.35$  (18) Å<sup>3</sup> $Z = 4$  $F(000) = 744$  $D_x = 1.558$  Mg m<sup>-3</sup>Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 9959 reflections

 $\theta = 2.3$ – $31.3$ ° $\mu = 0.56$  mm<sup>-1</sup> $T = 100$  K

Block, yellow

 $0.55 \times 0.40 \times 0.39$  mm*Data collection*

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\omega$  scans

Absorption correction: multi-scan

(SADABS; Bruker, 2008)

 $T_{\min} = 0.748$ ,  $T_{\max} = 0.811$ 

17429 measured reflections

4780 independent reflections

4373 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.021$  $\theta_{\text{max}} = 31.4$ °,  $\theta_{\text{min}} = 1.8$ ° $h = -10$ → $10$  $k = -13$ → $13$  $l = -31$ → $31$ *Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.030$  $wR(F^2) = 0.077$  $S = 0.97$ 

4780 reflections

209 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0374P)^2 + 1.0425P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\text{max}} = 0.001$  $\Delta\rho_{\text{max}} = 0.51$  e Å<sup>-3</sup> $\Delta\rho_{\text{min}} = -0.26$  e Å<sup>-3</sup>*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.97320 (4)	0.57727 (3)	0.666046 (12)	0.02001 (7)
Cl2	1.12452 (4)	1.16879 (3)	0.677155 (14)	0.02261 (7)
S1	1.08482 (4)	0.86262 (3)	0.702059 (12)	0.01784 (7)
O1	0.35474 (12)	-0.11321 (9)	0.30656 (4)	0.01761 (16)

O2	0.81055 (13)	0.85374 (9)	0.49460 (4)	0.02136 (18)
C1	0.93801 (15)	0.81919 (11)	0.59314 (5)	0.01381 (19)
C2	0.97881 (15)	0.96709 (12)	0.59807 (5)	0.01517 (19)
H18	0.9548	1.0307	0.5667	0.018*
C3	1.05603 (15)	1.00339 (12)	0.65347 (5)	0.0166 (2)
C4	0.99005 (15)	0.75105 (12)	0.64645 (5)	0.01516 (19)
C5	0.84960 (15)	0.76639 (12)	0.53422 (5)	0.01448 (19)
C6	0.81083 (16)	0.61554 (12)	0.52385 (5)	0.0159 (2)
H13	0.8391	0.5501	0.5544	0.019*
C7	0.73400 (15)	0.57431 (12)	0.46967 (5)	0.0159 (2)
H12	0.7128	0.6448	0.4409	0.019*
C8	0.67971 (15)	0.43260 (11)	0.45033 (5)	0.01457 (19)
C9	0.70770 (15)	0.31296 (12)	0.48855 (5)	0.01504 (19)
H9	0.7650	0.3241	0.5274	0.018*
C10	0.65144 (15)	0.18169 (12)	0.46900 (5)	0.01494 (19)
H8	0.6709	0.1050	0.4947	0.018*
C11	0.56360 (14)	0.16097 (11)	0.40986 (5)	0.01316 (19)
C12	0.53666 (14)	0.27935 (11)	0.37112 (5)	0.01365 (19)
C13	0.59605 (15)	0.41327 (11)	0.39266 (5)	0.0152 (2)
H11	0.5783	0.4907	0.3673	0.018*
C14	0.50474 (15)	0.02566 (11)	0.38918 (5)	0.01431 (19)
H2	0.5229	-0.0522	0.4143	0.017*
C15	0.42044 (15)	0.01046 (12)	0.33156 (5)	0.01419 (19)
C16	0.39546 (15)	0.12772 (12)	0.29236 (5)	0.0157 (2)
H6	0.3402	0.1154	0.2533	0.019*
C17	0.45213 (15)	0.25871 (12)	0.31159 (5)	0.0155 (2)
H5	0.4353	0.3350	0.2855	0.019*
C18	0.37100 (18)	-0.23386 (13)	0.34421 (6)	0.0225 (2)
H1A	0.4978	-0.2487	0.3586	0.034*
H1B	0.3246	-0.3149	0.3218	0.034*
H1C	0.3016	-0.2196	0.3777	0.034*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.02618 (14)	0.01646 (13)	0.01724 (13)	0.00090 (10)	0.00169 (10)	0.00379 (9)
C12	0.02353 (14)	0.01989 (14)	0.02423 (14)	-0.00595 (10)	0.00179 (10)	-0.00808 (10)
S1	0.01786 (13)	0.02130 (14)	0.01387 (12)	-0.00047 (10)	-0.00050 (9)	-0.00146 (10)
O1	0.0221 (4)	0.0145 (4)	0.0154 (4)	-0.0021 (3)	-0.0015 (3)	-0.0017 (3)
O2	0.0305 (5)	0.0157 (4)	0.0167 (4)	-0.0021 (3)	-0.0032 (3)	0.0020 (3)
C1	0.0133 (4)	0.0139 (4)	0.0142 (4)	-0.0005 (3)	0.0019 (3)	-0.0009 (4)
C2	0.0150 (5)	0.0142 (5)	0.0163 (5)	-0.0010 (4)	0.0016 (4)	-0.0012 (4)
C3	0.0154 (5)	0.0164 (5)	0.0181 (5)	-0.0025 (4)	0.0022 (4)	-0.0029 (4)
C4	0.0150 (5)	0.0154 (5)	0.0151 (5)	-0.0006 (4)	0.0016 (4)	-0.0007 (4)
C5	0.0143 (5)	0.0147 (5)	0.0145 (4)	-0.0014 (4)	0.0017 (3)	-0.0010 (4)
C6	0.0189 (5)	0.0129 (5)	0.0156 (5)	-0.0005 (4)	0.0009 (4)	0.0005 (4)
C7	0.0173 (5)	0.0141 (5)	0.0161 (5)	-0.0003 (4)	0.0015 (4)	0.0000 (4)
C8	0.0145 (5)	0.0139 (5)	0.0154 (5)	-0.0005 (4)	0.0020 (4)	-0.0006 (4)

C9	0.0163 (5)	0.0156 (5)	0.0128 (4)	-0.0007 (4)	-0.0004 (4)	-0.0003 (4)
C10	0.0164 (5)	0.0146 (5)	0.0134 (4)	0.0003 (4)	-0.0004 (4)	0.0016 (4)
C11	0.0122 (4)	0.0142 (5)	0.0131 (4)	0.0002 (3)	0.0013 (3)	-0.0002 (3)
C12	0.0127 (4)	0.0148 (5)	0.0135 (4)	0.0008 (4)	0.0012 (3)	0.0000 (4)
C13	0.0170 (5)	0.0135 (5)	0.0148 (5)	0.0001 (4)	0.0010 (4)	0.0014 (4)
C14	0.0154 (5)	0.0138 (5)	0.0134 (4)	0.0000 (4)	0.0003 (3)	0.0004 (4)
C15	0.0131 (4)	0.0147 (5)	0.0148 (5)	-0.0003 (4)	0.0018 (3)	-0.0019 (4)
C16	0.0152 (5)	0.0187 (5)	0.0130 (4)	0.0007 (4)	0.0000 (4)	-0.0006 (4)
C17	0.0166 (5)	0.0170 (5)	0.0126 (4)	0.0007 (4)	0.0004 (4)	0.0016 (4)
C18	0.0285 (6)	0.0152 (5)	0.0223 (6)	-0.0033 (4)	-0.0042 (5)	0.0007 (4)

*Geometric parameters (Å, °)*

C11—C4	1.7148 (12)	C9—C10	1.3694 (15)
C12—C3	1.7148 (12)	C9—H9	0.93
S1—C3	1.7220 (12)	C10—C11	1.4230 (15)
S1—C4	1.7231 (11)	C10—H8	0.93
O1—C15	1.3657 (13)	C11—C14	1.4167 (15)
O1—C18	1.4198 (14)	C11—C12	1.4206 (15)
O2—C5	1.2257 (14)	C12—C13	1.4116 (15)
C1—C4	1.3756 (15)	C12—C17	1.4220 (15)
C1—C2	1.4371 (15)	C13—H11	0.93
C1—C5	1.4930 (15)	C14—C15	1.3772 (15)
C2—C3	1.3519 (15)	C14—H2	0.93
C2—H18	0.93	C15—C16	1.4174 (15)
C5—C6	1.4732 (15)	C16—C17	1.3658 (16)
C6—C7	1.3403 (15)	C16—H6	0.93
C6—H13	0.93	C17—H5	0.93
C7—C8	1.4553 (15)	C18—H1A	0.96
C7—H12	0.93	C18—H1B	0.96
C8—C13	1.3811 (15)	C18—H1C	0.96
C8—C9	1.4238 (15)		
C3—S1—C4	90.43 (5)	C9—C10—H8	119.5
C15—O1—C18	116.50 (9)	C11—C10—H8	119.5
C4—C1—C2	110.89 (10)	C14—C11—C12	120.00 (10)
C4—C1—C5	131.54 (10)	C14—C11—C10	121.40 (10)
C2—C1—C5	117.57 (10)	C12—C11—C10	118.60 (10)
C3—C2—C1	112.49 (10)	C13—C12—C11	119.07 (10)
C3—C2—H18	123.8	C13—C12—C17	122.12 (10)
C1—C2—H18	123.8	C11—C12—C17	118.80 (10)
C2—C3—C12	126.65 (9)	C8—C13—C12	121.93 (10)
C2—C3—S1	113.15 (9)	C8—C13—H11	119.0
C12—C3—S1	120.20 (7)	C12—C13—H11	119.0
C1—C4—C11	130.82 (9)	C15—C14—C11	119.43 (10)
C1—C4—S1	113.03 (8)	C15—C14—H2	120.3
C11—C4—S1	116.14 (6)	C11—C14—H2	120.3
O2—C5—C6	121.16 (10)	O1—C15—C14	125.24 (10)

O2—C5—C1	117.28 (10)	O1—C15—C16	113.86 (9)
C6—C5—C1	121.57 (10)	C14—C15—C16	120.89 (10)
C7—C6—C5	118.79 (10)	C17—C16—C15	120.32 (10)
C7—C6—H13	120.6	C17—C16—H6	119.8
C5—C6—H13	120.6	C15—C16—H6	119.8
C6—C7—C8	127.89 (10)	C16—C17—C12	120.54 (10)
C6—C7—H12	116.1	C16—C17—H5	119.7
C8—C7—H12	116.1	C12—C17—H5	119.7
C13—C8—C9	118.50 (10)	O1—C18—H1A	109.5
C13—C8—C7	118.74 (10)	O1—C18—H1B	109.5
C9—C8—C7	122.76 (10)	H1A—C18—H1B	109.5
C10—C9—C8	120.97 (10)	O1—C18—H1C	109.5
C10—C9—H9	119.5	H1A—C18—H1C	109.5
C8—C9—H9	119.5	H1B—C18—H1C	109.5
C9—C10—C11	120.92 (10)		
C4—C1—C2—C3	-0.59 (14)	C8—C9—C10—C11	0.10 (17)
C5—C1—C2—C3	179.12 (10)	C9—C10—C11—C14	-179.96 (10)
C1—C2—C3—C12	179.91 (8)	C9—C10—C11—C12	0.54 (16)
C1—C2—C3—S1	0.38 (13)	C14—C11—C12—C13	179.89 (10)
C4—S1—C3—C2	-0.07 (9)	C10—C11—C12—C13	-0.61 (15)
C4—S1—C3—C12	-179.63 (8)	C14—C11—C12—C17	-0.70 (16)
C2—C1—C4—C11	179.61 (9)	C10—C11—C12—C17	178.80 (10)
C5—C1—C4—C11	-0.04 (19)	C9—C8—C13—C12	0.61 (17)
C2—C1—C4—S1	0.53 (12)	C7—C8—C13—C12	-178.84 (10)
C5—C1—C4—S1	-179.12 (10)	C11—C12—C13—C8	0.03 (16)
C3—S1—C4—C1	-0.28 (9)	C17—C12—C13—C8	-179.36 (10)
C3—S1—C4—C11	-179.50 (7)	C12—C11—C14—C15	-0.45 (16)
C4—C1—C5—O2	176.66 (12)	C10—C11—C14—C15	-179.94 (10)
C2—C1—C5—O2	-2.97 (15)	C18—O1—C15—C14	1.95 (16)
C4—C1—C5—C6	-3.43 (18)	C18—O1—C15—C16	-178.24 (10)
C2—C1—C5—C6	176.94 (10)	C11—C14—C15—O1	-178.83 (10)
O2—C5—C6—C7	1.01 (17)	C11—C14—C15—C16	1.37 (16)
C1—C5—C6—C7	-178.90 (10)	O1—C15—C16—C17	179.05 (10)
C5—C6—C7—C8	-178.11 (11)	C14—C15—C16—C17	-1.13 (17)
C6—C7—C8—C13	177.68 (12)	C15—C16—C17—C12	-0.06 (17)
C6—C7—C8—C9	-1.74 (19)	C13—C12—C17—C16	-179.65 (10)
C13—C8—C9—C10	-0.68 (17)	C11—C12—C17—C16	0.96 (16)
C7—C8—C9—C10	178.74 (11)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C2—H18...O2 <sup>i</sup>	0.93	2.56	3.2051 (14)	127

Symmetry code: (i)  $-x+2, -y+2, -z+1$ .