

N-(1,5-Dimethyl-3-oxo-2-phenyl-2,3-dihydro-1H-pyrazol-4-yl)acetamide-naphthalene-2,3-diol (1/1)

Abdullah M. Asiri,^a Salman A. Khan,^a Kong Wai Tan^b and Seik Weng Ng^{b*}

^aChemistry Department, Faculty of Science, King Abdul Aziz University, PO Box 80203, Jeddah 21589, Saudi Arabia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 16 June 2010; accepted 23 June 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.052; wR factor = 0.131; data-to-parameter ratio = 16.4.

In the reaction of naphthalene-2,3-diol and 4-aminoantipyrine in the presence of acetic acid, the amine function is acetylated and the resulting acetamide co-crystallizes with the diol in the title compound, $\text{C}_{13}\text{H}_{15}\text{N}_3\text{O}_2\cdot\text{C}_{10}\text{H}_8\text{O}_2$, with 1:1 molar stoichiometry. The two components are linked by two $\text{O}-\text{H}\cdots\text{O}=\text{C}$ hydrogen bonds. One of the hydroxy groups interacts with the pyrazolone carbonyl O atom and the other hydroxy group interacts with the amide O atom of another component, generating a chain motif. Adjacent chains are linked into a layer motif via $\text{N}-\text{H}\cdots\text{O}$ interactions involving only the heterocyclic acetamide component.

Related literature

For the crystal structure of 4-acetamido-2,3-dimethyl-1-phenyl-5-pyrazol-3-one, see: Kuznetsov *et al.* (1999). For co-crystals of naphthalene-2,3-diol, see: Fritchie & Johnston (1975); Herbert & Truter (1980); Kuo *et al.* (1974); Nakamatsu *et al.* (2003); Wang *et al.* (2008); Wells *et al.* (1974).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{15}\text{N}_3\text{O}_2\cdot\text{C}_{10}\text{H}_8\text{O}_2$

$M_r = 405.44$

Monoclinic, $P2_1/c$
 $a = 12.426 (1)\text{ \AA}$
 $b = 14.304 (2)\text{ \AA}$
 $c = 12.959 (1)\text{ \AA}$
 $\beta = 117.845 (1)^\circ$
 $V = 2036.7 (4)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.25 \times 0.25 \times 0.10\text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
19263 measured reflections

4683 independent reflections
3189 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.061$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.131$
 $S = 1.02$
4683 reflections
286 parameters
27 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.31\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.36\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}1-\text{H}1\cdots\text{O}2^i$	0.87 (1)	2.07 (1)	2.924 (2)	169 (2)
$\text{O}3-\text{H}3\cdots\text{O}2$	0.85 (3)	1.81 (3)	2.639 (2)	163 (3)
$\text{O}4-\text{H}4\cdots\text{O}1^{ii}$	0.85 (3)	1.81 (3)	2.646 (2)	168 (3)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x, -y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdul Aziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2213).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fritchie, C. J. & Johnston, R. M. (1975). *Acta Cryst.* **B31**, 454–461.
- Herbert, J. A. & Truter, M. R. (1980). *J. Chem. Soc. Perkin Trans. 2*, pp. 1253–1258.
- Kuo, M. C., Dunn, J. B. R. & Fritchie, C. J. (1974). *Acta Cryst.* **B30**, 1766–1771.
- Kuznetsov, M. L., Bel'skii, V. K., Dement'ev, A. I., Zaitsev, B. E., Lokshin, B. V. & Zhornik, V. G. (1999). *Russ. Chem. Bull.* **48**, 1274–1280.
- Nakamatsu, S., Yoshizawa, K., Toyota, S., Toda, F. & Matijasic, I. (2003). *Org. Biomol. Chem.* **1**, 2231–2234.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, Y.-T., Tang, G.-M. & Wan, W.-Z. (2008). *Acta Cryst.* **E64**, o1754.
- Wells, J. L., Trus, B. L., Johnston, R. M., Marsh, R. E. & Fritchie, C. J. (1974). *Acta Cryst.* **B30**, 1127–1134.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.

supporting information

Acta Cryst. (2010). E66, o1850 [doi:10.1107/S1600536810024438]

N-(1,5-Dimethyl-3-oxo-2-phenyl-2,3-dihydro-1H-pyrazol-4-yl)acetamide-naphthalene-2,3-diol (1/1)

Abdullah M. Asiri, Salman A. Khan, Kong Wai Tan and Seik Weng Ng

S1. Comment

Naphthalene-2,3-diol forms co-crystals with a number of neutral organic compounds (Fritchie & Johnston, 1975; Herbert & Truter, 1980; Kuo *et al.*, 1974; Nakamatsu *et al.*, 2003; Wang *et al.*, 2008; Wells *et al.*, 1974). The attempt to co-crystallize it with the drug 4-aminoantipyrine was successful when the reaction was carried out in the presence of acetic acid, but the acetic acid converted the amino group to an acetamido group instead. In the co-crystal (Scheme I, Fig. 1), the acetamide and the diol are linked by two $O\cdots H\cdots O=C$ hydrogen bonds. One of the hydroxy groups interacts with the pyrazolyl carbonyl O-atom and the other hydroxy group interacts with the amido O-atom of another component to generate a chain motif. Adjacent chains are linked into a layer motif *via* $N\cdots H\cdots O$ interactions that involves the acetamide component only.


S2. Experimental

Naphthalene-2,3-diol (0.35 g, 2.2 mol) and 4-aminoantipyrine (0.45 g, 2.2 mmol) were heated in methanol (15 ml) for 5 h; three drops of acetic acid were added. Crystals separated from the cool solution when it was set aside for a day.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions [$C-H$ 0.95 to 0.98 Å, $U(H) = 1.2$ to $1.5 U_{eq}(C)$] and were included in the refinement in the riding model approximation. The hydroxy and amino H-atoms were located in a difference Fourier map, and were refined with distance restraints of $O-H$ 0.84 ± 0.01 Å and $N-H$ 0.86 ± 0.01 Å. Their temperature factors were freely refined.

The anisotropic temperature factors of C17 to C20 atoms of the naphthalene fused-ring were restrained to be nearly isotropic and with the restraints, the ellipsoids were somewhat elongated.

**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of the $(C_{13}H_{15}N_3O_2)(C_{10}H_8O_2)$ co-crystal at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

*N-(1,5-Dimethyl-3-oxo-2-phenyl-2,3-dihydro-1*H*-pyrazol-4-yl)acetamide-naphthalene-2,3-diol (1/1)*

Crystal data



$M_r = 405.44$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 12.426(1)\text{ \AA}$

$b = 14.304(2)\text{ \AA}$

$c = 12.959(1)\text{ \AA}$

$\beta = 117.845(1)^\circ$

$V = 2036.7(4)\text{ \AA}^3$

$Z = 4$

$F(000) = 856$

$D_x = 1.322\text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\text{ \AA}$

Cell parameters from 2505 reflections

$\theta = 2.3\text{--}27.9^\circ$

$\mu = 0.09\text{ mm}^{-1}$

$T = 100\text{ K}$

Prism, colorless

$0.25 \times 0.25 \times 0.10\text{ mm}$

Data collection

Bruker SMART APEX

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

19263 measured reflections

4683 independent reflections

3189 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.061$

$\theta_{\max} = 27.5^\circ, \theta_{\min} = 1.9^\circ$

$h = -16 \rightarrow 16$

$k = -18 \rightarrow 18$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.131$
 $S = 1.02$
 4683 reflections
 286 parameters
 27 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0544P)^2 + 0.6929P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.31 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.33173 (12)	0.73209 (10)	0.32885 (12)	0.0290 (3)
O2	0.65387 (11)	0.56345 (9)	0.48381 (11)	0.0232 (3)
O3	0.56600 (12)	0.65279 (10)	0.60626 (11)	0.0253 (3)
O4	0.44923 (12)	0.70985 (10)	0.71521 (12)	0.0279 (3)
N1	0.37471 (14)	0.57790 (11)	0.36783 (14)	0.0224 (4)
N2	0.51486 (14)	0.55398 (11)	0.18066 (13)	0.0233 (4)
N3	0.61709 (14)	0.55929 (11)	0.29124 (13)	0.0219 (4)
C1	0.25557 (17)	0.65611 (14)	0.44515 (17)	0.0255 (4)
H1A	0.1879	0.7007	0.4123	0.038*
H1B	0.3105	0.6728	0.5264	0.038*
H1C	0.2237	0.5929	0.4422	0.038*
C2	0.32386 (16)	0.65897 (14)	0.37571 (16)	0.0219 (4)
C3	0.58105 (17)	0.56524 (12)	0.37598 (16)	0.0199 (4)
C4	0.45083 (17)	0.57041 (13)	0.31423 (16)	0.0218 (4)
C5	0.41487 (17)	0.56533 (13)	0.19785 (16)	0.0240 (4)
C6	0.29135 (19)	0.57192 (16)	0.09672 (18)	0.0326 (5)
H6A	0.2302	0.5591	0.1226	0.049*
H6B	0.2835	0.5261	0.0374	0.049*
H6C	0.2788	0.6350	0.0634	0.049*
C7	0.52538 (19)	0.60264 (14)	0.08616 (17)	0.0284 (5)
H7A	0.4540	0.5885	0.0116	0.043*
H7B	0.5992	0.5817	0.0836	0.043*
H7C	0.5300	0.6702	0.1002	0.043*
C8	0.73390 (18)	0.53349 (14)	0.30516 (17)	0.0266 (4)
C9	0.7468 (2)	0.45251 (15)	0.25275 (18)	0.0326 (5)
H9	0.6785	0.4135	0.2091	0.039*
C10	0.8600 (2)	0.4293 (2)	0.2648 (2)	0.0501 (7)
H10	0.8696	0.3745	0.2285	0.060*
C11	0.9594 (2)	0.4857 (2)	0.3296 (3)	0.0681 (10)
H11	1.0371	0.4698	0.3375	0.082*
C12	0.9456 (2)	0.5654 (2)	0.3830 (3)	0.0676 (10)
H12	1.0144	0.6035	0.4284	0.081*
C13	0.8323 (2)	0.59026 (18)	0.3707 (2)	0.0428 (6)

H13	0.8226	0.6452	0.4067	0.051*
C14	0.63471 (17)	0.66192 (13)	0.72372 (15)	0.0202 (4)
C15	0.75639 (18)	0.64515 (14)	0.78429 (17)	0.0266 (4)
H15	0.7987	0.6232	0.7440	0.032*
C16	0.82114 (19)	0.65991 (16)	0.90656 (18)	0.0339 (5)
C17	0.9490 (2)	0.6515 (2)	0.9714 (2)	0.0634 (9)
H17	0.9945	0.6316	0.9335	0.076*
C18	1.0082 (3)	0.6715 (3)	1.0883 (2)	0.0872 (13)
H18	1.0944	0.6674	1.1299	0.105*
C19	0.9427 (2)	0.6980 (3)	1.1467 (2)	0.0719 (10)
H19	0.9845	0.7104	1.2281	0.086*
C20	0.8197 (2)	0.70618 (18)	1.08762 (19)	0.0406 (6)
H20	0.7761	0.7235	1.1285	0.049*
C21	0.75571 (18)	0.68925 (14)	0.96659 (17)	0.0274 (4)
C22	0.62919 (17)	0.70463 (13)	0.90182 (16)	0.0211 (4)
H22	0.5848	0.7235	0.9413	0.025*
C23	0.56923 (16)	0.69285 (12)	0.78342 (16)	0.0198 (4)
H1	0.373 (2)	0.5318 (11)	0.4107 (17)	0.036 (6)*
H3	0.608 (2)	0.6281 (17)	0.577 (2)	0.054 (8)*
H4	0.419 (2)	0.7345 (19)	0.755 (2)	0.067 (9)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0304 (8)	0.0313 (8)	0.0338 (8)	0.0020 (6)	0.0222 (7)	0.0056 (6)
O2	0.0239 (7)	0.0308 (7)	0.0178 (7)	-0.0018 (6)	0.0121 (6)	-0.0039 (5)
O3	0.0269 (7)	0.0340 (8)	0.0172 (7)	0.0026 (6)	0.0121 (6)	-0.0035 (6)
O4	0.0215 (7)	0.0423 (9)	0.0207 (7)	0.0040 (6)	0.0106 (6)	-0.0036 (6)
N1	0.0230 (8)	0.0287 (9)	0.0210 (9)	-0.0039 (7)	0.0147 (7)	0.0004 (7)
N2	0.0266 (9)	0.0303 (9)	0.0165 (8)	-0.0067 (7)	0.0131 (7)	-0.0030 (7)
N3	0.0238 (8)	0.0281 (9)	0.0175 (8)	-0.0064 (7)	0.0127 (7)	-0.0067 (7)
C1	0.0186 (9)	0.0374 (11)	0.0224 (10)	0.0005 (8)	0.0111 (8)	0.0010 (8)
C2	0.0169 (9)	0.0310 (10)	0.0170 (9)	-0.0037 (8)	0.0073 (8)	-0.0025 (8)
C3	0.0263 (10)	0.0194 (9)	0.0196 (10)	-0.0040 (7)	0.0155 (8)	-0.0036 (7)
C4	0.0246 (10)	0.0240 (10)	0.0214 (10)	-0.0042 (8)	0.0147 (8)	-0.0025 (8)
C5	0.0273 (10)	0.0275 (10)	0.0203 (10)	-0.0063 (8)	0.0137 (8)	-0.0024 (8)
C6	0.0285 (11)	0.0463 (13)	0.0221 (11)	-0.0047 (10)	0.0110 (9)	-0.0027 (9)
C7	0.0387 (12)	0.0323 (11)	0.0211 (10)	-0.0084 (9)	0.0198 (10)	-0.0035 (8)
C8	0.0275 (10)	0.0352 (11)	0.0251 (11)	-0.0061 (8)	0.0191 (9)	-0.0079 (8)
C9	0.0346 (12)	0.0400 (12)	0.0291 (12)	-0.0036 (10)	0.0198 (10)	-0.0093 (9)
C10	0.0452 (15)	0.0678 (18)	0.0458 (15)	0.0041 (13)	0.0284 (13)	-0.0223 (13)
C11	0.0308 (14)	0.117 (3)	0.0654 (19)	-0.0044 (15)	0.0300 (14)	-0.0417 (18)
C12	0.0341 (14)	0.110 (3)	0.071 (2)	-0.0277 (15)	0.0349 (14)	-0.0542 (19)
C13	0.0349 (13)	0.0565 (15)	0.0477 (15)	-0.0169 (11)	0.0282 (12)	-0.0290 (12)
C14	0.0257 (10)	0.0207 (9)	0.0159 (9)	-0.0005 (8)	0.0111 (8)	-0.0022 (7)
C15	0.0262 (10)	0.0348 (11)	0.0247 (11)	0.0040 (8)	0.0169 (9)	-0.0030 (8)
C16	0.0252 (11)	0.0538 (14)	0.0230 (11)	0.0065 (10)	0.0115 (9)	-0.0049 (10)
C17	0.0274 (13)	0.125 (3)	0.0358 (14)	0.0178 (14)	0.0129 (11)	-0.0205 (15)

C18	0.0297 (15)	0.180 (4)	0.0390 (16)	0.0292 (19)	0.0051 (13)	-0.025 (2)
C19	0.0341 (14)	0.142 (3)	0.0275 (14)	0.0191 (16)	0.0041 (11)	-0.0231 (16)
C20	0.0315 (12)	0.0672 (17)	0.0221 (11)	0.0076 (11)	0.0117 (10)	-0.0076 (11)
C21	0.0251 (10)	0.0361 (11)	0.0215 (10)	0.0012 (9)	0.0115 (9)	-0.0030 (8)
C22	0.0259 (10)	0.0235 (9)	0.0186 (9)	-0.0005 (8)	0.0145 (8)	-0.0003 (7)
C23	0.0201 (9)	0.0205 (9)	0.0197 (10)	0.0000 (7)	0.0101 (8)	-0.0004 (7)

Geometric parameters (\AA , $^{\circ}$)

O1—C2	1.236 (2)	C8—C9	1.389 (3)
O2—C3	1.261 (2)	C9—C10	1.381 (3)
O3—C14	1.360 (2)	C9—H9	0.9500
O3—H3	0.85 (3)	C10—C11	1.383 (4)
O4—C23	1.354 (2)	C10—H10	0.9500
O4—H4	0.85 (3)	C11—C12	1.385 (4)
N1—C2	1.347 (2)	C11—H11	0.9500
N1—C4	1.415 (2)	C12—C13	1.388 (3)
N1—H1	0.868 (10)	C12—H12	0.9500
N2—C5	1.371 (2)	C13—H13	0.9500
N2—N3	1.404 (2)	C14—C15	1.361 (3)
N2—C7	1.466 (2)	C14—C23	1.430 (2)
N3—C3	1.369 (2)	C15—C16	1.418 (3)
N3—C8	1.425 (2)	C15—H15	0.9500
C1—C2	1.498 (2)	C16—C17	1.413 (3)
C1—H1A	0.9800	C16—C21	1.426 (3)
C1—H1B	0.9800	C17—C18	1.369 (4)
C1—H1C	0.9800	C17—H17	0.9500
C3—C4	1.433 (3)	C18—C19	1.398 (4)
C4—C5	1.360 (3)	C18—H18	0.9500
C5—C6	1.484 (3)	C19—C20	1.358 (3)
C6—H6A	0.9800	C19—H19	0.9500
C6—H6B	0.9800	C20—C21	1.409 (3)
C6—H6C	0.9800	C20—H20	0.9500
C7—H7A	0.9800	C21—C22	1.411 (3)
C7—H7B	0.9800	C22—C23	1.367 (3)
C7—H7C	0.9800	C22—H22	0.9500
C8—C13	1.380 (3)		
C14—O3—H3	109.9 (18)	C10—C9—H9	120.4
C23—O4—H4	110.2 (19)	C8—C9—H9	120.4
C2—N1—C4	123.16 (16)	C9—C10—C11	120.1 (2)
C2—N1—H1	117.1 (15)	C9—C10—H10	119.9
C4—N1—H1	118.4 (15)	C11—C10—H10	119.9
C5—N2—N3	106.52 (14)	C10—C11—C12	120.0 (2)
C5—N2—C7	121.38 (17)	C10—C11—H11	120.0
N3—N2—C7	115.82 (15)	C12—C11—H11	120.0
C3—N3—N2	110.07 (14)	C11—C12—C13	120.6 (2)
C3—N3—C8	127.60 (16)	C11—C12—H12	119.7

N2—N3—C8	119.84 (14)	C13—C12—H12	119.7
C2—C1—H1A	109.5	C8—C13—C12	118.7 (2)
C2—C1—H1B	109.5	C8—C13—H13	120.7
H1A—C1—H1B	109.5	C12—C13—H13	120.7
C2—C1—H1C	109.5	O3—C14—C15	125.21 (16)
H1A—C1—H1C	109.5	O3—C14—C23	114.64 (16)
H1B—C1—H1C	109.5	C15—C14—C23	120.15 (17)
O1—C2—N1	122.93 (17)	C14—C15—C16	121.07 (17)
O1—C2—C1	121.01 (17)	C14—C15—H15	119.5
N1—C2—C1	116.05 (17)	C16—C15—H15	119.5
O2—C3—N3	123.60 (17)	C17—C16—C15	123.06 (19)
O2—C3—C4	131.17 (16)	C17—C16—C21	117.99 (19)
N3—C3—C4	105.21 (15)	C15—C16—C21	118.87 (18)
C5—C4—N1	126.85 (17)	C18—C17—C16	121.0 (2)
C5—C4—C3	108.47 (16)	C18—C17—H17	119.5
N1—C4—C3	124.68 (16)	C16—C17—H17	119.5
C4—C5—N2	109.47 (17)	C17—C18—C19	120.5 (2)
C4—C5—C6	130.11 (18)	C17—C18—H18	119.8
N2—C5—C6	120.40 (17)	C19—C18—H18	119.8
C5—C6—H6A	109.5	C20—C19—C18	120.3 (2)
C5—C6—H6B	109.5	C20—C19—H19	119.8
H6A—C6—H6B	109.5	C18—C19—H19	119.8
C5—C6—H6C	109.5	C19—C20—C21	120.9 (2)
H6A—C6—H6C	109.5	C19—C20—H20	119.5
H6B—C6—H6C	109.5	C21—C20—H20	119.5
N2—C7—H7A	109.5	C20—C21—C22	121.83 (18)
N2—C7—H7B	109.5	C20—C21—C16	119.28 (19)
H7A—C7—H7B	109.5	C22—C21—C16	118.82 (18)
N2—C7—H7C	109.5	C23—C22—C21	121.33 (17)
H7A—C7—H7C	109.5	C23—C22—H22	119.3
H7B—C7—H7C	109.5	C21—C22—H22	119.3
C13—C8—C9	121.35 (19)	O4—C23—C22	124.57 (16)
C13—C8—N3	118.83 (18)	O4—C23—C14	115.71 (16)
C9—C8—N3	119.81 (18)	C22—C23—C14	119.71 (17)
C10—C9—C8	119.3 (2)		
C5—N2—N3—C3	5.4 (2)	C8—C9—C10—C11	0.8 (4)
C7—N2—N3—C3	143.68 (16)	C9—C10—C11—C12	0.3 (5)
C5—N2—N3—C8	168.77 (16)	C10—C11—C12—C13	-1.0 (5)
C7—N2—N3—C8	-52.9 (2)	C9—C8—C13—C12	0.5 (4)
C4—N1—C2—O1	5.8 (3)	N3—C8—C13—C12	-179.3 (2)
C4—N1—C2—C1	-174.38 (16)	C11—C12—C13—C8	0.6 (5)
N2—N3—C3—O2	174.23 (16)	O3—C14—C15—C16	177.70 (19)
C8—N3—C3—O2	12.4 (3)	C23—C14—C15—C16	-1.5 (3)
N2—N3—C3—C4	-4.09 (19)	C14—C15—C16—C17	-174.2 (2)
C8—N3—C3—C4	-165.87 (17)	C14—C15—C16—C21	2.5 (3)
C2—N1—C4—C5	-79.6 (3)	C15—C16—C17—C18	176.3 (3)
C2—N1—C4—C3	101.6 (2)	C21—C16—C17—C18	-0.4 (5)

O2—C3—C4—C5	−176.82 (19)	C16—C17—C18—C19	2.0 (6)
N3—C3—C4—C5	1.3 (2)	C17—C18—C19—C20	−1.5 (6)
O2—C3—C4—N1	2.2 (3)	C18—C19—C20—C21	−0.7 (5)
N3—C3—C4—N1	−179.66 (17)	C19—C20—C21—C22	−174.8 (3)
N1—C4—C5—N2	−176.99 (17)	C19—C20—C21—C16	2.3 (4)
C3—C4—C5—N2	2.0 (2)	C17—C16—C21—C20	−1.7 (3)
N1—C4—C5—C6	4.2 (3)	C15—C16—C21—C20	−178.5 (2)
C3—C4—C5—C6	−176.78 (19)	C17—C16—C21—C22	175.4 (2)
N3—N2—C5—C4	−4.4 (2)	C15—C16—C21—C22	−1.4 (3)
C7—N2—C5—C4	−139.91 (18)	C20—C21—C22—C23	176.4 (2)
N3—N2—C5—C6	174.48 (17)	C16—C21—C22—C23	−0.7 (3)
C7—N2—C5—C6	39.0 (3)	C21—C22—C23—O4	−177.26 (18)
C3—N3—C8—C13	−64.6 (3)	C21—C22—C23—C14	1.7 (3)
N2—N3—C8—C13	135.2 (2)	O3—C14—C23—O4	−0.8 (2)
C3—N3—C8—C9	115.5 (2)	C15—C14—C23—O4	178.42 (17)
N2—N3—C8—C9	−44.7 (3)	O3—C14—C23—C22	−179.90 (16)
C13—C8—C9—C10	−1.2 (3)	C15—C14—C23—C22	−0.6 (3)
N3—C8—C9—C10	178.6 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···O2 ⁱ	0.87 (1)	2.07 (1)	2.924 (2)	169 (2)
O3—H3···O2	0.85 (3)	1.81 (3)	2.639 (2)	163 (3)
O4—H4···O1 ⁱⁱ	0.85 (3)	1.81 (3)	2.646 (2)	168 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x, -y+3/2, z+1/2$.