

N-[4-(Dimethylamino)benzylidene]-3,4-dimethylisoxazol-5-amine

Abdullah M. Asiri,^a Salman A. Khan,^a Kong Wai Tan^b and Seik Weng Ng^{b*}

^aChemistry Department, Faculty of Science, King Abdul Aziz University, PO Box 80203, Jeddah 21589, Saudi Arabia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 15 June 2010; accepted 19 June 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.039; wR factor = 0.113; data-to-parameter ratio = 17.1.

The aromatic rings attached to the azomethine double bond in the title compound, $C_{14}H_{17}N_3O$, are *trans* to each other [$\text{C}=\text{C}\equiv\text{N}-\text{C}$ torsion angle = $179.5(1)^\circ$], and they are approximately coplanar [dihedral angle between the five- and six-membered rings = $13.7(1)^\circ$].

Related literature

For the spectroscopic characterization of a related Schiff base, see: Asiri *et al.* (2010).

**Experimental***Crystal data*

$C_{14}H_{17}N_3O$	$\gamma = 90.873(1)^\circ$
$M_r = 243.31$	$V = 628.86(10)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 6.5772(6)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.1246(9)\text{ \AA}$	$\mu = 0.08\text{ mm}^{-1}$
$c = 10.538(1)\text{ \AA}$	$T = 100\text{ K}$
$\alpha = 92.995(1)^\circ$	$0.35 \times 0.15 \times 0.10\text{ mm}$
$\beta = 95.183(1)^\circ$	

Data collection

Bruker SMART APEX	2866 independent reflections
diffractometer	2401 reflections with $I > 2\sigma(I)$
6092 measured reflections	$R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	168 parameters
$wR(F^2) = 0.113$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.25\text{ e \AA}^{-3}$
2866 reflections	$\Delta\rho_{\text{min}} = -0.24\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdul Aziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2170).

References

- Asiri, A. M., Khan, S. A. & Rasul, M. G. (2010). *Molbank*, **M684**, 3 pp.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, Submitted.

supporting information

Acta Cryst. (2010). E66, o1783 [doi:10.1107/S1600536810023780]

N-[4-(Dimethylamino)benzylidene]-3,4-dimethylisoxazol-5-amine

Abdullah M. Asiri, Salman A. Khan, Kong Wai Tan and Seik Weng Ng

S1. Comment

Although there is a large number of crystal structure studies of Schiff bases derived by condensing an aromatic aldehyde and an aromatic amine, there has not been any structural report on the condensation product involving 5-amino-3,4-dimethylisoxazole, a commercially available chemical. We have recently reported the spectroscopic characterization of the N-ethylcarbazole-3-aldehyde condensation product of this amine (Asiri *et al.*, 2010). The 4-dimethylaminobenzaldehyde condensation product (Scheme I, Fig. 1) features an azomethine double-bond whose aromatic substituents are located in *trans* positions. The rings are coplanar [C=C=N–C torsion angle 179.5 (1)°].

S2. Experimental

5-Amino-3,4-dimethylisoxazole (0.36 g, 3.2 mol) and *N,N*-dimethylaminobenzaldehyde (0.5 g, 3.2 mol) were heated in methanol (15 ml) for 5 h. The solvent was removed and the solid material recrystallized from methanol to give the crystalline Schiff base.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions [C–H 0.95 to 0.98 Å, $U(H)$ 1.2 to 1.5 $U_{eq}(C)$] and were included in the refinement in the riding model approximation.




Figure 1

Thermal ellipsoid plot (Barbour, 2001) of $C_{14}H_{13}N_3O$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

N-[4-(Dimethylamino)benzylidene]-3,4-dimethylisoxazol-5-amine*Crystal data*

C ₁₄ H ₁₇ N ₃ O	Z = 2
M _r = 243.31	F(000) = 260
Triclinic, P1	D _x = 1.285 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 6.5772 (6) Å	Cell parameters from 2610 reflections
b = 9.1246 (9) Å	θ = 2.2–28.3°
c = 10.538 (1) Å	μ = 0.08 mm ⁻¹
α = 92.995 (1)°	T = 100 K
β = 95.183 (1)°	Prism, yellow
γ = 90.873 (1)°	0.35 × 0.15 × 0.10 mm
V = 628.86 (10) Å ³	

Data collection

Bruker SMART APEX	2401 reflections with $I > 2\sigma(I)$
diffractometer	$R_{\text{int}} = 0.023$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 1.9^\circ$
Graphite monochromator	$h = -8 \rightarrow 8$
ω scans	$k = -11 \rightarrow 11$
6092 measured reflections	$l = -13 \rightarrow 13$
2866 independent reflections	

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.039$	$w = 1/[\sigma^2(F_o^2) + (0.0652P)^2 + 0.0819P]$
$wR(F^2) = 0.113$	where $P = (F_o^2 + 2F_c^2)/3$
S = 1.04	$(\Delta/\sigma)_{\text{max}} = 0.001$
2866 reflections	$\Delta\rho_{\text{max}} = 0.25 \text{ e } \text{\AA}^{-3}$
168 parameters	$\Delta\rho_{\text{min}} = -0.24 \text{ e } \text{\AA}^{-3}$
0 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $\text{Fc}^* = k\text{Fc}[1 + 0.001x\text{Fc}^2\lambda^3/\sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.024 (5)
Secondary atom site location: difference Fourier map	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.34307 (12)	0.64426 (8)	0.20359 (7)	0.0212 (2)
N2	0.46514 (14)	0.43648 (10)	0.31374 (9)	0.0188 (2)
N3	0.41200 (15)	0.73832 (10)	0.11248 (9)	0.0229 (2)
N1	0.13966 (15)	0.02064 (10)	0.73244 (9)	0.0214 (2)
C1	-0.06808 (18)	-0.00795 (14)	0.76262 (12)	0.0275 (3)
H1A	-0.1258	0.0831	0.7959	0.041*
H1B	-0.0673	-0.0816	0.8271	0.041*
H1C	-0.1511	-0.0446	0.6852	0.041*
C3	0.17610 (17)	0.11519 (11)	0.64040 (10)	0.0175 (2)
C2	0.29986 (19)	-0.07284 (12)	0.78465 (12)	0.0251 (3)
H2A	0.3270	-0.1499	0.7204	0.038*
H2B	0.2561	-0.1179	0.8604	0.038*

H2C	0.4245	-0.0138	0.8084	0.038*
C4	0.37269 (17)	0.13022 (12)	0.59705 (10)	0.0189 (2)
H4	0.4797	0.0699	0.6293	0.023*
C5	0.41094 (17)	0.23055 (12)	0.50920 (10)	0.0184 (2)
H5	0.5447	0.2391	0.4827	0.022*
C6	0.25736 (16)	0.32079 (12)	0.45762 (10)	0.0176 (2)
C7	0.06149 (17)	0.30343 (12)	0.49799 (11)	0.0194 (2)
H7	-0.0457	0.3622	0.4634	0.023*
C8	0.01988 (17)	0.20356 (12)	0.58643 (11)	0.0201 (2)
H8	-0.1148	0.1941	0.6113	0.024*
C9	0.29510 (17)	0.42739 (12)	0.36576 (10)	0.0185 (2)
H9	0.1903	0.4939	0.3424	0.022*
C10	0.49005 (17)	0.54187 (12)	0.22757 (10)	0.0180 (2)
C11	0.64745 (16)	0.56405 (11)	0.15557 (10)	0.0180 (2)
C12	0.83736 (17)	0.47749 (13)	0.14888 (11)	0.0237 (3)
H12A	0.8486	0.4097	0.2182	0.036*
H12B	0.9564	0.5444	0.1574	0.036*
H12C	0.8318	0.4215	0.0667	0.036*
C13	0.58994 (17)	0.68813 (12)	0.08652 (10)	0.0197 (2)
C14	0.71023 (19)	0.76199 (13)	-0.00676 (11)	0.0245 (3)
H14A	0.7154	0.6972	-0.0835	0.037*
H14B	0.8493	0.7833	0.0320	0.037*
H14C	0.6448	0.8538	-0.0299	0.037*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0190 (4)	0.0238 (4)	0.0218 (4)	-0.0006 (3)	0.0047 (3)	0.0072 (3)
N2	0.0200 (5)	0.0196 (5)	0.0168 (5)	-0.0031 (4)	0.0033 (4)	0.0010 (3)
N3	0.0236 (5)	0.0256 (5)	0.0203 (5)	-0.0034 (4)	0.0033 (4)	0.0088 (4)
N1	0.0200 (5)	0.0231 (5)	0.0224 (5)	-0.0003 (4)	0.0047 (4)	0.0078 (4)
C1	0.0240 (6)	0.0296 (6)	0.0311 (7)	-0.0021 (5)	0.0093 (5)	0.0117 (5)
C3	0.0197 (6)	0.0169 (5)	0.0160 (5)	-0.0017 (4)	0.0026 (4)	0.0006 (4)
C2	0.0285 (6)	0.0216 (6)	0.0267 (6)	0.0039 (5)	0.0053 (5)	0.0092 (5)
C4	0.0176 (5)	0.0198 (5)	0.0193 (5)	0.0015 (4)	0.0022 (4)	0.0015 (4)
C5	0.0157 (5)	0.0208 (5)	0.0190 (5)	-0.0020 (4)	0.0039 (4)	-0.0001 (4)
C6	0.0179 (5)	0.0188 (5)	0.0159 (5)	-0.0020 (4)	0.0018 (4)	0.0001 (4)
C7	0.0165 (5)	0.0224 (5)	0.0197 (5)	0.0010 (4)	0.0015 (4)	0.0034 (4)
C8	0.0160 (5)	0.0241 (6)	0.0207 (6)	-0.0013 (4)	0.0038 (4)	0.0026 (4)
C9	0.0173 (5)	0.0211 (5)	0.0168 (5)	-0.0012 (4)	0.0007 (4)	0.0012 (4)
C10	0.0182 (5)	0.0189 (5)	0.0165 (5)	-0.0020 (4)	0.0000 (4)	0.0003 (4)
C11	0.0184 (5)	0.0206 (5)	0.0146 (5)	-0.0047 (4)	0.0010 (4)	0.0004 (4)
C12	0.0197 (6)	0.0280 (6)	0.0242 (6)	-0.0016 (4)	0.0045 (5)	0.0038 (5)
C13	0.0205 (6)	0.0227 (5)	0.0156 (5)	-0.0052 (4)	0.0003 (4)	0.0006 (4)
C14	0.0267 (6)	0.0273 (6)	0.0201 (6)	-0.0054 (5)	0.0039 (5)	0.0046 (5)

Geometric parameters (\AA , $\text{^{\circ}}$)

O1—C10	1.3709 (13)	C5—C6	1.4025 (15)
O1—N3	1.4201 (11)	C5—H5	0.9500
N2—C9	1.2926 (14)	C6—C7	1.4020 (15)
N2—C10	1.3745 (14)	C6—C9	1.4419 (15)
N3—C13	1.3097 (15)	C7—C8	1.3788 (15)
N1—C3	1.3659 (14)	C7—H7	0.9500
N1—C2	1.4533 (14)	C8—H8	0.9500
N1—C1	1.4534 (14)	C9—H9	0.9500
C1—H1A	0.9800	C10—C11	1.3565 (15)
C1—H1B	0.9800	C11—C13	1.4155 (15)
C1—H1C	0.9800	C11—C12	1.4930 (15)
C3—C8	1.4132 (15)	C12—H12A	0.9800
C3—C4	1.4168 (15)	C12—H12B	0.9800
C2—H2A	0.9800	C12—H12C	0.9800
C2—H2B	0.9800	C13—C14	1.4966 (15)
C2—H2C	0.9800	C14—H14A	0.9800
C4—C5	1.3718 (15)	C14—H14B	0.9800
C4—H4	0.9500	C14—H14C	0.9800
C10—O1—N3	107.86 (8)	C8—C7—C6	121.91 (10)
C9—N2—C10	119.52 (10)	C8—C7—H7	119.0
C13—N3—O1	105.28 (9)	C6—C7—H7	119.0
C3—N1—C2	120.76 (9)	C7—C8—C3	120.48 (10)
C3—N1—C1	120.12 (9)	C7—C8—H8	119.8
C2—N1—C1	118.15 (9)	C3—C8—H8	119.8
N1—C1—H1A	109.5	N2—C9—C6	122.97 (10)
N1—C1—H1B	109.5	N2—C9—H9	118.5
H1A—C1—H1B	109.5	C6—C9—H9	118.5
N1—C1—H1C	109.5	C11—C10—O1	109.95 (10)
H1A—C1—H1C	109.5	C11—C10—N2	129.32 (10)
H1B—C1—H1C	109.5	O1—C10—N2	120.73 (9)
N1—C3—C8	121.22 (10)	C10—C11—C13	104.10 (10)
N1—C3—C4	121.27 (10)	C10—C11—C12	128.36 (10)
C8—C3—C4	117.50 (10)	C13—C11—C12	127.54 (10)
N1—C2—H2A	109.5	C11—C12—H12A	109.5
N1—C2—H2B	109.5	C11—C12—H12B	109.5
H2A—C2—H2B	109.5	H12A—C12—H12B	109.5
N1—C2—H2C	109.5	C11—C12—H12C	109.5
H2A—C2—H2C	109.5	H12A—C12—H12C	109.5
H2B—C2—H2C	109.5	H12B—C12—H12C	109.5
C5—C4—C3	121.04 (10)	N3—C13—C11	112.81 (10)
C5—C4—H4	119.5	N3—C13—C14	120.31 (10)
C3—C4—H4	119.5	C11—C13—C14	126.88 (11)
C4—C5—C6	121.58 (10)	C13—C14—H14A	109.5
C4—C5—H5	119.2	C13—C14—H14B	109.5
C6—C5—H5	119.2	H14A—C14—H14B	109.5

C7—C6—C5	117.44 (10)	C13—C14—H14C	109.5
C7—C6—C9	120.23 (10)	H14A—C14—H14C	109.5
C5—C6—C9	122.33 (10)	H14B—C14—H14C	109.5
C10—O1—N3—C13	0.59 (11)	C7—C6—C9—N2	-171.57 (10)
C2—N1—C3—C8	178.20 (10)	C5—C6—C9—N2	7.97 (17)
C1—N1—C3—C8	9.66 (16)	N3—O1—C10—C11	-0.86 (11)
C2—N1—C3—C4	-2.91 (16)	N3—O1—C10—N2	179.38 (9)
C1—N1—C3—C4	-171.45 (10)	C9—N2—C10—C11	-174.71 (11)
N1—C3—C4—C5	-176.74 (10)	C9—N2—C10—O1	5.00 (15)
C8—C3—C4—C5	2.19 (16)	O1—C10—C11—C13	0.75 (12)
C3—C4—C5—C6	-0.80 (17)	N2—C10—C11—C13	-179.51 (10)
C4—C5—C6—C7	-0.81 (16)	O1—C10—C11—C12	-178.70 (10)
C4—C5—C6—C9	179.64 (9)	N2—C10—C11—C12	1.03 (19)
C5—C6—C7—C8	1.00 (16)	O1—N3—C13—C11	-0.13 (12)
C9—C6—C7—C8	-179.45 (10)	O1—N3—C13—C14	-179.61 (9)
C6—C7—C8—C3	0.44 (17)	C10—C11—C13—N3	-0.38 (13)
N1—C3—C8—C7	176.93 (10)	C12—C11—C13—N3	179.08 (10)
C4—C3—C8—C7	-2.01 (16)	C10—C11—C13—C14	179.06 (10)
C10—N2—C9—C6	-179.54 (9)	C12—C11—C13—C14	-1.47 (18)