

***fac*-Tris(4-aminobenzohydroxamato)-iron(III) ethanol solvate**Ahmed B. M. Alagha,^a Declan Gaynor,^a Helge Müller-Bunz,^b Kevin B. Nolan^{a*} and Laavanya Parthasarathi^c^aCentre for Synthesis and Chemical Biology, Department of Pharmaceutical and Medicinal Chemistry, Royal College of Surgeons in Ireland, Dublin 2, Ireland,^bCentre for Synthesis and Chemical Biology, School of Chemistry and Chemical Biology, University College Dublin, Dublin 4, Ireland, and ^cSmurfit Institute of Genetics, Trinity College Dublin, Dublin 2, Ireland

Correspondence e-mail: kbnolan@rcsi.ie


Received 17 June 2010; accepted 21 June 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; disorder in solvent or counterion; R factor = 0.040; wR factor = 0.094; data-to-parameter ratio = 13.3.

In the structure of the title compound, $[\text{Fe}(\text{C}_7\text{H}_7\text{N}_2\text{O}_2)_3] \cdot \text{CH}_3\text{CH}_2\text{OH}$, the Fe^{III} atom is in a distorted octahedral O_6 environment with the three hydroxamate O atoms (and the three carbonyl O atoms) arranged in a *fac* configuration and one of the hydroxamate ligands being puckered. The methyl C atom of the ethanol solvent molecule is disordered over two positions with occupancies of 0.626 (13) and 0.374 (13), respectively. The cocrystallized ethanol molecule is hydrogen bonded to one of the hydroxamate O atoms. $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ interactions generate infinite three-dimensional networks along [100], [010] and [001].

Related literature

For a detailed account of the molecular and crystal structures of related tris(hydroxamato) Fe^{III} complexes, see: Rio-Echevarria *et al.* (2008); Mulcahy *et al.* (2007); Marmion *et al.* (2000). For ring puckering parameters, see: Cremer & Pople (1975); for pseudorotation parameters, see: Rao *et al.* (1981).

**Experimental***Crystal data*

$[\text{Fe}(\text{C}_7\text{H}_7\text{N}_2\text{O}_2)_3] \cdot \text{C}_2\text{H}_5\text{OH}$	$\gamma = 112.217(3)^\circ$
$M_r = 555.36$	$V = 1305.1(4)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 10.6232(18)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.6455(18)\text{ \AA}$	$\mu = 0.63\text{ mm}^{-1}$
$c = 13.225(2)\text{ \AA}$	$T = 100\text{ K}$
$\alpha = 107.550(3)^\circ$	$0.30 \times 0.20 \times 0.03\text{ mm}$
$\beta = 91.085(4)^\circ$	

Data collection

Bruker SMART CCD area detector diffractometer	24614 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2003)	5739 independent reflections
$T_{\min} = 0.883$, $T_{\max} = 0.981$	5065 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.094$	$\Delta\rho_{\max} = 0.43\text{ e \AA}^{-3}$
$S = 1.06$	$\Delta\rho_{\min} = -0.31\text{ e \AA}^{-3}$
5739 reflections	
433 parameters	

Table 1
Selected bond lengths (\AA).

Fe—O2	1.9615 (14)	Fe—O5	2.0444 (14)
Fe—O6	1.9769 (14)	Fe—O1	2.0470 (14)
Fe—O4	1.9822 (14)	Fe—O3	2.0654 (13)

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O7—H1O7 \cdots O6 ⁱ	0.78 (3)	1.91 (3)	2.694 (2)	175 (3)
N6—H1N6 \cdots O3 ⁱⁱ	0.84 (2)	2.15 (2)	2.968 (2)	163.9 (19)
N5—H1N5 \cdots O7	0.84 (2)	1.91 (2)	2.740 (2)	176 (2)
N3—H1N3 \cdots O4 ⁱⁱⁱ	0.86 (3)	1.98 (3)	2.755 (2)	150 (2)
N1—H1N1 \cdots O2 ^{iv}	0.84 (3)	1.99 (3)	2.737 (2)	149 (2)

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $-x, -y + 1, -z$; (iv) $-x, -y, -z$.

Data collection: *SMART* (Bruker, 2003); cell refinement: *SAINT* (Bruker, 2003); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

We thank the Irish Government under its Programme for Research in Third Level Institutions (Cycle 3), Enterprise Ireland (International Collaboration programme) and EU COST D8 for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5023).

References

- Bruker (2003). *SMART and SAINT*. Bruker AXS Inc, Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Marmion, C. J., Murphy, T., Starikova, Z. & Nolan, K. B. (2000). *Acta Cryst. C* **56**, e491–e492.
- Mulcahy, C., Krot, K. A., Griffith, D. M., Suponitsky, K. Y., Starikova, Z. A. & Marmion, C. J. (2007). *Eur. J. Inorg. Chem.* pp. 1373–1380.
- Rao, S. T., Westhof, E. & Sundaralingam, M. (1981). *Acta Cryst. A* **37**, 421–425.
- Rio-Echevarria, I. M., White, F. J., Brechin, E. K., Tasker, P. A. & Harris, S. G. (2008). *Chem. Commun.* pp. 4570–4572.
- Sheldrick, G. M. (2003). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155. 23

supporting information

Acta Cryst. (2010). E66, m853–m854 [doi:10.1107/S160053681002413X]

***fac*-Tris(4-aminobenzohydroxamato)iron(III) ethanol solvate**

Ahmed B. M. Alagha, Declan Gaynor, Helge Müller-Bunz, Kevin B. Nolan and Laavanya Parthasarathi

S1. Comment


The Fe^{III} core is in a distorted octahedral O6 environment with the three 4-aminophenylhydroxamato ligands coordinated through both O-atoms to form stable five-membered chelates (Fig. 1). The complex has a *fac* geometric configuration defined by the positioning of the three hydroxamate or the three carbonyl O atoms. The Fe—O3—C8—N3—O4 and Fe—O5—C15—N5—O6 five membered rings are not puckered whereas the Fe—O1—C1—N1—O2 ring exhibits a puckered configuration with the closest pucker descriptor being twisted on O2—Fe. The Cremer and Pople puckering parameters (Cremer & Pople, 1975) are $Q_2 = 0.120$ (2) Å & $\varphi_2 = 330.0$ (9) $^\circ$ and the pseudorotation parameters (Rao *et al.*, 1981) are $P = 130.8$ (6) $^\circ$ and $\tau(M) = 11.2$ (1) $^\circ$ for reference bond Fe—O1. The co-crystallized ethanol molecule is hydrogen bonded to one of the hydroxamate oxygen atoms through strong O—H···O bond (O7 H1O7 O6 interaction in Table 2). The O—H···O and N—H···O interactions (Table 2) form a set of clusters generating infinite two dimensional networks along the base vectors [0 1 0] and [1 0 0] and along the plane (0 0 1). These clusters then assemble to form infinite three dimensional networks (Fig. 2) along the base vectors [1 0 0], [0 1 0] and [0 0 1].

S2. Experimental


FeCl₃·6H₂O (59.3 mg, 0.22 mmol) in water (5 ml), was added to a solution of 4-aminophenylhydroxamic acid, (100 mg, 0.66 mmol) in ethanol (15 ml). The pH of the resulting solution was then raised to 5.7 using 0.1 M NaOH solution whereupon a dark red precipitate was obtained. This was removed by filtration and the filtrate was left to stand at room temperature for two months whereupon very dark red crystals were obtained. These were collected by filtration and dried. Yield (100 mg, 0.18 mmol, 82%). Elemental analysis (%), calcd C₂₃H₂₇N₆O₇Fe: C, 49.56; H, 5.26; N, 15.08; Fe, 10.02; found C 49.01, H 5.02, N 15.03, Fe 9.77. IR (KBr Disc); ν_{max} 3345br, 3211br, 1604 s, 1525 s, 1484 s cm⁻¹.

S3. Refinement

All H-atoms (except those attached to the methylene and the methyl carbon atoms of the ethanol solvate) were located in difference maps and their positions and isotropic displacement parameters were freely refined. The CH₂ and CH₃ H atoms of the solvate were constrained to ride on their parent C atoms.

**Figure 1**

The asymmetric unit of (I) with the atoms labelled and displacement ellipsoids depicted at the 50% probability level for all non-H atoms. H-atoms are drawn as spheres of arbitrary radius

**Figure 2**

The molecular packing viewed down the b -axis. Dashed lines represent the O—H···O and N—H···O interactions within the lattice.

fac-Tris(4-aminobenzohydroxamato)iron(III) ethanol solvate

Crystal data

$[\text{Fe}(\text{C}_7\text{H}_7\text{N}_2\text{O}_2)_3] \cdot \text{C}_2\text{H}_6\text{O}$

$M_r = 555.36$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 10.6232 (18)$ Å

$b = 10.6455 (18)$ Å

$c = 13.225 (2)$ Å

$\alpha = 107.550 (3)^\circ$

$\beta = 91.085 (4)^\circ$

$\gamma = 112.217 (3)^\circ$

$V = 1305.1 (4)$ Å³

$Z = 2$

$F(000) = 578$

$D_x = 1.413 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4743 reflections

$\theta = 2.2\text{--}24.9^\circ$

$\mu = 0.63 \text{ mm}^{-1}$

$T = 100$ K

Plate, red

$0.30 \times 0.20 \times 0.03$ mm

Data collection

Bruker SMART CCD area detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 8.366 pixels mm⁻¹

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2003)

$T_{\min} = 0.883$, $T_{\max} = 0.981$

24614 measured reflections

5739 independent reflections

5065 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.044$
 $\theta_{\text{max}} = 27.1^\circ$, $\theta_{\text{min}} = 1.6^\circ$
 $h = -13 \rightarrow 13$

$k = -13 \rightarrow 13$
 $l = -16 \rightarrow 16$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.094$
 $S = 1.06$
5739 reflections
433 parameters
0 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: difference Fourier map
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0438P)^2 + 0.6552P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.004$
 $\Delta\rho_{\text{max}} = 0.43 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.31 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^* / U_{\text{eq}}$	Occ. (<1)
Fe	0.13879 (3)	0.36163 (3)	0.16284 (2)	0.01220 (9)	
O2	0.06647 (14)	0.16143 (14)	0.06621 (11)	0.0180 (3)	
N1	-0.05305 (18)	0.07800 (18)	0.09477 (14)	0.0187 (4)	
H1N1	-0.087 (3)	-0.007 (3)	0.052 (2)	0.035 (7)*	
C1	-0.0862 (2)	0.1334 (2)	0.18804 (16)	0.0161 (4)	
O1	-0.01237 (14)	0.26557 (14)	0.24206 (11)	0.0162 (3)	
C2	-0.2049 (2)	0.0445 (2)	0.22618 (16)	0.0179 (4)	
C3	-0.2669 (2)	0.1117 (2)	0.30398 (18)	0.0239 (5)	
H3	-0.231 (3)	0.211 (3)	0.331 (2)	0.039 (7)*	
C4	-0.3816 (2)	0.0325 (3)	0.3394 (2)	0.0279 (5)	
H4	-0.425 (3)	0.078 (3)	0.391 (2)	0.031 (7)*	
C5	-0.4376 (2)	-0.1172 (2)	0.29942 (18)	0.0243 (5)	
N2	-0.5509 (2)	-0.1970 (3)	0.3375 (2)	0.0340 (5)	
H1N2	-0.595 (3)	-0.277 (3)	0.296 (2)	0.040 (9)*	
H2N2	-0.602 (3)	-0.147 (3)	0.364 (2)	0.048 (9)*	
C6	-0.3739 (3)	-0.1848 (2)	0.2234 (2)	0.0317 (5)	
H6	-0.410 (3)	-0.288 (3)	0.198 (2)	0.043 (8)*	
C7	-0.2604 (2)	-0.1056 (2)	0.1865 (2)	0.0286 (5)	
H7	-0.219 (3)	-0.152 (3)	0.136 (2)	0.037 (7)*	
O4	0.04007 (15)	0.41933 (14)	0.06861 (11)	0.0175 (3)	
N3	0.03983 (18)	0.55267 (18)	0.11997 (14)	0.0188 (4)	
H1N3	0.008 (3)	0.585 (3)	0.078 (2)	0.029 (7)*	

C8	0.1040 (2)	0.6232 (2)	0.21800 (15)	0.0146 (4)	
O3	0.15716 (14)	0.56033 (14)	0.26357 (10)	0.0149 (3)	
C9	0.1150 (2)	0.7704 (2)	0.27167 (16)	0.0154 (4)	
C10	0.1173 (2)	0.8613 (2)	0.21373 (17)	0.0196 (4)	
H10	0.116 (2)	0.832 (2)	0.1403 (18)	0.014 (5)*	
C11	0.1315 (2)	1.0008 (2)	0.26579 (17)	0.0205 (4)	
H11	0.135 (2)	1.060 (2)	0.2244 (17)	0.015 (5)*	
C12	0.1403 (2)	1.0527 (2)	0.37769 (17)	0.0184 (4)	
N4	0.1479 (2)	1.1887 (2)	0.42768 (17)	0.0238 (4)	
H1N4	0.165 (3)	1.217 (3)	0.494 (2)	0.027 (7)*	
H2N4	0.161 (3)	1.247 (3)	0.388 (2)	0.042 (8)*	
C13	0.1409 (2)	0.9625 (2)	0.43591 (17)	0.0182 (4)	
H13	0.148 (2)	0.995 (3)	0.508 (2)	0.024 (6)*	
C14	0.1295 (2)	0.8241 (2)	0.38380 (16)	0.0168 (4)	
H14	0.129 (2)	0.765 (2)	0.4247 (17)	0.010 (5)*	
O6	0.30277 (14)	0.41024 (15)	0.09015 (11)	0.0178 (3)	
N5	0.41597 (18)	0.41538 (19)	0.14684 (14)	0.0173 (4)	
H1N5	0.480 (2)	0.412 (2)	0.1112 (18)	0.016 (6)*	
C15	0.4028 (2)	0.3985 (2)	0.24097 (15)	0.0145 (4)	
O5	0.28464 (14)	0.37299 (14)	0.27232 (11)	0.0157 (3)	
C16	0.5226 (2)	0.4146 (2)	0.30955 (15)	0.0154 (4)	
C17	0.5013 (2)	0.3612 (2)	0.39495 (16)	0.0178 (4)	
H17	0.408 (2)	0.311 (2)	0.4045 (18)	0.021 (6)*	
C18	0.6104 (2)	0.3822 (2)	0.46526 (16)	0.0175 (4)	
H18	0.592 (2)	0.346 (3)	0.5211 (19)	0.024 (6)*	
C19	0.7457 (2)	0.4587 (2)	0.45299 (15)	0.0158 (4)	
N6	0.8551 (2)	0.4808 (2)	0.52404 (15)	0.0198 (4)	
H1N6	0.835 (2)	0.467 (2)	0.5823 (18)	0.008 (5)*	
H2N6	0.927 (3)	0.549 (3)	0.529 (2)	0.037 (8)*	
C20	0.7675 (2)	0.5081 (2)	0.36565 (17)	0.0186 (4)	
H20	0.854 (2)	0.559 (2)	0.3568 (17)	0.015 (5)*	
C21	0.6581 (2)	0.4880 (2)	0.29572 (17)	0.0187 (4)	
H21	0.677 (2)	0.527 (2)	0.2400 (17)	0.012 (5)*	
O7	0.62865 (18)	0.41913 (19)	0.03203 (14)	0.0300 (4)	
H1O7	0.651 (3)	0.473 (3)	0.000 (2)	0.030 (7)*	
C22	0.6632 (3)	0.2992 (3)	-0.0101 (2)	0.0450 (7)	
H22A	0.7633	0.3304	0.0097	0.054*	0.626 (13)
H22B	0.6425	0.2649	-0.0893	0.054*	0.626 (13)
H22C	0.5911	0.2283	-0.0711	0.054*	0.374 (13)
H22D	0.7502	0.3321	-0.0395	0.054*	0.374 (13)
C23A	0.5904 (9)	0.1809 (6)	0.0278 (5)	0.048 (2)	0.626 (13)
H23A	0.5992	0.2168	0.1062	0.073*	0.626 (13)
H23B	0.6297	0.1088	0.0061	0.073*	0.626 (13)
H23C	0.4929	0.1371	-0.0032	0.073*	0.626 (13)
C23B	0.6799 (14)	0.2255 (11)	0.0608 (8)	0.052 (3)	0.374 (13)
H23D	0.7001	0.1432	0.0207	0.077*	0.374 (13)
H23E	0.5950	0.1914	0.0912	0.077*	0.374 (13)
H23F	0.7561	0.2917	0.1189	0.077*	0.374 (13)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe	0.01437 (15)	0.00964 (14)	0.01186 (14)	0.00477 (11)	0.00097 (10)	0.00273 (10)
O2	0.0186 (7)	0.0124 (7)	0.0189 (7)	0.0040 (6)	0.0068 (6)	0.0023 (6)
N1	0.0189 (9)	0.0108 (8)	0.0199 (9)	0.0027 (7)	0.0037 (7)	0.0002 (7)
C1	0.0155 (9)	0.0125 (9)	0.0187 (10)	0.0061 (8)	0.0006 (8)	0.0028 (8)
O1	0.0176 (7)	0.0102 (6)	0.0168 (7)	0.0036 (5)	0.0032 (5)	0.0015 (5)
C2	0.0178 (10)	0.0148 (10)	0.0201 (10)	0.0057 (8)	0.0022 (8)	0.0058 (8)
C3	0.0252 (11)	0.0143 (10)	0.0286 (12)	0.0054 (9)	0.0085 (9)	0.0052 (9)
C4	0.0281 (12)	0.0259 (12)	0.0317 (13)	0.0118 (10)	0.0146 (10)	0.0106 (10)
C5	0.0181 (10)	0.0253 (11)	0.0301 (12)	0.0043 (9)	0.0001 (9)	0.0161 (10)
N2	0.0229 (11)	0.0333 (13)	0.0448 (14)	0.0024 (10)	0.0056 (10)	0.0229 (11)
C6	0.0296 (13)	0.0151 (11)	0.0439 (15)	0.0021 (10)	0.0070 (11)	0.0099 (10)
C7	0.0299 (12)	0.0150 (10)	0.0382 (14)	0.0087 (9)	0.0126 (11)	0.0051 (10)
O4	0.0266 (8)	0.0127 (7)	0.0132 (7)	0.0111 (6)	-0.0023 (6)	0.0004 (5)
N3	0.0263 (9)	0.0166 (9)	0.0158 (9)	0.0130 (7)	-0.0014 (7)	0.0033 (7)
C8	0.0161 (9)	0.0137 (9)	0.0142 (9)	0.0061 (8)	0.0024 (7)	0.0050 (7)
O3	0.0191 (7)	0.0115 (6)	0.0140 (7)	0.0063 (5)	-0.0011 (5)	0.0038 (5)
C9	0.0161 (9)	0.0143 (9)	0.0161 (9)	0.0073 (8)	0.0018 (7)	0.0041 (8)
C10	0.0257 (11)	0.0190 (10)	0.0162 (10)	0.0112 (9)	0.0029 (8)	0.0060 (8)
C11	0.0253 (11)	0.0186 (10)	0.0231 (11)	0.0117 (9)	0.0041 (9)	0.0108 (9)
C12	0.0146 (9)	0.0151 (10)	0.0256 (11)	0.0063 (8)	0.0057 (8)	0.0065 (8)
N4	0.0326 (11)	0.0153 (9)	0.0239 (11)	0.0109 (8)	0.0077 (8)	0.0050 (8)
C13	0.0202 (10)	0.0168 (10)	0.0151 (10)	0.0065 (8)	0.0036 (8)	0.0032 (8)
C14	0.0185 (10)	0.0144 (9)	0.0181 (10)	0.0062 (8)	0.0028 (8)	0.0069 (8)
O6	0.0160 (7)	0.0245 (8)	0.0167 (7)	0.0092 (6)	0.0019 (6)	0.0105 (6)
N5	0.0152 (8)	0.0225 (9)	0.0181 (9)	0.0091 (7)	0.0045 (7)	0.0102 (7)
C15	0.0175 (10)	0.0103 (9)	0.0162 (9)	0.0063 (7)	0.0027 (8)	0.0042 (7)
O5	0.0158 (7)	0.0171 (7)	0.0160 (7)	0.0071 (6)	0.0025 (5)	0.0073 (6)
C16	0.0180 (10)	0.0141 (9)	0.0159 (9)	0.0086 (8)	0.0028 (8)	0.0046 (8)
C17	0.0181 (10)	0.0159 (10)	0.0202 (10)	0.0070 (8)	0.0052 (8)	0.0068 (8)
C18	0.0227 (10)	0.0196 (10)	0.0151 (10)	0.0109 (8)	0.0047 (8)	0.0094 (8)
C19	0.0195 (10)	0.0141 (9)	0.0141 (9)	0.0102 (8)	0.0022 (8)	0.0007 (7)
N6	0.0197 (10)	0.0247 (10)	0.0168 (9)	0.0100 (8)	0.0037 (7)	0.0083 (8)
C20	0.0142 (10)	0.0190 (10)	0.0209 (10)	0.0046 (8)	0.0045 (8)	0.0069 (8)
C21	0.0198 (10)	0.0203 (10)	0.0171 (10)	0.0069 (8)	0.0043 (8)	0.0092 (8)
O7	0.0367 (10)	0.0372 (10)	0.0326 (9)	0.0221 (8)	0.0197 (8)	0.0238 (8)
C22	0.0638 (19)	0.0503 (17)	0.0404 (16)	0.0374 (16)	0.0226 (14)	0.0222 (14)
C23A	0.063 (5)	0.036 (3)	0.058 (4)	0.026 (3)	0.021 (3)	0.023 (3)
C23B	0.048 (7)	0.039 (5)	0.069 (6)	0.022 (5)	-0.016 (5)	0.015 (4)

Geometric parameters (\AA , ^\circ)

Fe—O2	1.9615 (14)	N4—H1N4	0.83 (3)
Fe—O6	1.9769 (14)	N4—H2N4	0.90 (3)
Fe—O4	1.9822 (14)	C13—C14	1.382 (3)
Fe—O5	2.0444 (14)	C13—H13	0.90 (2)

Fe—O1	2.0470 (14)	C14—H14	0.94 (2)
Fe—O3	2.0654 (13)	O6—N5	1.379 (2)
O2—N1	1.383 (2)	N5—C15	1.313 (3)
N1—C1	1.315 (3)	N5—H1N5	0.84 (2)
N1—H1N1	0.84 (3)	C15—O5	1.285 (2)
C1—O1	1.282 (2)	C15—C16	1.474 (3)
C1—C2	1.466 (3)	C16—C17	1.396 (3)
C2—C3	1.394 (3)	C16—C21	1.400 (3)
C2—C7	1.396 (3)	C17—C18	1.377 (3)
C3—C4	1.378 (3)	C17—H17	0.96 (2)
C3—H3	0.93 (3)	C18—C19	1.400 (3)
C4—C5	1.391 (3)	C18—H18	0.92 (2)
C4—H4	0.94 (3)	C19—N6	1.381 (3)
C5—N2	1.390 (3)	C19—C20	1.396 (3)
C5—C6	1.395 (3)	N6—H1N6	0.84 (2)
N2—H1N2	0.80 (3)	N6—H2N6	0.82 (3)
N2—H2N2	0.90 (3)	C20—C21	1.380 (3)
C6—C7	1.380 (3)	C20—H20	0.90 (2)
C6—H6	0.96 (3)	C21—H21	0.94 (2)
C7—H7	0.92 (3)	O7—C22	1.417 (3)
O4—N3	1.378 (2)	O7—H1O7	0.78 (3)
N3—C8	1.304 (3)	C22—C23B	1.438 (8)
N3—H1N3	0.86 (3)	C22—C23A	1.446 (5)
C8—O3	1.283 (2)	C22—H22A	0.9900
C8—C9	1.470 (3)	C22—H22B	0.9900
C9—C10	1.398 (3)	C22—H22C	0.9900
C9—C14	1.401 (3)	C22—H22D	0.9900
C10—C11	1.382 (3)	C23A—H23A	0.9800
C10—H10	0.92 (2)	C23A—H23B	0.9800
C11—C12	1.402 (3)	C23A—H23C	0.9800
C11—H11	0.95 (2)	C23B—H23D	0.9800
C12—N4	1.368 (3)	C23B—H23E	0.9800
C12—C13	1.402 (3)	C23B—H23F	0.9800
O2—Fe—O6	90.72 (6)	C12—C13—H13	119.8 (15)
O2—Fe—O4	92.73 (6)	C13—C14—C9	120.68 (18)
O6—Fe—O4	91.70 (6)	C13—C14—H14	119.0 (13)
O2—Fe—O5	104.13 (6)	C9—C14—H14	120.3 (13)
O6—Fe—O5	78.92 (5)	N5—O6—Fe	112.13 (11)
O4—Fe—O5	160.68 (6)	C15—N5—O6	117.38 (16)
O2—Fe—O1	79.40 (5)	C15—N5—H1N5	127.0 (15)
O6—Fe—O1	162.99 (6)	O6—N5—H1N5	114.4 (15)
O4—Fe—O1	102.49 (6)	O5—C15—N5	118.19 (17)
O5—Fe—O1	90.01 (6)	O5—C15—C16	121.59 (17)
O2—Fe—O3	163.96 (6)	N5—C15—C16	120.18 (17)
O6—Fe—O3	102.91 (6)	C15—O5—Fe	113.19 (12)
O4—Fe—O3	78.63 (5)	C17—C16—C21	118.42 (18)
O5—Fe—O3	86.98 (5)	C17—C16—C15	119.09 (18)

O1—Fe—O3	89.21 (5)	C21—C16—C15	122.41 (18)
N1—O2—Fe	111.18 (11)	C18—C17—C16	121.01 (19)
C1—N1—O2	117.85 (16)	C18—C17—H17	120.6 (14)
C1—N1—H1N1	129.3 (18)	C16—C17—H17	118.4 (14)
O2—N1—H1N1	112.1 (18)	C17—C18—C19	120.58 (19)
O1—C1—N1	117.72 (18)	C17—C18—H18	118.3 (15)
O1—C1—C2	122.48 (17)	C19—C18—H18	121.1 (15)
N1—C1—C2	119.79 (17)	N6—C19—C20	120.91 (19)
C1—O1—Fe	112.64 (12)	N6—C19—C18	120.57 (19)
C3—C2—C7	118.2 (2)	C20—C19—C18	118.49 (18)
C3—C2—C1	119.10 (18)	C19—N6—H1N6	115.4 (14)
C7—C2—C1	122.69 (19)	C19—N6—H2N6	117 (2)
C4—C3—C2	121.1 (2)	H1N6—N6—H2N6	115 (2)
C4—C3—H3	119.6 (17)	C21—C20—C19	120.82 (19)
C2—C3—H3	119.3 (17)	C21—C20—H20	119.1 (14)
C3—C4—C5	120.8 (2)	C19—C20—H20	119.9 (14)
C3—C4—H4	121.3 (16)	C20—C21—C16	120.62 (19)
C5—C4—H4	117.9 (16)	C20—C21—H21	118.1 (13)
N2—C5—C4	120.7 (2)	C16—C21—H21	121.3 (13)
N2—C5—C6	121.0 (2)	C22—O7—H1O7	113 (2)
C4—C5—C6	118.3 (2)	O7—C22—C23B	118.2 (4)
C5—N2—H1N2	114 (2)	O7—C22—C23A	113.5 (3)
C5—N2—H2N2	112.1 (19)	C23B—C22—C23A	37.0 (4)
H1N2—N2—H2N2	114 (3)	O7—C22—H22A	108.9
C7—C6—C5	121.0 (2)	C23B—C22—H22A	73.0
C7—C6—H6	120.4 (17)	C23A—C22—H22A	108.9
C5—C6—H6	118.6 (17)	O7—C22—H22B	108.9
C6—C7—C2	120.6 (2)	C23B—C22—H22B	129.9
C6—C7—H7	120.1 (17)	C23A—C22—H22B	108.9
C2—C7—H7	119.2 (17)	H22A—C22—H22B	107.7
N3—O4—Fe	111.77 (11)	O7—C22—H22C	107.7
C8—N3—O4	118.68 (16)	C23B—C22—H22C	107.9
C8—N3—H1N3	127.3 (17)	C23A—C22—H22C	75.6
O4—N3—H1N3	113.2 (17)	H22A—C22—H22C	136.9
O3—C8—N3	117.74 (17)	H22B—C22—H22C	37.7
O3—C8—C9	122.39 (17)	O7—C22—H22D	107.7
N3—C8—C9	119.87 (17)	C23B—C22—H22D	107.6
C8—O3—Fe	113.04 (12)	C23A—C22—H22D	135.6
C10—C9—C14	118.69 (18)	H22A—C22—H22D	39.5
C10—C9—C8	121.53 (18)	H22B—C22—H22D	71.1
C14—C9—C8	119.73 (17)	H22C—C22—H22D	107.1
C11—C10—C9	120.73 (19)	C22—C23A—H22C	38.6
C11—C10—H10	118.5 (13)	C22—C23A—H23A	109.5
C9—C10—H10	120.7 (13)	H22C—C23A—H23A	142.0
C10—C11—C12	120.61 (19)	C22—C23A—H23B	109.5
C10—C11—H11	118.6 (13)	H22C—C23A—H23B	102.8
C12—C11—H11	120.8 (13)	C22—C23A—H23C	109.5
N4—C12—C13	121.5 (2)	H22C—C23A—H23C	76.8

N4—C12—C11	119.96 (19)	C22—C23B—H23D	109.5
C13—C12—C11	118.58 (18)	C22—C23B—H23E	109.5
C12—N4—H1N4	114.5 (18)	H23D—C23B—H23E	109.5
C12—N4—H2N4	117.8 (18)	C22—C23B—H23F	109.5
H1N4—N4—H2N4	125 (3)	H23D—C23B—H23F	109.5
C14—C13—C12	120.63 (19)	H23E—C23B—H23F	109.5
C14—C13—H13	119.5 (15)		
O6—Fe—O2—N1	175.55 (12)	O3—C8—C9—C10	149.8 (2)
O4—Fe—O2—N1	-92.72 (12)	N3—C8—C9—C10	-29.2 (3)
O5—Fe—O2—N1	96.80 (12)	O3—C8—C9—C14	-27.4 (3)
O1—Fe—O2—N1	9.49 (11)	N3—C8—C9—C14	153.54 (19)
O3—Fe—O2—N1	-36.0 (3)	C14—C9—C10—C11	-1.0 (3)
Fe—O2—N1—C1	-11.7 (2)	C8—C9—C10—C11	-178.25 (19)
O2—N1—C1—O1	6.1 (3)	C9—C10—C11—C12	-1.7 (3)
O2—N1—C1—C2	-174.36 (16)	C10—C11—C12—N4	-177.0 (2)
N1—C1—O1—Fe	2.5 (2)	C10—C11—C12—C13	2.9 (3)
C2—C1—O1—Fe	-176.99 (14)	N4—C12—C13—C14	178.27 (19)
O2—Fe—O1—C1	-6.88 (13)	C11—C12—C13—C14	-1.6 (3)
O6—Fe—O1—C1	-62.3 (2)	C12—C13—C14—C9	-1.0 (3)
O4—Fe—O1—C1	83.59 (13)	C10—C9—C14—C13	2.3 (3)
O5—Fe—O1—C1	-111.26 (13)	C8—C9—C14—C13	179.61 (18)
O3—Fe—O1—C1	161.76 (13)	O2—Fe—O6—N5	-100.53 (12)
O1—C1—C2—C3	20.6 (3)	O4—Fe—O6—N5	166.72 (12)
N1—C1—C2—C3	-158.8 (2)	O5—Fe—O6—N5	3.73 (11)
O1—C1—C2—C7	-160.5 (2)	O1—Fe—O6—N5	-46.5 (3)
N1—C1—C2—C7	20.0 (3)	O3—Fe—O6—N5	88.00 (12)
C7—C2—C3—C4	-1.3 (3)	Fe—O6—N5—C15	-4.7 (2)
C1—C2—C3—C4	177.7 (2)	O6—N5—C15—O5	2.6 (3)
C2—C3—C4—C5	0.9 (4)	O6—N5—C15—C16	-175.01 (16)
C3—C4—C5—N2	178.1 (2)	N5—C15—O5—Fe	0.8 (2)
C3—C4—C5—C6	0.6 (4)	C16—C15—O5—Fe	178.37 (13)
N2—C5—C6—C7	-179.2 (2)	O2—Fe—O5—C15	85.37 (13)
C4—C5—C6—C7	-1.7 (4)	O6—Fe—O5—C15	-2.56 (12)
C5—C6—C7—C2	1.3 (4)	O4—Fe—O5—C15	-64.7 (2)
C3—C2—C7—C6	0.2 (4)	O1—Fe—O5—C15	164.44 (12)
C1—C2—C7—C6	-178.7 (2)	O3—Fe—O5—C15	-106.35 (12)
O2—Fe—O4—N3	166.76 (12)	O5—C15—C16—C17	18.8 (3)
O6—Fe—O4—N3	-102.43 (12)	N5—C15—C16—C17	-163.68 (18)
O5—Fe—O4—N3	-42.2 (2)	O5—C15—C16—C21	-157.92 (18)
O1—Fe—O4—N3	87.02 (12)	N5—C15—C16—C21	19.6 (3)
O3—Fe—O4—N3	0.40 (11)	C21—C16—C17—C18	1.2 (3)
Fe—O4—N3—C8	1.7 (2)	C15—C16—C17—C18	-175.66 (18)
O4—N3—C8—O3	-4.0 (3)	C16—C17—C18—C19	0.4 (3)
O4—N3—C8—C9	175.04 (16)	C17—C18—C19—N6	179.44 (18)
N3—C8—O3—Fe	4.1 (2)	C17—C18—C19—C20	-2.4 (3)
C9—C8—O3—Fe	-174.89 (14)	N6—C19—C20—C21	-178.92 (19)
O2—Fe—O3—C8	-60.9 (3)	C18—C19—C20—C21	3.0 (3)

O6—Fe—O3—C8	86.72 (13)	C19—C20—C21—C16	−1.4 (3)
O4—Fe—O3—C8	−2.41 (12)	C17—C16—C21—C20	−0.6 (3)
O5—Fe—O3—C8	164.62 (13)	C15—C16—C21—C20	176.07 (18)
O1—Fe—O3—C8	−105.33 (13)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O7—H1O7···O6 ⁱ	0.78 (3)	1.91 (3)	2.694 (2)	175 (3)
N6—H1N6···O3 ⁱⁱ	0.84 (2)	2.15 (2)	2.968 (2)	163.9 (19)
N5—H1N5···O7	0.84 (2)	1.91 (2)	2.740 (2)	176 (2)
N3—H1N3···O4 ⁱⁱⁱ	0.86 (3)	1.98 (3)	2.755 (2)	150 (2)
N1—H1N1···O2 ^{iv}	0.84 (3)	1.99 (3)	2.737 (2)	149 (2)

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x+1, -y+1, -z+1$; (iii) $-x, -y+1, -z$; (iv) $-x, -y, -z$.