

**8-Quinolyl 5-(dimethylamino)-naphthalene-1-sulfonate****Zuo-an Xiao\*** and **Dan Zhan**

School of Chemical Engineering and Food Science, Xiangfan University, Xiangfan 441053, People's Republic of China

Correspondence e-mail: blueice8250@yahoo.com.cn


Received 21 July 2010; accepted 27 July 2010

Key indicators: single-crystal X-ray study;  $T = 298\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  $R$  factor = 0.050;  $wR$  factor = 0.132; data-to-parameter ratio = 14.3.

In the title compound,  $\text{C}_{21}\text{H}_{18}\text{N}_2\text{O}_3\text{S}$ , the dihedral angle between the naphthalene and quinoline ring systems is  $55.53(2)^\circ$ , and the torsion angle involving the connecting  $\text{C}-\text{S}-\text{O}-\text{C}$  atoms is  $87.60(3)^\circ$ . In the crystal structure, weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds connect molecules into chains along [100] and there are  $\pi-\pi$  stacking interactions between pairs of chains with a centroid–centroid distance of  $3.5485(15)\text{ \AA}$ .

**Related literature**

For background information and the applications of compounds containing the 5-(dimethylamino)naphthalene-1-sulfonyl group, see: Li *et al.* (1975); Walkup & Imperiali (1997); Chen & Chen (2004).

**Experimental***Crystal data* $\text{C}_{21}\text{H}_{18}\text{N}_2\text{O}_3\text{S}$  $M_r = 378.43$ 

Triclinic, $P\bar{1}$	$V = 912.30(19)\text{ \AA}^3$
$a = 9.5556(12)\text{ \AA}$	$Z = 2$
$b = 10.1237(12)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 11.4182(14)\text{ \AA}$	$\mu = 0.20\text{ mm}^{-1}$
$\alpha = 108.736(2)^\circ$	$T = 298\text{ K}$
$\beta = 100.426(2)^\circ$	$0.20 \times 0.20 \times 0.20\text{ mm}$
$\gamma = 111.860(2)^\circ$	

*Data collection*

Bruker SMART CCD diffractometer	5269 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1997)	3526 independent reflections
$T_{\min} = 0.961$ , $T_{\max} = 0.980$	2959 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.054$

*Refinement*

$R[F^2 > 2\sigma(F^2)] = 0.050$	246 parameters
$wR(F^2) = 0.132$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$
3526 reflections	$\Delta\rho_{\min} = -0.34\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C16}-\text{H16}\cdots\text{O1}^{\text{i}}$	0.93	2.52	3.411 (3)	160

Symmetry code: (i)  $x + 1, y, z$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors are grateful to the Science Technology Research Programme of the Education Office of Hubei Province (grant No. Q20092503) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5093).

**References**

- Bruker (2007). *SAINT-Plus* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, C. & Chen, Q. (2004). *Tetrahedron Lett.* **45**, 3957–3960.
- Li, Y., Chan, L., Tyer, L., Moody, R. T., Hirnel, C. M. & Hercules, D. M. (1975). *J. Am. Chem. Soc.* **97**, 3118–3126.
- Sheldrick, G. M. (1997). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Walkup, G. K. & Imperiali, B. (1997). *J. Am. Chem. Soc.* **119**, 3443–3450.

# supporting information

*Acta Cryst.* (2010). E66, o2180 [https://doi.org/10.1107/S160053681002979X]

## 8-Quinolyl 5-(dimethylamino)naphthalene-1-sulfonate

Zuo-an Xiao and Dan Zhan

### S1. Comment

The dansyl fluorophore (5-(dimethylamino)naphthalene-1-sulfonyl) is characterized by a charge transfer excited state exhibiting solvatochromism and high emission quantum yields (Li *et al.*, 1975). These characteristics, together with the synthetic flexibility of the sulfonic acid group, have led the dansyl fluorophore to be a core-structure present in many fluorescent sensors and labels for the detection of both metal cations and anions (Walkup & Imperiali, 1997; Chen & Chen, 2004). We are interested in designing fluorescent drug or ligand analogs that are expected to bind to hydrophobic sites in proteins or membranes. With this mind, the title compound, (I), was prepared and we report the crystal structure herein.


In the molecular structure (Fig. 1), the dihedral angle between the naphthalene and quinoline ring systems is 55.53 (2) $^{\circ}$ , and these aromatic ring systems are connected by the atoms C8—S1—O3—C13, giving a torsion angle of 87.60 (3) $^{\circ}$ . In the crystal structure (Fig. 2) molecules are linked by weak intermolecular C—H $\cdots$ O hydrogen bonds forming 1-D chains along [100]. Pairs of chains are connected by weak  $\pi$  $\cdots$  $\pi$  stacking interactions with Cg $\cdots$ Cg(2-x, -y, -z) = 3.5485 (15), where Cg is the centroid defined by ring atoms C13-C17/C21.

### S2. Experimental


8-Hydroxyquinolin (0.16 g, 1 mmol) was added to a stirred solution of dansyl chloride (0.27 g, 1 mmol) in dry acetone (40 ml). The reaction mixture was allowed to stir for 12 hr at 293 K. The solvent was evaporated and the residue was purified by column chromatography (petroleum ether-ethyl acetate, 1:4 v/v) to afford the title compound as a yellow solid. Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in ethyl acetate at room temperature.

### S3. Refinement

All H atoms were placed in idealized positions [C—H(methyl)=0.96 Å, and 0.93 Å (aromatic), with  $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}(\text{methyl C})$  1.2 $U_{\text{eq}}(\text{other C})$ ].

**Figure 1**

The molecular structure of (I), with displacement ellipsoids drawn at the 50% probability level.

**Figure 2**

Part of the crystal structure of (I) showing hydrogen bonds as dashed lines.

### 8-Quinolyl 5-(dimethylamino)naphthalene-1-sulfonate

#### Crystal data

C<sub>21</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub>S

M<sub>r</sub> = 378.43

Triclinic, P<sup>1</sup>

Hall symbol: -P 1

a = 9.5556 (12) Å

b = 10.1237 (12) Å

c = 11.4182 (14) Å

α = 108.736 (2)°

β = 100.426 (2)°

γ = 111.860 (2)°

V = 912.30 (19) Å<sup>3</sup>

Z = 2

F(000) = 396

D<sub>x</sub> = 1.378 Mg m<sup>-3</sup>

Mo Kα radiation, λ = 0.71073 Å

Cell parameters from 2502 reflections

θ = 1.7–22.5°

μ = 0.20 mm<sup>-1</sup>

T = 298 K

Block, yellow

0.20 × 0.20 × 0.20 mm

*Data collection*

Bruker SMART CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Sheldrick, 1997)  
 $T_{\min} = 0.961$ ,  $T_{\max} = 0.980$

5269 measured reflections  
3526 independent reflections  
2959 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.054$   
 $\theta_{\max} = 26.0^\circ$ ,  $\theta_{\min} = 2.0^\circ$   
 $h = -11 \rightarrow 11$   
 $k = -12 \rightarrow 12$   
 $l = -14 \rightarrow 14$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.050$   
 $wR(F^2) = 0.132$   
 $S = 1.04$   
3526 reflections  
246 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0622P)^2 + 0.1786P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3811 (4)	-0.3563 (4)	0.4404 (3)	0.0808 (9)
H1A	0.3092	-0.3111	0.4357	0.121*
H1B	0.3945	-0.3739	0.5185	0.121*
H1C	0.3375	-0.4546	0.3645	0.121*
C2	0.6414 (4)	-0.3214 (4)	0.4384 (3)	0.0776 (8)
H2A	0.5946	-0.4150	0.3577	0.116*
H2B	0.6564	-0.3480	0.5116	0.116*
H2C	0.7433	-0.2490	0.4419	0.116*
C3	0.5317 (3)	-0.1772 (2)	0.3569 (2)	0.0460 (5)
C4	0.4046 (3)	-0.2372 (3)	0.2452 (2)	0.0549 (6)
H4	0.3129	-0.3287	0.2251	0.066*
C5	0.4103 (3)	-0.1632 (3)	0.1607 (2)	0.0543 (6)
H5	0.3217	-0.2059	0.0857	0.065*
C6	0.5421 (3)	-0.0303 (3)	0.1855 (2)	0.0469 (5)
H6	0.5452	0.0133	0.1251	0.056*
C7	0.6746 (2)	0.0422 (2)	0.30273 (19)	0.0386 (4)

C8	0.8158 (2)	0.1871 (2)	0.34276 (19)	0.0407 (5)
C9	0.9371 (3)	0.2558 (3)	0.4599 (2)	0.0498 (5)
H9	1.0277	0.3491	0.4818	0.060*
C10	0.9242 (3)	0.1849 (3)	0.5467 (2)	0.0593 (6)
H10	1.0052	0.2328	0.6279	0.071*
C11	0.7943 (3)	0.0466 (3)	0.5135 (2)	0.0550 (6)
H11	0.7878	0.0016	0.5729	0.066*
C12	0.6684 (2)	-0.0309 (2)	0.39138 (19)	0.0423 (5)
C13	1.0085 (2)	0.1803 (2)	0.12403 (19)	0.0406 (4)
C14	1.0695 (3)	0.1102 (3)	0.1871 (2)	0.0514 (5)
H14	1.0128	0.0581	0.2302	0.062*
C15	1.2188 (3)	0.1174 (3)	0.1866 (2)	0.0616 (6)
H15	1.2622	0.0718	0.2315	0.074*
C16	1.3009 (3)	0.1902 (3)	0.1213 (2)	0.0599 (6)
H16	1.3999	0.1942	0.1221	0.072*
C17	1.2370 (3)	0.2596 (3)	0.0523 (2)	0.0487 (5)
C18	1.3112 (3)	0.3301 (3)	-0.0238 (2)	0.0623 (7)
H18	1.4073	0.3321	-0.0306	0.075*
C19	1.2415 (3)	0.3946 (3)	-0.0863 (3)	0.0643 (7)
H19	1.2891	0.4411	-0.1366	0.077*
C20	1.0978 (3)	0.3904 (3)	-0.0744 (2)	0.0584 (6)
H20	1.0533	0.4377	-0.1164	0.070*
C21	1.0888 (2)	0.2580 (2)	0.05481 (18)	0.0399 (4)
N1	0.5359 (2)	-0.2493 (2)	0.4443 (2)	0.0572 (5)
N2	1.0200 (2)	0.3242 (2)	-0.00774 (17)	0.0479 (4)
O1	0.6994 (2)	0.2943 (2)	0.18886 (19)	0.0656 (5)
O2	0.9897 (2)	0.42788 (18)	0.30555 (16)	0.0607 (4)
O3	0.85533 (16)	0.16883 (17)	0.11789 (13)	0.0442 (4)
S1	0.84187 (7)	0.28769 (6)	0.24117 (5)	0.04560 (19)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.084 (2)	0.085 (2)	0.118 (2)	0.0459 (17)	0.0604 (19)	0.073 (2)
C2	0.081 (2)	0.089 (2)	0.100 (2)	0.0529 (17)	0.0357 (17)	0.0648 (19)
C3	0.0476 (12)	0.0491 (12)	0.0530 (12)	0.0252 (10)	0.0223 (10)	0.0291 (10)
C4	0.0446 (13)	0.0498 (13)	0.0624 (14)	0.0144 (10)	0.0138 (11)	0.0260 (11)
C5	0.0417 (12)	0.0600 (14)	0.0521 (13)	0.0197 (11)	0.0044 (10)	0.0239 (11)
C6	0.0462 (12)	0.0550 (13)	0.0448 (11)	0.0258 (10)	0.0115 (10)	0.0268 (10)
C7	0.0400 (11)	0.0433 (11)	0.0403 (10)	0.0227 (9)	0.0167 (9)	0.0212 (9)
C8	0.0438 (11)	0.0435 (11)	0.0417 (11)	0.0228 (9)	0.0208 (9)	0.0198 (9)
C9	0.0454 (12)	0.0483 (12)	0.0469 (12)	0.0162 (10)	0.0157 (10)	0.0165 (10)
C10	0.0515 (14)	0.0681 (15)	0.0400 (12)	0.0201 (12)	0.0035 (10)	0.0171 (11)
C11	0.0587 (14)	0.0654 (15)	0.0412 (12)	0.0257 (12)	0.0126 (10)	0.0289 (11)
C12	0.0443 (12)	0.0480 (11)	0.0427 (11)	0.0245 (10)	0.0167 (9)	0.0240 (9)
C13	0.0396 (11)	0.0424 (11)	0.0377 (10)	0.0196 (9)	0.0132 (9)	0.0138 (9)
C14	0.0632 (15)	0.0557 (13)	0.0456 (12)	0.0330 (12)	0.0221 (11)	0.0249 (10)
C15	0.0694 (17)	0.0730 (16)	0.0583 (14)	0.0493 (14)	0.0174 (13)	0.0295 (13)

C16	0.0451 (13)	0.0693 (15)	0.0659 (15)	0.0349 (12)	0.0163 (12)	0.0208 (13)
C17	0.0404 (11)	0.0451 (11)	0.0524 (12)	0.0182 (9)	0.0171 (10)	0.0124 (10)
C18	0.0472 (14)	0.0584 (14)	0.0719 (16)	0.0169 (11)	0.0325 (12)	0.0189 (13)
C19	0.0682 (17)	0.0630 (15)	0.0685 (16)	0.0237 (13)	0.0401 (14)	0.0347 (13)
C20	0.0682 (16)	0.0602 (14)	0.0589 (14)	0.0313 (13)	0.0294 (12)	0.0334 (12)
C21	0.0376 (11)	0.0390 (10)	0.0380 (10)	0.0170 (9)	0.0127 (8)	0.0114 (8)
N1	0.0615 (13)	0.0614 (12)	0.0719 (13)	0.0315 (10)	0.0320 (10)	0.0466 (11)
N2	0.0497 (11)	0.0542 (11)	0.0505 (10)	0.0262 (9)	0.0226 (9)	0.0291 (9)
O1	0.0683 (11)	0.0834 (12)	0.0966 (13)	0.0545 (10)	0.0485 (10)	0.0642 (11)
O2	0.0660 (11)	0.0423 (8)	0.0713 (10)	0.0186 (8)	0.0341 (9)	0.0231 (8)
O3	0.0389 (8)	0.0519 (8)	0.0446 (8)	0.0201 (7)	0.0182 (6)	0.0227 (7)
S1	0.0499 (3)	0.0456 (3)	0.0582 (3)	0.0270 (3)	0.0298 (3)	0.0300 (3)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

C1—N1	1.454 (3)	C10—H10	0.9300
C1—H1A	0.9600	C11—C12	1.412 (3)
C1—H1B	0.9600	C11—H11	0.9300
C1—H1C	0.9600	C13—C14	1.358 (3)
C2—N1	1.448 (3)	C13—O3	1.411 (2)
C2—H2A	0.9600	C13—C21	1.415 (3)
C2—H2B	0.9600	C14—C15	1.403 (3)
C2—H2C	0.9600	C14—H14	0.9300
C3—C4	1.364 (3)	C15—C16	1.360 (4)
C3—N1	1.417 (3)	C15—H15	0.9300
C3—C12	1.433 (3)	C16—C17	1.410 (3)
C4—C5	1.396 (3)	C16—H16	0.9300
C4—H4	0.9300	C17—C21	1.416 (3)
C5—C6	1.356 (3)	C17—C18	1.418 (3)
C5—H5	0.9300	C18—C19	1.353 (4)
C6—C7	1.413 (3)	C18—H18	0.9300
C6—H6	0.9300	C19—C20	1.391 (4)
C7—C12	1.430 (3)	C19—H19	0.9300
C7—C8	1.434 (3)	C20—N2	1.320 (3)
C8—C9	1.362 (3)	C20—H20	0.9300
C8—S1	1.766 (2)	C21—N2	1.363 (3)
C9—C10	1.396 (3)	O1—S1	1.4188 (17)
C9—H9	0.9300	O2—S1	1.4183 (17)
C10—C11	1.356 (3)	O3—S1	1.5933 (15)
N1—C1—H1A	109.5	C11—C12—C3	121.60 (18)
N1—C1—H1B	109.5	C7—C12—C3	119.32 (18)
H1A—C1—H1B	109.5	C14—C13—O3	120.52 (19)
N1—C1—H1C	109.5	C14—C13—C21	122.3 (2)
H1A—C1—H1C	109.5	O3—C13—C21	117.06 (17)
H1B—C1—H1C	109.5	C13—C14—C15	119.3 (2)
N1—C2—H2A	109.5	C13—C14—H14	120.3
N1—C2—H2B	109.5	C15—C14—H14	120.3

H2A—C2—H2B	109.5	C16—C15—C14	120.8 (2)
N1—C2—H2C	109.5	C16—C15—H15	119.6
H2A—C2—H2C	109.5	C14—C15—H15	119.6
H2B—C2—H2C	109.5	C15—C16—C17	120.5 (2)
C4—C3—N1	123.8 (2)	C15—C16—H16	119.7
C4—C3—C12	119.17 (18)	C17—C16—H16	119.7
N1—C3—C12	117.06 (19)	C16—C17—C21	119.7 (2)
C3—C4—C5	120.9 (2)	C16—C17—C18	123.8 (2)
C3—C4—H4	119.5	C21—C17—C18	116.5 (2)
C5—C4—H4	119.5	C19—C18—C17	119.8 (2)
C6—C5—C4	121.6 (2)	C19—C18—H18	120.1
C6—C5—H5	119.2	C17—C18—H18	120.1
C4—C5—H5	119.2	C18—C19—C20	119.0 (2)
C5—C6—C7	120.30 (19)	C18—C19—H19	120.5
C5—C6—H6	119.8	C20—C19—H19	120.5
C7—C6—H6	119.8	N2—C20—C19	124.7 (2)
C6—C7—C12	118.50 (18)	N2—C20—H20	117.6
C6—C7—C8	125.16 (18)	C19—C20—H20	117.6
C12—C7—C8	116.32 (18)	N2—C21—C13	119.30 (18)
C9—C8—C7	122.78 (18)	N2—C21—C17	123.32 (19)
C9—C8—S1	116.00 (16)	C13—C21—C17	117.37 (19)
C7—C8—S1	121.22 (15)	C3—N1—C2	113.78 (19)
C8—C9—C10	119.4 (2)	C3—N1—C1	116.1 (2)
C8—C9—H9	120.3	C2—N1—C1	110.5 (2)
C10—C9—H9	120.3	C20—N2—C21	116.59 (19)
C11—C10—C9	120.4 (2)	C13—O3—S1	117.87 (12)
C11—C10—H10	119.8	O2—S1—O1	119.57 (11)
C9—C10—H10	119.8	O2—S1—O3	108.77 (8)
C10—C11—C12	121.9 (2)	O1—S1—O3	104.37 (10)
C10—C11—H11	119.0	O2—S1—C8	109.32 (10)
C12—C11—H11	119.0	O1—S1—C8	110.89 (10)
C11—C12—C7	119.01 (19)	O3—S1—C8	102.41 (8)
N1—C3—C4—C5	-177.7 (2)	C16—C17—C18—C19	-179.7 (2)
C12—C3—C4—C5	3.9 (3)	C21—C17—C18—C19	1.6 (3)
C3—C4—C5—C6	0.6 (4)	C17—C18—C19—C20	0.1 (4)
C4—C5—C6—C7	-3.7 (4)	C18—C19—C20—N2	-1.5 (4)
C5—C6—C7—C12	2.2 (3)	C14—C13—C21—N2	179.21 (19)
C5—C6—C7—C8	-176.3 (2)	O3—C13—C21—N2	3.3 (3)
C6—C7—C8—C9	176.8 (2)	C14—C13—C21—C17	0.5 (3)
C12—C7—C8—C9	-1.7 (3)	O3—C13—C21—C17	-175.38 (17)
C6—C7—C8—S1	-3.1 (3)	C16—C17—C21—N2	179.12 (19)
C12—C7—C8—S1	178.40 (14)	C18—C17—C21—N2	-2.1 (3)
C7—C8—C9—C10	-1.0 (3)	C16—C17—C21—C13	-2.2 (3)
S1—C8—C9—C10	178.88 (17)	C18—C17—C21—C13	176.51 (18)
C8—C9—C10—C11	1.8 (4)	C4—C3—N1—C2	107.4 (3)
C9—C10—C11—C12	0.2 (4)	C12—C3—N1—C2	-74.3 (3)
C10—C11—C12—C7	-3.0 (3)	C4—C3—N1—C1	-22.5 (3)

C10—C11—C12—C3	179.9 (2)	C12—C3—N1—C1	155.8 (2)
C6—C7—C12—C11	-175.00 (19)	C19—C20—N2—C21	1.0 (4)
C8—C7—C12—C11	3.6 (3)	C13—C21—N2—C20	-177.77 (19)
C6—C7—C12—C3	2.2 (3)	C17—C21—N2—C20	0.9 (3)
C8—C7—C12—C3	-179.18 (17)	C14—C13—O3—S1	82.5 (2)
C4—C3—C12—C11	171.9 (2)	C21—C13—O3—S1	-101.55 (17)
N1—C3—C12—C11	-6.6 (3)	C13—O3—S1—O2	28.05 (16)
C4—C3—C12—C7	-5.2 (3)	C13—O3—S1—O1	156.75 (14)
N1—C3—C12—C7	176.32 (17)	C13—O3—S1—C8	-87.57 (15)
O3—C13—C14—C15	177.12 (19)	C9—C8—S1—O2	-0.92 (19)
C21—C13—C14—C15	1.4 (3)	C7—C8—S1—O2	178.98 (15)
C13—C14—C15—C16	-1.6 (4)	C9—C8—S1—O1	-134.83 (17)
C14—C15—C16—C17	-0.2 (4)	C7—C8—S1—O1	45.06 (19)
C15—C16—C17—C21	2.1 (3)	C9—C8—S1—O3	114.31 (17)
C15—C16—C17—C18	-176.5 (2)	C7—C8—S1—O3	-65.80 (17)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
C16—H16···O1 <sup>i</sup>	0.93	2.52	3.411 (3)	160

Symmetry code: (i)  $x+1, y, z$ .